



12a₀₈₈₂ (K12a₀₈₈₂)



A knot diagram¹

Linearized knot diagram



Solving Sequence

$$8,12 \xrightarrow{c_8} 9 \xrightarrow{c_{12}} 1,4 \xrightarrow{c_1} 2 \xrightarrow{c_9} 10 \xrightarrow{c_4} 5 \xrightarrow{c_{11}} 11 \xrightarrow{c_7} 7 \xrightarrow{c_3} 3 \xrightarrow{c_6} 6 \rightsquigarrow c_2, c_5, c_{10}$$

Ideals for irreducible components² of X_{par}

$$I_1^u = \langle 4u^{74} - 16u^{73} + \dots + 4b - 5, -u^{74} + 8u^{73} + \dots + 4a + 13, u^{75} - 4u^{74} + \dots - 6u + 1 \rangle$$

$$I_2^u = \langle u^2a + b + a, u^2a + a^2 + u^2 + a + u + 1, u^3 + u^2 + 2u + 1 \rangle$$

$$I_3^u = \langle -u^3 + b - u, a + u, u^7 + 3u^5 + 2u^3 - u - 1 \rangle$$

$$I_4^u = \langle u^2 + b + u + 1, a + u, u^3 + u^2 + 2u + 1 \rangle$$

* 4 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 91 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/maths/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$\langle 4u^{74} - 16u^{73} + \dots + 4b - 5, -u^{74} + 8u^{73} + \dots + 4a + 13, u^{75} - 4u^{74} + \dots - 6u + 1 \rangle$$

I. $I_1^u =$

(i) Arc colorings

$$a_8 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ -u^2 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} u \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} \frac{1}{4}u^{74} - 2u^{73} + \dots + \frac{65}{4}u - \frac{13}{4} \\ -u^{74} + 4u^{73} + \dots - 4u + \frac{5}{4} \end{pmatrix}$$

$$a_2 = \begin{pmatrix} \frac{1}{4}u^{74} - \frac{3}{4}u^{73} + \dots + 7u + \frac{3}{4} \\ -\frac{1}{4}u^{74} + u^{73} + \dots - \frac{9}{4}u + \frac{1}{2} \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -u^4 - u^2 + 1 \\ -u^4 - 2u^2 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} \frac{15}{4}u^{74} - 10u^{73} + \dots + \frac{23}{4}u - \frac{3}{4} \\ 2u^{74} - u^{73} + \dots - 8u + \frac{7}{4} \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -u \\ u^3 + u \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -u^4 - u^2 + 1 \\ u^6 + 2u^4 + u^2 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} \frac{13}{4}u^{74} - \frac{55}{4}u^{73} + \dots + \frac{53}{2}u - \frac{17}{4} \\ u^{74} - \frac{15}{4}u^{73} + \dots - \frac{11}{4}u + \frac{5}{4} \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -u^{74} + \frac{19}{4}u^{73} + \dots - \frac{7}{4}u + \frac{3}{4} \\ -\frac{3}{4}u^{74} + 3u^{73} + \dots + \frac{9}{4}u - \frac{1}{2} \end{pmatrix}$$

(ii) Obstruction class = -1

$$(iii) \text{ Cusp Shapes} = -\frac{9}{2}u^{74} + \frac{19}{2}u^{73} + \dots - \frac{39}{2}u - \frac{9}{4}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{75} - 14u^{74} + \dots + 204u + 801$
c_2, c_5, c_6	$u^{75} - 4u^{74} + \dots + 10u - 1$
c_3	$u^{75} + 4u^{74} + \dots + 2274u - 153$
c_4, c_{10}	$u^{75} + 6u^{74} + \dots - 3072u - 512$
c_7, c_9	$u^{75} - 4u^{74} + \dots - 750u - 153$
c_8, c_{11}, c_{12}	$u^{75} + 4u^{74} + \dots - 6u - 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{75} + 42y^{74} + \dots + 136940526y - 641601$
c_2, c_5, c_6	$y^{75} + 70y^{74} + \dots + 34y - 1$
c_3	$y^{75} + 14y^{74} + \dots + 593622y - 23409$
c_4, c_{10}	$y^{75} - 42y^{74} + \dots + 3276800y - 262144$
c_7, c_9	$y^{75} - 58y^{74} + \dots - 431082y - 23409$
c_8, c_{11}, c_{12}	$y^{75} + 62y^{74} + \dots - 14y - 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.129331 + 1.117110I$ $a = 0.29237 - 1.85419I$ $b = 0.11903 - 1.46059I$	$1.67101 - 2.28476I$	0
$u = -0.129331 - 1.117110I$ $a = 0.29237 + 1.85419I$ $b = 0.11903 + 1.46059I$	$1.67101 + 2.28476I$	0
$u = 0.864283 + 0.073603I$ $a = -0.630215 - 0.318792I$ $b = -0.46463 - 1.50057I$	$14.07080 + 0.97614I$	$11.89444 - 0.69772I$
$u = 0.864283 - 0.073603I$ $a = -0.630215 + 0.318792I$ $b = -0.46463 + 1.50057I$	$14.07080 - 0.97614I$	$11.89444 + 0.69772I$
$u = 0.851962 + 0.130511I$ $a = 0.189706 + 0.407751I$ $b = -0.20536 + 2.41576I$	$12.0915 + 11.1738I$	$9.92293 - 6.69276I$
$u = 0.851962 - 0.130511I$ $a = 0.189706 - 0.407751I$ $b = -0.20536 - 2.41576I$	$12.0915 - 11.1738I$	$9.92293 + 6.69276I$
$u = 0.838542 + 0.118303I$ $a = -0.263667 - 0.275157I$ $b = 0.30731 - 2.10324I$	$6.28260 + 7.50607I$	$6.31085 - 6.58507I$
$u = 0.838542 - 0.118303I$ $a = -0.263667 + 0.275157I$ $b = 0.30731 + 2.10324I$	$6.28260 - 7.50607I$	$6.31085 + 6.58507I$
$u = 0.834761 + 0.091281I$ $a = 0.443254 + 0.200964I$ $b = -0.17545 + 1.64029I$	$7.24740 + 3.22206I$	$8.45543 - 0.86065I$
$u = 0.834761 - 0.091281I$ $a = 0.443254 - 0.200964I$ $b = -0.17545 - 1.64029I$	$7.24740 - 3.22206I$	$8.45543 + 0.86065I$

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.806243 + 0.042985I$ $a = 0.254905 + 0.770452I$ $b = 0.64512 + 2.35548I$	$8.28866 - 4.61277I$	$9.57496 + 3.37665I$
$u = -0.806243 - 0.042985I$ $a = 0.254905 - 0.770452I$ $b = 0.64512 - 2.35548I$	$8.28866 + 4.61277I$	$9.57496 - 3.37665I$
$u = 0.417667 + 1.122760I$ $a = 2.24416 - 1.33147I$ $b = 0.82696 - 1.40165I$	$9.05288 - 6.61172I$	0
$u = 0.417667 - 1.122760I$ $a = 2.24416 + 1.33147I$ $b = 0.82696 + 1.40165I$	$9.05288 + 6.61172I$	0
$u = 0.394262 + 1.136570I$ $a = -1.88247 + 1.23446I$ $b = -0.590542 + 1.186780I$	$3.16758 - 3.06037I$	0
$u = 0.394262 - 1.136570I$ $a = -1.88247 - 1.23446I$ $b = -0.590542 - 1.186780I$	$3.16758 + 3.06037I$	0
$u = -0.767081 + 0.032918I$ $a = -0.147823 - 0.586406I$ $b = -0.41161 - 1.84908I$	$2.54564 - 1.71234I$	$5.68970 + 3.98160I$
$u = -0.767081 - 0.032918I$ $a = -0.147823 + 0.586406I$ $b = -0.41161 + 1.84908I$	$2.54564 + 1.71234I$	$5.68970 - 3.98160I$
$u = 0.387354 + 1.172960I$ $a = 1.54778 - 0.81015I$ $b = 0.496497 - 0.735271I$	$3.93329 + 1.17961I$	0
$u = 0.387354 - 1.172960I$ $a = 1.54778 + 0.81015I$ $b = 0.496497 + 0.735271I$	$3.93329 - 1.17961I$	0

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.761899 + 0.044601I$ $a = 0.670042 - 0.120108I$ $b = -1.140900 + 0.673722I$	$5.34357 + 3.86193I$	$9.73760 - 4.61433I$
$u = 0.761899 - 0.044601I$ $a = 0.670042 + 0.120108I$ $b = -1.140900 - 0.673722I$	$5.34357 - 3.86193I$	$9.73760 + 4.61433I$
$u = -0.502648 + 0.555980I$ $a = 1.35292 - 0.76273I$ $b = 0.424196 + 0.870841I$	$6.99841 - 6.56505I$	$7.94758 + 6.80228I$
$u = -0.502648 - 0.555980I$ $a = 1.35292 + 0.76273I$ $b = 0.424196 - 0.870841I$	$6.99841 + 6.56505I$	$7.94758 - 6.80228I$
$u = 0.417996 + 1.195510I$ $a = -1.81445 + 0.17291I$ $b = -0.943590 + 0.346371I$	$10.62020 + 3.61647I$	0
$u = 0.417996 - 1.195510I$ $a = -1.81445 - 0.17291I$ $b = -0.943590 - 0.346371I$	$10.62020 - 3.61647I$	0
$u = -0.586451 + 0.420780I$ $a = -0.745096 - 0.020246I$ $b = 0.568271 - 1.084970I$	$7.43119 + 2.66733I$	$9.42694 + 0.01688I$
$u = -0.586451 - 0.420780I$ $a = -0.745096 + 0.020246I$ $b = 0.568271 + 1.084970I$	$7.43119 - 2.66733I$	$9.42694 - 0.01688I$
$u = -0.352653 + 1.230150I$ $a = -2.08694 - 2.25380I$ $b = -1.15302 - 2.28427I$	$4.63286 + 0.43633I$	0
$u = -0.352653 - 1.230150I$ $a = -2.08694 + 2.25380I$ $b = -1.15302 + 2.28427I$	$4.63286 - 0.43633I$	0

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.316269 + 1.249700I$ $a = 1.82642 + 1.56645I$ $b = 1.16268 + 1.70863I$	$-1.20445 - 2.19229I$	0
$u = -0.316269 - 1.249700I$ $a = 1.82642 - 1.56645I$ $b = 1.16268 - 1.70863I$	$-1.20445 + 2.19229I$	0
$u = 0.053632 + 1.289800I$ $a = -1.55559 + 0.97168I$ $b = -0.72624 + 1.73649I$	$-0.99866 + 4.23513I$	0
$u = 0.053632 - 1.289800I$ $a = -1.55559 - 0.97168I$ $b = -0.72624 - 1.73649I$	$-0.99866 - 4.23513I$	0
$u = 0.016620 + 1.304390I$ $a = 1.194330 - 0.735726I$ $b = 0.69581 - 1.49341I$	$-5.82265 + 0.81095I$	0
$u = 0.016620 - 1.304390I$ $a = 1.194330 + 0.735726I$ $b = 0.69581 + 1.49341I$	$-5.82265 - 0.81095I$	0
$u = -0.667774 + 0.163365I$ $a = 0.235125 + 0.105097I$ $b = 1.132700 + 0.443466I$	$4.25302 - 0.78723I$	$8.08019 + 0.69488I$
$u = -0.667774 - 0.163365I$ $a = 0.235125 - 0.105097I$ $b = 1.132700 - 0.443466I$	$4.25302 + 0.78723I$	$8.08019 - 0.69488I$
$u = 0.321229 + 1.276340I$ $a = 0.975526 + 0.758046I$ $b = 1.38917 - 0.31193I$	$-2.32532 + 3.88876I$	0
$u = 0.321229 - 1.276340I$ $a = 0.975526 - 0.758046I$ $b = 1.38917 + 0.31193I$	$-2.32532 - 3.88876I$	0

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.099287 + 1.314930I$ $a = -0.318472 + 0.820675I$ $b = -0.439087 + 1.148180I$	$-3.46404 - 2.05434I$	0
$u = -0.099287 - 1.314930I$ $a = -0.318472 - 0.820675I$ $b = -0.439087 - 1.148180I$	$-3.46404 + 2.05434I$	0
$u = -0.450416 + 0.509884I$ $a = -0.960127 + 0.859309I$ $b = -0.166013 - 0.591858I$	$1.28106 - 3.47044I$	$3.88375 + 7.68592I$
$u = -0.450416 - 0.509884I$ $a = -0.960127 - 0.859309I$ $b = -0.166013 + 0.591858I$	$1.28106 + 3.47044I$	$3.88375 - 7.68592I$
$u = -0.332166 + 1.293110I$ $a = -2.54722 - 1.00893I$ $b = -1.85697 - 1.49469I$	$-1.59371 - 5.68239I$	0
$u = -0.332166 - 1.293110I$ $a = -2.54722 + 1.00893I$ $b = -1.85697 + 1.49469I$	$-1.59371 + 5.68239I$	0
$u = 0.331305 + 1.302290I$ $a = -1.56104 - 0.14984I$ $b = -1.55504 + 0.97937I$	$1.12844 + 7.81474I$	0
$u = 0.331305 - 1.302290I$ $a = -1.56104 + 0.14984I$ $b = -1.55504 - 0.97937I$	$1.12844 - 7.81474I$	0
$u = -0.355420 + 1.298670I$ $a = 3.07198 + 1.18787I$ $b = 2.20157 + 1.78185I$	$4.10044 - 8.79609I$	0
$u = -0.355420 - 1.298670I$ $a = 3.07198 - 1.18787I$ $b = 2.20157 - 1.78185I$	$4.10044 + 8.79609I$	0

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.036427 + 1.346260I$ $a = -0.607791 + 0.422715I$ $b = -0.629673 + 1.192930I$	$-3.35604 - 2.35859I$	0
$u = -0.036427 - 1.346260I$ $a = -0.607791 - 0.422715I$ $b = -0.629673 - 1.192930I$	$-3.35604 + 2.35859I$	0
$u = -0.204057 + 1.350880I$ $a = -0.508818 + 0.849448I$ $b = -0.815090 + 0.663922I$	$-3.59967 - 2.38098I$	0
$u = -0.204057 - 1.350880I$ $a = -0.508818 - 0.849448I$ $b = -0.815090 - 0.663922I$	$-3.59967 + 2.38098I$	0
$u = -0.269785 + 1.347060I$ $a = 1.64477 - 0.56880I$ $b = 1.62647 + 0.01247I$	$-0.50518 - 4.19710I$	0
$u = -0.269785 - 1.347060I$ $a = 1.64477 + 0.56880I$ $b = 1.62647 - 0.01247I$	$-0.50518 + 4.19710I$	0
$u = 0.391046 + 1.322010I$ $a = 1.06116 - 1.80811I$ $b = 0.48619 - 2.18356I$	$9.70536 + 5.47983I$	0
$u = 0.391046 - 1.322010I$ $a = 1.06116 + 1.80811I$ $b = 0.48619 + 2.18356I$	$9.70536 - 5.47983I$	0
$u = 0.368532 + 1.330510I$ $a = -1.73222 + 1.34594I$ $b = -1.12286 + 2.08477I$	$2.78873 + 7.54981I$	0
$u = 0.368532 - 1.330510I$ $a = -1.73222 - 1.34594I$ $b = -1.12286 - 2.08477I$	$2.78873 - 7.54981I$	0

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.368054 + 1.347150I$ $a = 2.22544 - 1.63294I$ $b = 1.36977 - 2.45177I$	$1.67578 + 11.84830I$	0
$u = 0.368054 - 1.347150I$ $a = 2.22544 + 1.63294I$ $b = 1.36977 + 2.45177I$	$1.67578 - 11.84830I$	0
$u = -0.103421 + 1.398990I$ $a = -0.156439 - 0.588353I$ $b = 0.497683 - 1.119190I$	$-4.75437 - 5.19853I$	0
$u = -0.103421 - 1.398990I$ $a = -0.156439 + 0.588353I$ $b = 0.497683 + 1.119190I$	$-4.75437 + 5.19853I$	0
$u = 0.373793 + 1.356110I$ $a = -2.39427 + 1.96966I$ $b = -1.37738 + 2.74063I$	$7.4137 + 15.5822I$	0
$u = 0.373793 - 1.356110I$ $a = -2.39427 - 1.96966I$ $b = -1.37738 - 2.74063I$	$7.4137 - 15.5822I$	0
$u = -0.11241 + 1.42300I$ $a = 0.455015 + 0.662998I$ $b = -0.434609 + 1.177910I$	$0.66003 - 8.48529I$	0
$u = -0.11241 - 1.42300I$ $a = 0.455015 - 0.662998I$ $b = -0.434609 - 1.177910I$	$0.66003 + 8.48529I$	0
$u = -0.484381$ $a = -0.253063$ $b = -0.395089$	0.828246	12.7290
$u = -0.040668 + 0.478942I$ $a = 0.06428 - 2.03562I$ $b = -0.380831 - 0.239183I$	$2.10140 - 1.97255I$	$1.62971 + 4.07824I$

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.040668 - 0.478942I$		
$a = 0.06428 + 2.03562I$	$2.10140 + 1.97255I$	$1.62971 - 4.07824I$
$b = -0.380831 + 0.239183I$		
$u = 0.268652 + 0.175360I$		
$a = -0.22486 - 2.74712I$	$3.41208 + 3.23701I$	$-0.51319 - 4.60923I$
$b = -0.722155 + 0.751154I$		
$u = 0.268652 - 0.175360I$		
$a = -0.22486 + 2.74712I$	$3.41208 - 3.23701I$	$-0.51319 + 4.60923I$
$b = -0.722155 - 0.751154I$		
$u = 0.113105 + 0.236302I$		
$a = 0.51485 + 2.54920I$	$-1.187190 + 0.455593I$	$-6.24430 - 1.84121I$
$b = 0.559155 - 0.256506I$		
$u = 0.113105 - 0.236302I$		
$a = 0.51485 - 2.54920I$	$-1.187190 - 0.455593I$	$-6.24430 + 1.84121I$
$b = 0.559155 + 0.256506I$		

$$\text{II. } I_2^u = \langle u^2a + b + a, u^2a + a^2 + u^2 + a + u + 1, u^3 + u^2 + 2u + 1 \rangle$$

(i) Arc colorings

$$a_8 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ -u^2 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} u \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} a \\ -u^2a - a \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -u^2a - au - 2a + u - 1 \\ u^2a + u^2 + 2a + u + 1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -u \\ -u^2 - u - 1 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} a \\ -u^2a - a \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -u \\ -u^2 - u - 1 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -u \\ -u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -u^2a - au \\ -2u^2a - au - 2a \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -au - u - 1 \\ u^2a + 2u^2 + 2a + 3 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = $-3u^2a + 5u^2 - a + 5u + 8$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_3	$(u^3 + u^2 - 1)^2$
c_2, c_{11}, c_{12}	$(u^3 - u^2 + 2u - 1)^2$
c_4, c_{10}	u^6
c_5, c_6, c_8	$(u^3 + u^2 + 2u + 1)^2$
c_7, c_9	$(u^3 - u^2 + 1)^2$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_3, c_7 c_9	$(y^3 - y^2 + 2y - 1)^2$
c_2, c_5, c_6 c_8, c_{11}, c_{12}	$(y^3 + 3y^2 + 2y - 1)^2$
c_4, c_{10}	y^6

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.215080 + 1.307140I$ $a = -0.662359 + 0.562280I$ $b = -0.754878$	$-4.13758 - 2.82812I$	$-4.97655 + 4.84887I$
$u = -0.215080 + 1.307140I$ $a = 1.32472$ $b = 0.877439 + 0.744862I$	$-5.65624I$	$3.89456 + 5.95889I$
$u = -0.215080 - 1.307140I$ $a = -0.662359 - 0.562280I$ $b = -0.754878$	$-4.13758 + 2.82812I$	$-4.97655 - 4.84887I$
$u = -0.215080 - 1.307140I$ $a = 1.32472$ $b = 0.877439 - 0.744862I$	$5.65624I$	$3.89456 - 5.95889I$
$u = -0.569840$ $a = -0.662359 + 0.562280I$ $b = 0.877439 - 0.744862I$	$4.13758 + 2.82812I$	$8.08199 - 1.11003I$
$u = -0.569840$ $a = -0.662359 - 0.562280I$ $b = 0.877439 + 0.744862I$	$4.13758 - 2.82812I$	$8.08199 + 1.11003I$

$$\text{III. } I_3^u = \langle -u^3 + b - u, a + u, u^7 + 3u^5 + 2u^3 - u - 1 \rangle$$

(i) Arc colorings

$$a_8 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ -u^2 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} u \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -u \\ u^3 + u \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -u^5 - 2u^3 + u \\ 2u + 1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -u^4 - u^2 + 1 \\ -u^4 - 2u^2 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} u^4 + u^2 - u - 1 \\ u^4 + u^3 + 2u^2 + u \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -u \\ u^3 + u \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -u^4 - u^2 + 1 \\ u^6 + 2u^4 + u^2 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} u^5 + 2u^3 - u - 1 \\ u^5 + 3u^3 + u^2 + u \end{pmatrix}$$

$$a_6 = \begin{pmatrix} u^6 + u^4 - 2u^2 + 1 \\ u^6 + 2u^4 - u^2 - u \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes = 6

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^7 - 2u^6 + u^5 + 2u^4 - 4u^3 + 6u^2 - 3u + 3$
c_2, c_5, c_6 c_8, c_{11}, c_{12}	$u^7 + 3u^5 + 2u^3 - u + 1$
c_3, c_7, c_9	$u^7 - u^5 - 2u^4 + 2u^3 + 2u^2 - 3u + 2$
c_4, c_{10}	$(u - 1)^7$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^7 - 2y^6 + y^5 + 6y^4 - 2y^3 - 24y^2 - 27y - 9$
c_2, c_5, c_6 c_8, c_{11}, c_{12}	$y^7 + 6y^6 + 13y^5 + 10y^4 - 2y^3 - 4y^2 + y - 1$
c_3, c_7, c_9	$y^7 - 2y^6 + 5y^5 - 14y^4 + 18y^3 - 8y^2 + y - 4$
c_4, c_{10}	$(y - 1)^7$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.757137$ $a = -0.757137$ $b = 1.19117$	1.64493	6.00000
$u = 0.311114 + 1.246820I$ $a = -0.311114 - 1.246820I$ $b = -1.109710 - 0.329390I$	1.64493	6.00000
$u = 0.311114 - 1.246820I$ $a = -0.311114 + 1.246820I$ $b = -1.109710 + 0.329390I$	1.64493	6.00000
$u = -0.501027 + 0.385135I$ $a = 0.501027 - 0.385135I$ $b = -0.403848 + 0.618048I$	1.64493	6.00000
$u = -0.501027 - 0.385135I$ $a = 0.501027 + 0.385135I$ $b = -0.403848 - 0.618048I$	1.64493	6.00000
$u = -0.18866 + 1.40255I$ $a = 0.18866 - 1.40255I$ $b = 0.91797 - 1.20672I$	1.64493	6.00000
$u = -0.18866 - 1.40255I$ $a = 0.18866 + 1.40255I$ $b = 0.91797 + 1.20672I$	1.64493	6.00000

$$\text{IV. } I_4^u = \langle u^2 + b + u + 1, a + u, u^3 + u^2 + 2u + 1 \rangle$$

(i) Arc colorings

$$a_8 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ -u^2 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} u \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -u \\ -u^2 - u - 1 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 2u + 1 \\ 2u \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -u \\ -u^2 - u - 1 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -u \\ -u^2 - u - 1 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -u \\ -u^2 - u - 1 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -u \\ -u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -2u - 1 \\ -u^2 - 2u - 2 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -2u^2 - 2u \\ -2u^2 - u \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = 0

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_3	$u^3 + u^2 - 1$
c_2, c_{11}, c_{12}	$u^3 - u^2 + 2u - 1$
c_4, c_{10}	u^3
c_5, c_6, c_8	$u^3 + u^2 + 2u + 1$
c_7, c_9	$u^3 - u^2 + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_3, c_7 c_9	$y^3 - y^2 + 2y - 1$
c_2, c_5, c_6 c_8, c_{11}, c_{12}	$y^3 + 3y^2 + 2y - 1$
c_4, c_{10}	y^3

(vi) Complex Volumes and Cusp Shapes

Solutions to I_4^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.215080 + 1.307140I$ $a = 0.215080 - 1.307140I$ $b = 0.877439 - 0.744862I$	0	0
$u = -0.215080 - 1.307140I$ $a = 0.215080 + 1.307140I$ $b = 0.877439 + 0.744862I$	0	0
$u = -0.569840$ $a = 0.569840$ $b = -0.754878$	0	0

V. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1	$(u^3 + u^2 - 1)^3(u^7 - 2u^6 + u^5 + 2u^4 - 4u^3 + 6u^2 - 3u + 3)$ $\cdot (u^{75} - 14u^{74} + \dots + 204u + 801)$
c_2	$((u^3 - u^2 + 2u - 1)^3)(u^7 + 3u^5 + 2u^3 - u + 1)(u^{75} - 4u^{74} + \dots + 10u - 1)$
c_3	$(u^3 + u^2 - 1)^3(u^7 - u^5 - 2u^4 + 2u^3 + 2u^2 - 3u + 2)$ $\cdot (u^{75} + 4u^{74} + \dots + 2274u - 153)$
c_4, c_{10}	$u^9(u - 1)^7(u^{75} + 6u^{74} + \dots - 3072u - 512)$
c_5, c_6	$((u^3 + u^2 + 2u + 1)^3)(u^7 + 3u^5 + 2u^3 - u + 1)(u^{75} - 4u^{74} + \dots + 10u - 1)$
c_7, c_9	$(u^3 - u^2 + 1)^3(u^7 - u^5 - 2u^4 + 2u^3 + 2u^2 - 3u + 2)$ $\cdot (u^{75} - 4u^{74} + \dots - 750u - 153)$
c_8	$((u^3 + u^2 + 2u + 1)^3)(u^7 + 3u^5 + 2u^3 - u + 1)(u^{75} + 4u^{74} + \dots - 6u - 1)$
c_{11}, c_{12}	$((u^3 - u^2 + 2u - 1)^3)(u^7 + 3u^5 + 2u^3 - u + 1)(u^{75} + 4u^{74} + \dots - 6u - 1)$

VI. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1	$(y^3 - y^2 + 2y - 1)^3(y^7 - 2y^6 + y^5 + 6y^4 - 2y^3 - 24y^2 - 27y - 9)$ $\cdot (y^{75} + 42y^{74} + \dots + 136940526y - 641601)$
c_2, c_5, c_6	$(y^3 + 3y^2 + 2y - 1)^3(y^7 + 6y^6 + 13y^5 + 10y^4 - 2y^3 - 4y^2 + y - 1)$ $\cdot (y^{75} + 70y^{74} + \dots + 34y - 1)$
c_3	$(y^3 - y^2 + 2y - 1)^3(y^7 - 2y^6 + 5y^5 - 14y^4 + 18y^3 - 8y^2 + y - 4)$ $\cdot (y^{75} + 14y^{74} + \dots + 593622y - 23409)$
c_4, c_{10}	$y^9(y - 1)^7(y^{75} - 42y^{74} + \dots + 3276800y - 262144)$
c_7, c_9	$(y^3 - y^2 + 2y - 1)^3(y^7 - 2y^6 + 5y^5 - 14y^4 + 18y^3 - 8y^2 + y - 4)$ $\cdot (y^{75} - 58y^{74} + \dots - 431082y - 23409)$
c_8, c_{11}, c_{12}	$(y^3 + 3y^2 + 2y - 1)^3(y^7 + 6y^6 + 13y^5 + 10y^4 - 2y^3 - 4y^2 + y - 1)$ $\cdot (y^{75} + 62y^{74} + \dots - 14y - 1)$