


$12n_{0140}$ ($K12n_{0140}$)



Ideals for irreducible components² of X_{par}

$$\begin{aligned}
 I_1^u &= \langle -219034042585u^{29} + 477658347892u^{28} + \dots + 4962190966784b + 2219335787081, \\
 &\quad -1283291111693u^{29} + 4041034486657u^{28} + \dots + 2481095483392a + 21380565133380, \\
 &\quad u^{30} - 3u^{29} + \dots - 14u - 1 \rangle \\
 I_2^u &= \langle 9.19271 \times 10^{50}u^{41} + 6.53340 \times 10^{51}u^{40} + \dots + 1.68743 \times 10^{52}b + 4.76895 \times 10^{52}, \\
 &\quad -4.05666 \times 10^{51}u^{41} - 1.46559 \times 10^{52}u^{40} + \dots + 1.18120 \times 10^{53}a - 9.05537 \times 10^{53}, \\
 &\quad u^{42} + 8u^{41} + \dots + 406u + 49 \rangle \\
 I_3^u &= \langle b, -u^3 + u^2 + 4a + 2u + 3, u^4 + u^2 - u + 1 \rangle \\
 I_4^u &= \langle 8a^2 + b + 18a + 4, 8a^3 + 20a^2 + 8a + 1, u - 1 \rangle \\
 I_5^u &= \langle b, -u^3 + a - u - 1, u^6 + u^5 + 2u^4 + 2u^3 + 2u^2 + 2u + 1 \rangle \\
 I_6^u &= \langle au + 4b + a + u - 5, a^2 + 4au - 2a + 6u - 3, u^2 + 1 \rangle
 \end{aligned}$$

* 6 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 89 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/math/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$\text{I. } I_1^u = \langle -2.19 \times 10^{11}u^{29} + 4.78 \times 10^{11}u^{28} + \dots + 4.96 \times 10^{12}b + 2.22 \times 10^{12}, -1.28 \times 10^{12}u^{29} + 4.04 \times 10^{12}u^{28} + \dots + 2.48 \times 10^{12}a + 2.14 \times 10^{13}, u^{30} - 3u^{29} + \dots - 14u - 1 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_8 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_9 &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_3 &= \begin{pmatrix} 0.517228u^{29} - 1.62873u^{28} + \dots + 26.8709u - 8.61739 \\ 0.0441406u^{29} - 0.0962596u^{28} + \dots + 1.16893u - 0.447249 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} -u \\ u \end{pmatrix} \\ a_7 &= \begin{pmatrix} 0.146114u^{29} - 0.558009u^{28} + \dots + 11.5186u - 3.67862 \\ 0.114897u^{29} - 0.286853u^{28} + \dots - 0.242495u - 0.299101 \end{pmatrix} \\ a_6 &= \begin{pmatrix} 0.120091u^{29} - 0.480458u^{28} + \dots + 10.2319u - 3.49919 \\ 0.0593413u^{29} - 0.0838619u^{28} + \dots - 0.275735u - 0.299617 \end{pmatrix} \\ a_5 &= \begin{pmatrix} 0.120607u^{29} - 0.426449u^{28} + \dots + 10.4168u - 3.47317 \\ 0.0588257u^{29} - 0.137871u^{28} + \dots - 0.460615u - 0.325639 \end{pmatrix} \\ a_2 &= \begin{pmatrix} 0.366931u^{29} - 1.21295u^{28} + \dots + 20.1769u - 5.68931 \\ 0.0588257u^{29} - 0.137871u^{28} + \dots - 0.460615u - 0.325639 \end{pmatrix} \\ a_1 &= \begin{pmatrix} -0.0625000u^{29} + 0.125000u^{28} + \dots + 2.93750u + 0.0625000 \\ \frac{1}{16}u^{29} - \frac{1}{8}u^{28} + \dots - \frac{31}{16}u - \frac{1}{16} \end{pmatrix} \\ a_{10} &= \begin{pmatrix} \frac{1}{16}u^{29} - \frac{1}{4}u^{28} + \dots + \frac{13}{16}u + \frac{17}{16} \\ -0.0625000u^{29} + 0.250000u^{28} + \dots - 0.812500u - 0.0625000 \end{pmatrix} \\ a_4 &= \begin{pmatrix} 0.473087u^{29} - 1.53247u^{28} + \dots + 25.7020u - 8.17014 \\ 0.0441406u^{29} - 0.0962596u^{28} + \dots + 1.16893u - 0.447249 \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes

$$= -\frac{29246499640993}{39697527734272}u^{29} + \frac{4877178426605}{2481095483392}u^{28} + \dots - \frac{648015885883539}{39697527734272}u - \frac{374986321947515}{39697527734272}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{30} + 14u^{29} + \cdots - 2399u + 256$
c_2, c_4	$u^{30} - 6u^{29} + \cdots - 31u + 16$
c_3, c_7	$u^{30} - 2u^{29} + \cdots - 96u - 256$
c_5, c_6	$8(8u^{30} + 20u^{29} + \cdots + 12u + 4)$
c_8, c_9, c_{11} c_{12}	$u^{30} + 3u^{29} + \cdots + 14u - 1$
c_{10}	$u^{30} - 6u^{29} + \cdots + 64u + 256$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{30} + 10y^{29} + \cdots - 6849345y + 65536$
c_2, c_4	$y^{30} - 14y^{29} + \cdots + 2399y + 256$
c_3, c_7	$y^{30} + 18y^{29} + \cdots + 76800y + 65536$
c_5, c_6	$64(64y^{30} - 1360y^{29} + \cdots - 192y + 16)$
c_8, c_9, c_{11} c_{12}	$y^{30} + 23y^{29} + \cdots - 290y + 1$
c_{10}	$y^{30} + 6y^{29} + \cdots - 1789952y + 65536$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.151717 + 0.986186I$		
$a = -1.190140 - 0.027428I$	$-5.68347 - 0.58233I$	$-8.3451 + 12.8590I$
$b = 1.54321 - 0.20318I$		
$u = 0.151717 - 0.986186I$		
$a = -1.190140 + 0.027428I$	$-5.68347 + 0.58233I$	$-8.3451 - 12.8590I$
$b = 1.54321 + 0.20318I$		
$u = 0.783732 + 0.743673I$		
$a = 0.631228 - 0.256038I$	$-1.33282 - 2.23553I$	$-4.59779 + 3.41546I$
$b = -0.083764 + 0.794094I$		
$u = 0.783732 - 0.743673I$		
$a = 0.631228 + 0.256038I$	$-1.33282 + 2.23553I$	$-4.59779 - 3.41546I$
$b = -0.083764 - 0.794094I$		
$u = 1.09098$		
$a = 1.70930$	-2.65754	30.3480
$b = 0.608605$		
$u = -0.576049 + 1.073970I$		
$a = -0.261951 - 0.195372I$	$1.34016 + 8.01200I$	$2.34579 - 12.49324I$
$b = -0.151381 + 0.594028I$		
$u = -0.576049 - 1.073970I$		
$a = -0.261951 + 0.195372I$	$1.34016 - 8.01200I$	$2.34579 + 12.49324I$
$b = -0.151381 - 0.594028I$		
$u = 0.625133 + 0.199278I$		
$a = 1.38297 + 2.15219I$	$-2.83148 - 0.66530I$	$-18.2240 - 6.7846I$
$b = 0.386290 - 0.372621I$		
$u = 0.625133 - 0.199278I$		
$a = 1.38297 - 2.15219I$	$-2.83148 + 0.66530I$	$-18.2240 + 6.7846I$
$b = 0.386290 + 0.372621I$		
$u = -0.117957 + 1.367060I$		
$a = 0.699932 - 0.475481I$	$6.76007 + 1.66777I$	$0.57274 - 1.46728I$
$b = -1.78516 + 0.43773I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.117957 - 1.367060I$		
$a = 0.699932 + 0.475481I$	$6.76007 - 1.66777I$	$0.57274 + 1.46728I$
$b = -1.78516 - 0.43773I$		
$u = -0.244597 + 1.352860I$		
$a = 0.31073 + 1.77800I$	$5.09939 + 5.42433I$	$-0.68852 - 4.20178I$
$b = -0.22783 - 1.65229I$		
$u = -0.244597 - 1.352860I$		
$a = 0.31073 - 1.77800I$	$5.09939 - 5.42433I$	$-0.68852 + 4.20178I$
$b = -0.22783 + 1.65229I$		
$u = 0.259417 + 1.365310I$		
$a = -0.19856 + 1.74019I$	$10.78970 - 8.07891I$	$-0.53452 + 5.22614I$
$b = -0.91810 - 1.71117I$		
$u = 0.259417 - 1.365310I$		
$a = -0.19856 - 1.74019I$	$10.78970 + 8.07891I$	$-0.53452 - 5.22614I$
$u = -0.35445 + 1.40429I$		
$a = -0.544865 + 0.390463I$	$7.38806 + 8.84406I$	$-1.28461 - 6.03116I$
$b = 1.65162 + 0.07394I$		
$u = -0.35445 - 1.40429I$		
$a = -0.544865 - 0.390463I$	$7.38806 - 8.84406I$	$-1.28461 + 6.03116I$
$b = 1.65162 - 0.07394I$		
$u = 0.06550 + 1.45294I$		
$a = -0.04154 - 1.68423I$	$13.33180 - 0.21203I$	$1.73134 + 0.I$
$b = 0.69837 + 1.94525I$		
$u = 0.06550 - 1.45294I$		
$a = -0.04154 + 1.68423I$	$13.33180 + 0.21203I$	$1.73134 + 0.I$
$b = 0.69837 - 1.94525I$		
$u = 0.339570 + 0.379677I$		
$a = 0.592119 + 0.913683I$	$-0.505428 - 1.105530I$	$-5.83661 + 6.57696I$
$b = -0.452974 - 0.318485I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.339570 - 0.379677I$		
$a = 0.592119 - 0.913683I$	$-0.505428 + 1.105530I$	$-5.83661 - 6.57696I$
$b = -0.452974 + 0.318485I$		
$u = 1.52530 + 0.11249I$		
$a = 0.155403 + 0.304847I$	$2.40331 + 3.17859I$	$0. - 3.79440I$
$b = 0.24118 - 1.48607I$		
$u = 1.52530 - 0.11249I$		
$a = 0.155403 - 0.304847I$	$2.40331 - 3.17859I$	$0. + 3.79440I$
$b = 0.24118 + 1.48607I$		
$u = -0.58141 + 1.45298I$		
$a = 0.76540 + 1.55713I$	$12.0126 + 17.2447I$	$-2.29441 - 8.43337I$
$b = 0.78764 - 1.58037I$		
$u = -0.58141 - 1.45298I$		
$a = 0.76540 - 1.55713I$	$12.0126 - 17.2447I$	$-2.29441 + 8.43337I$
$b = 0.78764 + 1.58037I$		
$u = -0.414284 + 0.113025I$		
$a = 0.170265 - 1.051000I$	$2.31731 - 2.61856I$	$1.56244 + 2.01080I$
$b = -0.229191 - 1.205720I$		
$u = -0.414284 - 0.113025I$		
$a = 0.170265 + 1.051000I$	$2.31731 + 2.61856I$	$1.56244 - 2.01080I$
$b = -0.229191 + 1.205720I$		
$u = -0.47748 + 1.50384I$		
$a = -0.58934 - 1.58578I$	$14.1647 + 9.9224I$	$0. - 4.39684I$
$b = -0.50245 + 1.81791I$		
$u = -0.47748 - 1.50384I$		
$a = -0.58934 + 1.58578I$	$14.1647 - 9.9224I$	$0. + 4.39684I$
$b = -0.50245 - 1.81791I$		
$u = -0.0592723$		
$a = -10.2226$	-1.19030	-8.24210
$b = -0.523532$		

$$\text{II. } I_2^u = \langle 9.19 \times 10^{50}u^{41} + 6.53 \times 10^{51}u^{40} + \dots + 1.69 \times 10^{52}b + 4.77 \times 10^{52}, -4.06 \times 10^{51}u^{41} - 1.47 \times 10^{52}u^{40} + \dots + 1.18 \times 10^{53}a - 9.06 \times 10^{53}, u^{42} + 8u^{41} + \dots + 406u + 49 \rangle$$

(i) **Arc colorings**

$$\begin{aligned} a_8 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_9 &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_3 &= \begin{pmatrix} 0.0343435u^{41} + 0.124076u^{40} + \dots + 50.6244u + 7.66624 \\ -0.0544776u^{41} - 0.387181u^{40} + \dots - 17.0063u - 2.82616 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} -u \\ u \end{pmatrix} \\ a_7 &= \begin{pmatrix} -0.137949u^{41} - 1.03303u^{40} + \dots - 26.6535u - 2.02469 \\ -0.0157309u^{41} - 0.0881611u^{40} + \dots + 22.8491u + 3.77512 \end{pmatrix} \\ a_6 &= \begin{pmatrix} -0.162962u^{41} - 1.21826u^{40} + \dots - 71.3909u - 9.25731 \\ -0.0118963u^{41} - 0.0998927u^{40} + \dots + 18.0364u + 3.04637 \end{pmatrix} \\ a_5 &= \begin{pmatrix} -0.146119u^{41} - 1.10192u^{40} + \dots - 47.1902u - 5.30246 \\ -0.0287391u^{41} - 0.216234u^{40} + \dots - 6.16428u - 0.908484 \end{pmatrix} \\ a_2 &= \begin{pmatrix} 0.187628u^{41} + 1.31124u^{40} + \dots + 100.721u + 14.2062 \\ -0.0287391u^{41} - 0.216234u^{40} + \dots - 6.16428u - 0.908484 \end{pmatrix} \\ a_1 &= \begin{pmatrix} 0.0204082u^{41} + 0.163265u^{40} + \dots + 36.6122u + 8.28571 \\ 0.0941921u^{41} + 0.708146u^{40} + \dots + 54.1974u + 8.58973 \end{pmatrix} \\ a_{10} &= \begin{pmatrix} 0.175301u^{41} + 1.30821u^{40} + \dots + 114.966u + 16.9746 \\ -1 \end{pmatrix} \\ a_4 &= \begin{pmatrix} 0.0888211u^{41} + 0.511257u^{40} + \dots + 67.6307u + 10.4924 \\ -0.0544776u^{41} - 0.387181u^{40} + \dots - 17.0063u - 2.82616 \end{pmatrix} \end{aligned}$$

(ii) **Obstruction class** = -1

(iii) **Cusp Shapes** = $-0.137513u^{41} - 1.10677u^{40} + \dots - 86.1411u - 17.7083$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$(u^{21} + 6u^{20} + \cdots - 2u + 1)^2$
c_2, c_4	$(u^{21} - 4u^{20} + \cdots - 2u + 1)^2$
c_3, c_7	$(u^{21} - u^{20} + \cdots + 4u + 8)^2$
c_5, c_6	$u^{42} + 8u^{41} + \cdots + 859266u + 387139$
c_8, c_9, c_{11} c_{12}	$u^{42} - 8u^{41} + \cdots - 406u + 49$
c_{10}	$(u^{21} + 2u^{20} + \cdots + u - 1)^2$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$(y^{21} + 22y^{20} + \cdots + 66y - 1)^2$
c_2, c_4	$(y^{21} - 6y^{20} + \cdots - 2y - 1)^2$
c_3, c_7	$(y^{21} + 21y^{20} + \cdots - 176y - 64)^2$
c_5, c_6	$y^{42} - 26y^{41} + \cdots - 934716639340y + 149876605321$
c_8, c_9, c_{11} c_{12}	$y^{42} + 30y^{41} + \cdots + 10976y + 2401$
c_{10}	$(y^{21} - 8y^{20} + \cdots + 17y - 1)^2$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.041887 + 1.002630I$		
$a = -8.74756 - 5.85633I$	2.07989	$-13.37190 + 0.I$
$b = -0.492750$		
$u = -0.041887 - 1.002630I$		
$a = -8.74756 + 5.85633I$	2.07989	$-13.37190 + 0.I$
$b = -0.492750$		
$u = -0.870370 + 0.220126I$		
$a = 1.199530 - 0.425810I$	2.22124 + 4.45806I	$-4.43689 - 6.14529I$
$b = 1.088250 - 0.021385I$		
$u = -0.870370 - 0.220126I$		
$a = 1.199530 + 0.425810I$	2.22124 - 4.45806I	$-4.43689 + 6.14529I$
$b = 1.088250 + 0.021385I$		
$u = -0.769906 + 0.433901I$		
$a = 0.483221 - 0.092871I$	-0.56968 - 2.93752I	$-1.02400 + 3.43881I$
$b = -0.006772 - 0.621655I$		
$u = -0.769906 - 0.433901I$		
$a = 0.483221 + 0.092871I$	-0.56968 + 2.93752I	$-1.02400 - 3.43881I$
$b = -0.006772 + 0.621655I$		
$u = 0.600601 + 0.944887I$		
$a = -0.154802 + 0.082043I$	-0.56968 - 2.93752I	$0. + 3.43881I$
$b = -0.006772 - 0.621655I$		
$u = 0.600601 - 0.944887I$		
$a = -0.154802 - 0.082043I$	-0.56968 + 2.93752I	$0. - 3.43881I$
$b = -0.006772 + 0.621655I$		
$u = -0.475850 + 1.016220I$		
$a = 0.272799 + 0.705259I$	4.75904 + 0.34630I	$1.96536 + 0.I$
$b = 0.528856 + 0.467306I$		
$u = -0.475850 - 1.016220I$		
$a = 0.272799 - 0.705259I$	4.75904 - 0.34630I	$1.96536 + 0.I$
$b = 0.528856 - 0.467306I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.053090 + 1.155400I$		
$a = -0.058655 - 0.325592I$	$1.45515 - 0.21101I$	$-6.00000 + 0.I$
$b = -0.899194 - 0.226112I$		
$u = 0.053090 - 1.155400I$		
$a = -0.058655 + 0.325592I$	$1.45515 + 0.21101I$	$-6.00000 + 0.I$
$b = -0.899194 + 0.226112I$		
$u = 0.216321 + 1.202960I$		
$a = -0.33507 - 2.15389I$	$0.26332 - 2.36605I$	0
$b = -0.157544 + 0.891019I$		
$u = 0.216321 - 1.202960I$		
$a = -0.33507 + 2.15389I$	$0.26332 + 2.36605I$	0
$b = -0.157544 - 0.891019I$		
$u = 0.474335 + 0.591031I$		
$a = 1.05367 + 1.22545I$	$6.58039 + 1.36266I$	$-4.18856 - 2.27516I$
$b = 0.00145 + 1.46011I$		
$u = 0.474335 - 0.591031I$		
$a = 1.05367 - 1.22545I$	$6.58039 - 1.36266I$	$-4.18856 + 2.27516I$
$b = 0.00145 - 1.46011I$		
$u = -0.185639 + 1.238440I$		
$a = -0.70977 - 2.01041I$	$5.71484 + 4.94435I$	0
$b = -0.45321 + 1.45865I$		
$u = -0.185639 - 1.238440I$		
$a = -0.70977 + 2.01041I$	$5.71484 - 4.94435I$	0
$b = -0.45321 - 1.45865I$		
$u = -0.021801 + 1.271800I$		
$a = 0.50394 + 2.30405I$	$6.58039 - 1.36266I$	0
$b = 0.00145 - 1.46011I$		
$u = -0.021801 - 1.271800I$		
$a = 0.50394 - 2.30405I$	$6.58039 + 1.36266I$	0
$b = 0.00145 + 1.46011I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.252620 + 0.261440I$		
$a = -0.093957 - 0.166544I$	$8.50490 + 3.89686I$	0
$b = -0.15224 + 1.62071I$		
$u = -1.252620 - 0.261440I$		
$a = -0.093957 + 0.166544I$	$8.50490 - 3.89686I$	0
$b = -0.15224 - 1.62071I$		
$u = -1.302650 + 0.047271I$		
$a = 0.370885 + 0.223986I$	$7.25306 + 10.68720I$	0
$b = 0.55439 - 1.54207I$		
$u = -1.302650 - 0.047271I$		
$a = 0.370885 - 0.223986I$	$7.25306 - 10.68720I$	0
$b = 0.55439 + 1.54207I$		
$u = -0.226174 + 1.289570I$		
$a = 0.560692 + 0.745306I$	$4.75904 - 0.34630I$	0
$b = 0.528856 - 0.467306I$		
$u = -0.226174 - 1.289570I$		
$a = 0.560692 - 0.745306I$	$4.75904 + 0.34630I$	0
$b = 0.528856 + 0.467306I$		
$u = 0.588336 + 0.271246I$		
$a = -0.350608 - 1.017080I$	$5.71484 - 4.94435I$	$-5.24866 + 2.70559I$
$b = -0.45321 - 1.45865I$		
$u = 0.588336 - 0.271246I$		
$a = -0.350608 + 1.017080I$	$5.71484 + 4.94435I$	$-5.24866 - 2.70559I$
$b = -0.45321 + 1.45865I$		
$u = 0.319588 + 1.353120I$		
$a = -0.014969 - 0.198168I$	$2.22124 - 4.45806I$	0
$b = 1.088250 + 0.021385I$		
$u = 0.319588 - 1.353120I$		
$a = -0.014969 + 0.198168I$	$2.22124 + 4.45806I$	0
$b = 1.088250 - 0.021385I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.579219 + 0.187158I$		
$a = -1.56373 + 0.83045I$	$0.26332 + 2.36605I$	$-4.59037 - 2.67274I$
$b = -0.157544 - 0.891019I$		
$u = -0.579219 - 0.187158I$		
$a = -1.56373 - 0.83045I$	$0.26332 - 2.36605I$	$-4.59037 + 2.67274I$
$b = -0.157544 + 0.891019I$		
$u = -0.239978 + 0.362151I$		
$a = -3.20155 + 1.86451I$	$1.45515 + 0.21101I$	$-7.18710 - 0.57244I$
$b = -0.899194 + 0.226112I$		
$u = -0.239978 - 0.362151I$		
$a = -3.20155 - 1.86451I$	$1.45515 - 0.21101I$	$-7.18710 + 0.57244I$
$b = -0.899194 - 0.226112I$		
$u = 0.62069 + 1.53093I$		
$a = 0.66906 - 1.34449I$	$7.25306 - 10.68720I$	0
$b = 0.55439 + 1.54207I$		
$u = 0.62069 - 1.53093I$		
$a = 0.66906 + 1.34449I$	$7.25306 + 10.68720I$	0
$b = 0.55439 - 1.54207I$		
$u = 0.45894 + 1.59821I$		
$a = -0.42722 + 1.36484I$	$8.50490 - 3.89686I$	0
$b = -0.15224 - 1.62071I$		
$u = 0.45894 - 1.59821I$		
$a = -0.42722 - 1.36484I$	$8.50490 + 3.89686I$	0
$b = -0.15224 + 1.62071I$		
$u = -0.76320 + 1.48676I$		
$a = 0.794802 + 1.042100I$	$12.12580 + 3.51416I$	0
$b = 0.24239 - 1.67299I$		
$u = -0.76320 - 1.48676I$		
$a = 0.794802 - 1.042100I$	$12.12580 - 3.51416I$	0
$b = 0.24239 + 1.67299I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.60260 + 1.60806I$		
$a = -0.536403 - 1.053510I$	$12.12580 - 3.51416I$	0
$b = 0.24239 + 1.67299I$		
$u = -0.60260 - 1.60806I$		
$a = -0.536403 + 1.053510I$	$12.12580 + 3.51416I$	0
$b = 0.24239 - 1.67299I$		

$$\text{III. } I_3^u = \langle b, -u^3 + u^2 + 4a + 2u + 3, u^4 + u^2 - u + 1 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_8 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_9 &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_3 &= \begin{pmatrix} \frac{1}{4}u^3 - \frac{1}{4}u^2 - \frac{1}{2}u - \frac{3}{4} \\ 0 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} -u \\ u \end{pmatrix} \\ a_7 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_6 &= \begin{pmatrix} u^2 + 1 \\ -u^2 + u - 1 \end{pmatrix} \\ a_5 &= \begin{pmatrix} -u^3 + u^2 + 1 \\ u^3 - u^2 + u - 1 \end{pmatrix} \\ a_2 &= \begin{pmatrix} \frac{5}{4}u^3 - \frac{5}{4}u^2 - \frac{1}{2}u - \frac{7}{4} \\ -u^3 + u^2 - u + 1 \end{pmatrix} \\ a_1 &= \begin{pmatrix} u^3 - u^2 - 1 \\ -u^3 + u^2 - u + 1 \end{pmatrix} \\ a_{10} &= \begin{pmatrix} -u^3 - u^2 \\ u^3 + u^2 + 1 \end{pmatrix} \\ a_4 &= \begin{pmatrix} \frac{1}{4}u^3 - \frac{1}{4}u^2 - \frac{1}{2}u - \frac{3}{4} \\ 0 \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = 1

$$(iii) \text{ Cusp Shapes} = \frac{49}{16}u^3 + \frac{43}{16}u^2 + \frac{21}{8}u - \frac{163}{16}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_2	$(u - 1)^4$
c_3, c_7	u^4
c_4	$(u + 1)^4$
c_5, c_6	$u^4 + 2u^3 + 3u^2 + u + 1$
c_8, c_9	$u^4 + u^2 - u + 1$
c_{10}	$u^4 + 3u^3 + 4u^2 + 3u + 2$
c_{11}, c_{12}	$u^4 + u^2 + u + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_4	$(y - 1)^4$
c_3, c_7	y^4
c_5, c_6	$y^4 + 2y^3 + 7y^2 + 5y + 1$
c_8, c_9, c_{11} c_{12}	$y^4 + 2y^3 + 3y^2 + y + 1$
c_{10}	$y^4 - y^3 + 2y^2 + 7y + 4$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.547424 + 0.585652I$		
$a = -1.112690 - 0.371716I$	$-2.62503 - 1.39709I$	$-10.08957 + 4.25783I$
$b = 0$		
$u = 0.547424 - 0.585652I$		
$a = -1.112690 + 0.371716I$	$-2.62503 + 1.39709I$	$-10.08957 - 4.25783I$
$b = 0$		
$u = -0.547424 + 1.120870I$		
$a = 0.237691 - 0.353773I$	$0.98010 + 7.64338I$	$-8.37918 - 1.58240I$
$b = 0$		
$u = -0.547424 - 1.120870I$		
$a = 0.237691 + 0.353773I$	$0.98010 - 7.64338I$	$-8.37918 + 1.58240I$
$b = 0$		

$$\text{IV. } I_4^u = \langle 8a^2 + b + 18a + 4, \ 8a^3 + 20a^2 + 8a + 1, \ u - 1 \rangle$$

(i) Arc colorings

$$a_8 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} a \\ -8a^2 - 18a - 4 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -2a^2 - 4a \\ -4a^2 - 8a \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 0 \\ -2a^2 - 4a \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 2a^2 + 4a \\ -4a^2 - 8a \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -2a^2 - 4a - 2 \\ -4a^2 - 8a \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} 8a^2 + 19a + 4 \\ -8a^2 - 18a - 4 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = $3a^2 + \frac{61}{2}a - \frac{5}{4}$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_3	$u^3 - u^2 + 2u - 1$
c_2	$u^3 + u^2 - 1$
c_4	$u^3 - u^2 + 1$
c_5	$8(8u^3 - 12u^2 + 4u + 1)$
c_6	$8(8u^3 + 12u^2 + 4u - 1)$
c_7	$u^3 + u^2 + 2u + 1$
c_8, c_9	$(u - 1)^3$
c_{10}	u^3
c_{11}, c_{12}	$(u + 1)^3$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_3, c_7	$y^3 + 3y^2 + 2y - 1$
c_2, c_4	$y^3 - y^2 + 2y - 1$
c_5, c_6	$64(64y^3 - 80y^2 + 40y - 1)$
c_8, c_9, c_{11} c_{12}	$(y - 1)^3$
c_{10}	y^3

(vi) Complex Volumes and Cusp Shapes

Solutions to I_4^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.00000$		
$a = -0.230101 + 0.091291I$	$1.37919 - 2.82812I$	$-8.13425 + 2.65834I$
$b = -0.215080 - 1.307140I$		
$u = 1.00000$		
$a = -0.230101 - 0.091291I$	$1.37919 + 2.82812I$	$-8.13425 - 2.65834I$
$b = -0.215080 + 1.307140I$		
$u = 1.00000$		
$a = -2.03980$	-2.75839	-50.9820
$b = -0.569840$		

$$\mathbf{V} \cdot I_5^u = \langle b, -u^3 + a - u - 1, u^6 + u^5 + 2u^4 + 2u^3 + 2u^2 + 2u + 1 \rangle$$

(i) Arc colorings

$$a_8 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} u^3 + u + 1 \\ 0 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -u \\ u \end{pmatrix}$$

$$a_7 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} u^2 + 1 \\ u^4 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} u^5 + u^4 + 2u^3 + 2u^2 + 2u + 2 \\ -u^5 - 2u^3 - u^2 - 2u - 1 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -u^5 - u^4 - u^3 - 2u^2 - u - 1 \\ u^5 + 2u^3 + u^2 + 2u + 1 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -u^5 - u^4 - 2u^3 - 2u^2 - 2u - 2 \\ u^5 + 2u^3 + u^2 + 2u + 1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -u^5 - 2u^3 - u \\ -1 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u^3 + u + 1 \\ 0 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = $-2u^5 + 3u^3 - 2u^2 + 3u - 8$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_2	$(u - 1)^6$
c_3, c_7	u^6
c_4	$(u + 1)^6$
c_5, c_6	$u^6 + 3u^5 + 4u^4 + 2u^3 + 1$
c_8, c_9	$u^6 + u^5 + 2u^4 + 2u^3 + 2u^2 + 2u + 1$
c_{10}	$(u^3 - u^2 + 1)^2$
c_{11}, c_{12}	$u^6 - u^5 + 2u^4 - 2u^3 + 2u^2 - 2u + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_4	$(y - 1)^6$
c_3, c_7	y^6
c_5, c_6	$y^6 - y^5 + 4y^4 - 2y^3 + 8y^2 + 1$
c_8, c_9, c_{11} c_{12}	$y^6 + 3y^5 + 4y^4 + 2y^3 + 1$
c_{10}	$(y^3 - y^2 + 2y - 1)^2$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_5^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.498832 + 1.001300I$		
$a = 0.122561 + 0.744862I$	$-1.37919 - 2.82812I$	$-11.71191 + 2.59975I$
$b = 0$		
$u = 0.498832 - 1.001300I$		
$a = 0.122561 - 0.744862I$	$-1.37919 + 2.82812I$	$-11.71191 - 2.59975I$
$b = 0$		
$u = -0.284920 + 1.115140I$		
$a = 1.75488$	2.75839	$-60.423824 + 0.10I$
$b = 0$		
$u = -0.284920 - 1.115140I$		
$a = 1.75488$	2.75839	$-60.423824 + 0.10I$
$b = 0$		
$u = -0.713912 + 0.305839I$		
$a = 0.122561 + 0.744862I$	$-1.37919 - 2.82812I$	$-11.71191 + 2.59975I$
$b = 0$		
$u = -0.713912 - 0.305839I$		
$a = 0.122561 - 0.744862I$	$-1.37919 + 2.82812I$	$-11.71191 - 2.59975I$
$b = 0$		

$$\text{VI. } I_6^u = \langle au + 4b + a + u - 5, a^2 + 4au - 2a + 6u - 3, u^2 + 1 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_8 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_9 &= \begin{pmatrix} 1 \\ -1 \end{pmatrix} \\ a_3 &= \begin{pmatrix} a \\ -\frac{1}{4}au - \frac{1}{4}a - \frac{1}{4}u + \frac{5}{4} \end{pmatrix} \\ a_{12} &= \begin{pmatrix} -u \\ u \end{pmatrix} \\ a_7 &= \begin{pmatrix} -\frac{1}{4}au + \frac{1}{4}a - \frac{3}{4}u + \frac{13}{4} \\ \frac{1}{4}au + \frac{1}{4}a + \frac{1}{4}u - \frac{9}{4} \end{pmatrix} \\ a_6 &= \begin{pmatrix} -\frac{1}{4}au - \frac{1}{4}a - \frac{1}{4}u + \frac{9}{4} \\ \frac{1}{4}au + \frac{3}{4}a - \frac{1}{4}u - \frac{5}{4} \end{pmatrix} \\ a_5 &= \begin{pmatrix} -\frac{1}{4}au + \frac{1}{4}a - \frac{3}{4}u + \frac{13}{4} \\ \frac{1}{4}au + \frac{1}{4}a + \frac{1}{4}u - \frac{9}{4} \end{pmatrix} \\ a_2 &= \begin{pmatrix} -\frac{1}{2}au - u + \frac{7}{2} \\ \frac{1}{4}au + \frac{1}{4}a + \frac{1}{4}u - \frac{9}{4} \end{pmatrix} \\ a_1 &= \begin{pmatrix} -\frac{3}{4}au - \frac{3}{4}a - \frac{7}{4}u + \frac{23}{4} \\ \frac{3}{4}au + \frac{3}{4}a + \frac{3}{4}u - \frac{23}{4} \end{pmatrix} \\ a_{10} &= \begin{pmatrix} \frac{3}{4}au - \frac{3}{4}a - \frac{23}{4}u - \frac{3}{4} \\ -\frac{3}{4}au + \frac{3}{4}a + \frac{23}{4}u - \frac{1}{4} \end{pmatrix} \\ a_4 &= \begin{pmatrix} \frac{1}{4}au + \frac{5}{4}a + \frac{1}{4}u - \frac{5}{4} \\ -\frac{1}{4}au - \frac{1}{4}a - \frac{1}{4}u + \frac{5}{4} \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = -4

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$(u^2 - 3u + 1)^2$
c_2, c_3	$(u^2 + u - 1)^2$
c_4, c_7	$(u^2 - u - 1)^2$
c_5	$u^4 - 6u^3 + 18u^2 - 12u + 4$
c_6	$u^4 + 6u^3 + 18u^2 + 12u + 4$
c_8, c_9, c_{11} c_{12}	$(u^2 + 1)^2$
c_{10}	$u^4 + 7u^2 + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$(y^2 - 7y + 1)^2$
c_2, c_3, c_4 c_7	$(y^2 - 3y + 1)^2$
c_5, c_6	$y^4 + 188y^2 + 16$
c_8, c_9, c_{11} c_{12}	$(y + 1)^4$
c_{10}	$(y^2 + 7y + 1)^2$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_6^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.000000I$		
$a = -1.236070 + 0.236068I$	-5.59278	-4.00000
$b = 1.61803$		
$u = 1.000000I$		
$a = 3.23607 - 4.23607I$	2.30291	-4.00000
$b = -0.618034$		
$u = -1.000000I$		
$a = -1.236070 - 0.236068I$	-5.59278	-4.00000
$b = 1.61803$		
$u = -1.000000I$		
$a = 3.23607 + 4.23607I$	2.30291	-4.00000
$b = -0.618034$		

VII. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1	$((u - 1)^{10})(u^2 - 3u + 1)^2(u^3 - u^2 + 2u - 1)(u^{21} + 6u^{20} + \dots - 2u + 1)^2$ $\cdot (u^{30} + 14u^{29} + \dots - 2399u + 256)$
c_2	$((u - 1)^{10})(u^2 + u - 1)^2(u^3 + u^2 - 1)(u^{21} - 4u^{20} + \dots - 2u + 1)^2$ $\cdot (u^{30} - 6u^{29} + \dots - 31u + 16)$
c_3	$u^{10}(u^2 + u - 1)^2(u^3 - u^2 + 2u - 1)(u^{21} - u^{20} + \dots + 4u + 8)^2$ $\cdot (u^{30} - 2u^{29} + \dots - 96u - 256)$
c_4	$((u + 1)^{10})(u^2 - u - 1)^2(u^3 - u^2 + 1)(u^{21} - 4u^{20} + \dots - 2u + 1)^2$ $\cdot (u^{30} - 6u^{29} + \dots - 31u + 16)$
c_5	$64(8u^3 - 12u^2 + 4u + 1)(u^4 - 6u^3 + 18u^2 - 12u + 4)$ $\cdot (u^4 + 2u^3 + 3u^2 + u + 1)(u^6 + 3u^5 + 4u^4 + 2u^3 + 1)$ $\cdot (8u^{30} + 20u^{29} + \dots + 12u + 4)(u^{42} + 8u^{41} + \dots + 859266u + 387139)$
c_6	$64(8u^3 + 12u^2 + 4u - 1)(u^4 + 2u^3 + 3u^2 + u + 1)$ $\cdot (u^4 + 6u^3 + 18u^2 + 12u + 4)(u^6 + 3u^5 + 4u^4 + 2u^3 + 1)$ $\cdot (8u^{30} + 20u^{29} + \dots + 12u + 4)(u^{42} + 8u^{41} + \dots + 859266u + 387139)$
c_7	$u^{10}(u^2 - u - 1)^2(u^3 + u^2 + 2u + 1)(u^{21} - u^{20} + \dots + 4u + 8)^2$ $\cdot (u^{30} - 2u^{29} + \dots - 96u - 256)$
c_8, c_9	$((u - 1)^3)(u^2 + 1)^2(u^4 + u^2 - u + 1)(u^6 + u^5 + \dots + 2u + 1)$ $\cdot (u^{30} + 3u^{29} + \dots + 14u - 1)(u^{42} - 8u^{41} + \dots - 406u + 49)$
c_{10}	$u^3(u^3 - u^2 + 1)^2(u^4 + 7u^2 + 1)(u^4 + 3u^3 + 4u^2 + 3u + 2)$ $\cdot ((u^{21} + 2u^{20} + \dots + u - 1)^2)(u^{30} - 6u^{29} + \dots + 64u + 256)$
c_{11}, c_{12}	$((u + 1)^3)(u^2 + 1)^2(u^4 + u^2 + u + 1)(u^6 - u^5 + \dots - 2u + 1)$ $\cdot (u^{30} + 3u^{29} + \dots + 14u - 1)(u^{42} - 8u^{41} + \dots - 406u + 49)$

VIII. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1	$(y - 1)^{10}(y^2 - 7y + 1)^2(y^3 + 3y^2 + 2y - 1)$ $\cdot (y^{21} + 22y^{20} + \dots + 66y - 1)^2$ $\cdot (y^{30} + 10y^{29} + \dots - 6849345y + 65536)$
c_2, c_4	$((y - 1)^{10})(y^2 - 3y + 1)^2(y^3 - y^2 + 2y - 1)(y^{21} - 6y^{20} + \dots - 2y - 1)^2$ $\cdot (y^{30} - 14y^{29} + \dots + 2399y + 256)$
c_3, c_7	$y^{10}(y^2 - 3y + 1)^2(y^3 + 3y^2 + 2y - 1)(y^{21} + 21y^{20} + \dots - 176y - 64)^2$ $\cdot (y^{30} + 18y^{29} + \dots + 76800y + 65536)$
c_5, c_6	$4096(64y^3 - 80y^2 + 40y - 1)(y^4 + 188y^2 + 16)(y^4 + 2y^3 + \dots + 5y + 1)$ $\cdot (y^6 - y^5 + 4y^4 - 2y^3 + 8y^2 + 1)(64y^{30} - 1360y^{29} + \dots - 192y + 16)$ $\cdot (y^{42} - 26y^{41} + \dots - 934716639340y + 149876605321)$
c_8, c_9, c_{11} c_{12}	$((y - 1)^3)(y + 1)^4(y^4 + 2y^3 + \dots + y + 1)(y^6 + 3y^5 + \dots + 2y^3 + 1)$ $\cdot (y^{30} + 23y^{29} + \dots - 290y + 1)(y^{42} + 30y^{41} + \dots + 10976y + 2401)$
c_{10}	$y^3(y^2 + 7y + 1)^2(y^3 - y^2 + 2y - 1)^2(y^4 - y^3 + 2y^2 + 7y + 4)$ $\cdot ((y^{21} - 8y^{20} + \dots + 17y - 1)^2)(y^{30} + 6y^{29} + \dots - 1789952y + 65536)$