



12n₀₆₁₂ (K12n₀₆₁₂)



A knot diagram¹

Linearized knot diagram



Solving Sequence

$$1,4 \xrightarrow{c_4} 5,10 \xrightarrow{c_5} 6 \xrightarrow{c_9} 9 \xrightarrow{c_8} 8 \xrightarrow{c_{12}} 12 \xrightarrow{c_{11}} 11 \xrightarrow{c_3} 3 \xrightarrow{c_1} 2 \xrightarrow{c_7} 7 \twoheadrightarrow c_2, c_6, c_{10}$$

Ideals for irreducible components² of X_{par}

$$I_1^u = \langle b - u,$$

$$65065875226664520u^{27} + 9394396496362895u^{26} + \dots + 76364734116743a + 26735659168418070, \\ u^{28} + 23u^{26} + \dots - 3u + 1 \rangle$$

$$I_2^u = \langle b + u, 5u^{16} + u^{15} + \dots + a - 5,$$

$$u^{17} + 3u^{15} - u^{14} - 4u^{13} - 5u^{11} + 5u^{10} + 11u^9 - 6u^8 - 2u^7 - 2u^6 - 7u^5 + 8u^4 + 5u^3 - 5u^2 - u + 1 \rangle$$

$$I_3^u = \langle 1.90741 \times 10^{76}u^{35} + 2.50394 \times 10^{76}u^{34} + \dots + 5.34816 \times 10^{78}b - 7.63438 \times 10^{78},$$

$$4.20379 \times 10^{66}u^{35} + 2.74371 \times 10^{66}u^{34} + \dots + 1.04184 \times 10^{69}a - 1.38242 \times 10^{70},$$

$$u^{36} + u^{35} + \dots - 2560u + 121 \rangle$$

* 3 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 81 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/maths/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$\mathbf{I. } I_1^u = \langle b - u, 6.51 \times 10^{16}u^{27} + 9.39 \times 10^{15}u^{26} + \dots + 7.64 \times 10^{13}a + 2.67 \times 10^{16}, u^{28} + 23u^{26} + \dots - 3u + 1 \rangle$$

(i) Arc colorings

$$a_1 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -852.041u^{27} - 123.020u^{26} + \dots - 1960.13u - 350.105 \\ u \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -104.065u^{27} + 370.478u^{26} + \dots - 4216.19u + 1309.36 \\ -42.7858u^{27} + 67.8173u^{26} + \dots - 845.364u + 240.781 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -852.041u^{27} - 123.020u^{26} + \dots - 1959.13u - 350.105 \\ u \end{pmatrix}$$

$$a_8 = \begin{pmatrix} -852.041u^{27} - 123.020u^{26} + \dots - 1959.13u - 350.105 \\ 240.781u^{27} + 42.7858u^{26} + \dots + 483.981u + 123.020 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -456.322u^{27} + 18.9555u^{26} + \dots - 1801.95u + 59.0018 \\ -240.781u^{27} - 42.7858u^{26} + \dots - 481.981u - 123.020 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -697.103u^{27} - 23.8304u^{26} + \dots - 2283.93u - 64.0183 \\ -240.781u^{27} - 42.7858u^{26} + \dots - 481.981u - 123.020 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -186.897u^{27} + 111.670u^{26} + \dots - 1633.52u + 381.526 \\ 80.2342u^{27} - 172.964u^{26} + \dots + 2060.86u - 611.260 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 643.875u^{27} + 247.704u^{26} + \dots - 217.453u + 810.960 \\ -403.888u^{27} + 1.74743u^{26} + \dots - 1518.50u + 43.0095 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} 20.8672u^{27} - 274.777u^{26} + \dots + 2601.82u - 917.955 \\ -35.7010u^{27} + 222.640u^{26} + \dots - 2384.15u + 774.366 \end{pmatrix}$$

(ii) Obstruction class = -1

$$\mathbf{(iii) Cusp Shapes = } -\frac{57088194451595465}{76364734116743}u^{27} + \frac{88581311353921087}{76364734116743}u^{26} + \dots - \frac{1111026347572146836}{76364734116743}u + \frac{313608959053702506}{76364734116743}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_6	$u^{28} + 9u^{27} + \dots + 84u + 16$
c_2, c_7	$u^{28} + 5u^{27} + \dots + 2u + 4$
c_3, c_5, c_{10} c_{11}	$u^{28} + u^{27} + \dots + 4u + 1$
c_4, c_9	$u^{28} + 23u^{26} + \dots + 3u + 1$
c_8	$u^{28} - 25u^{27} + \dots - 98862u + 9028$
c_{12}	$u^{28} + 21u^{27} + \dots + 3584u + 512$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_6	$y^{28} + 21y^{27} + \dots + 2064y + 256$
c_2, c_7	$y^{28} + 9y^{27} + \dots + 84y + 16$
c_3, c_5, c_{10} c_{11}	$y^{28} - 11y^{27} + \dots - 6y + 1$
c_4, c_9	$y^{28} + 46y^{27} + \dots + 49y + 1$
c_8	$y^{28} - 3y^{27} + \dots + 785869044y + 81504784$
c_{12}	$y^{28} - 17y^{27} + \dots + 4456448y + 262144$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.062917 + 0.794694I$		
$a = -1.44001 - 0.70376I$	$-1.93668 - 4.60321I$	$-5.64825 + 3.26019I$
$b = -0.062917 + 0.794694I$		
$u = -0.062917 - 0.794694I$		
$a = -1.44001 + 0.70376I$	$-1.93668 + 4.60321I$	$-5.64825 - 3.26019I$
$b = -0.062917 - 0.794694I$		
$u = 0.193820 + 0.693750I$		
$a = 1.312680 - 0.250146I$	$-0.409488 - 0.484105I$	$-3.68035 + 2.98454I$
$b = 0.193820 + 0.693750I$		
$u = 0.193820 - 0.693750I$		
$a = 1.312680 + 0.250146I$	$-0.409488 + 0.484105I$	$-3.68035 - 2.98454I$
$b = 0.193820 - 0.693750I$		
$u = -0.015958 + 0.620215I$		
$a = -2.15814 - 0.25579I$	$-5.53121 + 1.96442I$	$-8.05770 - 3.63784I$
$b = -0.015958 + 0.620215I$		
$u = -0.015958 - 0.620215I$		
$a = -2.15814 + 0.25579I$	$-5.53121 - 1.96442I$	$-8.05770 + 3.63784I$
$b = -0.015958 - 0.620215I$		
$u = 0.007916 + 0.540155I$		
$a = -2.68846 + 0.36830I$	$-0.95872 + 8.32918I$	$-2.85757 - 9.28705I$
$b = 0.007916 + 0.540155I$		
$u = 0.007916 - 0.540155I$		
$a = -2.68846 - 0.36830I$	$-0.95872 - 8.32918I$	$-2.85757 + 9.28705I$
$b = 0.007916 - 0.540155I$		
$u = 0.018995 + 0.532352I$		
$a = 2.36847 + 0.52484I$	$0.30181 - 2.82865I$	$-1.29087 + 4.58370I$
$b = 0.018995 + 0.532352I$		
$u = 0.018995 - 0.532352I$		
$a = 2.36847 - 0.52484I$	$0.30181 + 2.82865I$	$-1.29087 - 4.58370I$
$b = 0.018995 - 0.532352I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.25002 + 1.56539I$ $a = 0.238631 - 0.969260I$ $b = -0.25002 + 1.56539I$	$0.93824 + 1.99231I$	0
$u = -0.25002 - 1.56539I$ $a = 0.238631 + 0.969260I$ $b = -0.25002 - 1.56539I$	$0.93824 - 1.99231I$	0
$u = 0.005308 + 0.404634I$ $a = 0.23723 + 2.20308I$ $b = 0.005308 + 0.404634I$	$3.88624 - 2.72415I$	$0.13151 + 3.72503I$
$u = 0.005308 - 0.404634I$ $a = 0.23723 - 2.20308I$ $b = 0.005308 - 0.404634I$	$3.88624 + 2.72415I$	$0.13151 - 3.72503I$
$u = 0.182846 + 0.331511I$ $a = 0.690101 + 0.486506I$ $b = 0.182846 + 0.331511I$	$-0.224067 - 0.893165I$	$-4.75660 + 7.55885I$
$u = 0.182846 - 0.331511I$ $a = 0.690101 - 0.486506I$ $b = 0.182846 - 0.331511I$	$-0.224067 + 0.893165I$	$-4.75660 - 7.55885I$
$u = -0.50624 + 1.65356I$ $a = 0.039303 - 0.997212I$ $b = -0.50624 + 1.65356I$	$-0.51665 + 9.12206I$	0
$u = -0.50624 - 1.65356I$ $a = 0.039303 + 0.997212I$ $b = -0.50624 - 1.65356I$	$-0.51665 - 9.12206I$	0
$u = 0.35393 + 1.72346I$ $a = -0.116049 - 0.916399I$ $b = 0.35393 + 1.72346I$	$3.78322 - 6.15036I$	0
$u = 0.35393 - 1.72346I$ $a = -0.116049 + 0.916399I$ $b = 0.35393 - 1.72346I$	$3.78322 + 6.15036I$	0

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.62895 + 1.78797I$	$5.5259 + 15.3858I$	0
$a = -0.052590 - 0.928175I$		
$b = -0.62895 + 1.78797I$		
$u = -0.62895 - 1.78797I$	$5.5259 - 15.3858I$	0
$a = -0.052590 + 0.928175I$		
$b = -0.62895 - 1.78797I$		
$u = 0.57825 + 1.81066I$	$6.63527 - 9.39925I$	0
$a = 0.026177 - 0.911601I$		
$b = 0.57825 + 1.81066I$		
$u = 0.57825 - 1.81066I$	$6.63527 + 9.39925I$	0
$a = 0.026177 + 0.911601I$		
$b = 0.57825 - 1.81066I$		
$u = 0.16208 + 1.89611I$	$9.28183 - 2.22011I$	0
$a = 0.245895 - 0.654376I$		
$b = 0.16208 + 1.89611I$		
$u = 0.16208 - 1.89611I$	$9.28183 + 2.22011I$	0
$a = 0.245895 + 0.654376I$		
$b = 0.16208 - 1.89611I$		
$u = -0.03905 + 1.92499I$	$9.65564 - 4.10797I$	0
$a = -0.203229 - 0.694596I$		
$b = -0.03905 + 1.92499I$		
$u = -0.03905 - 1.92499I$	$9.65564 + 4.10797I$	0
$a = -0.203229 + 0.694596I$		
$b = -0.03905 - 1.92499I$		

$$\text{II. } I_2^u = \langle b + u, 5u^{16} + u^{15} + \dots + a - 5, u^{17} + 3u^{15} + \dots - u + 1 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_1 &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_4 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_5 &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_{10} &= \begin{pmatrix} -5u^{16} - u^{15} + \dots + 12u + 5 \\ -u \end{pmatrix} \\ a_6 &= \begin{pmatrix} 5u^{16} + 5u^{15} + \dots + u - 16 \\ u^2 + 1 \end{pmatrix} \\ a_9 &= \begin{pmatrix} -5u^{16} - u^{15} + \dots + 11u + 5 \\ -u \end{pmatrix} \\ a_8 &= \begin{pmatrix} -5u^{16} - u^{15} + \dots + 11u + 5 \\ -u^{16} - 3u^{14} + \dots + 3u + 1 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} 12u^{16} + 4u^{15} + \dots - 17u - 11 \\ -u^{16} - 3u^{14} + \dots + 5u + 1 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 11u^{16} + 4u^{15} + \dots - 12u - 10 \\ -u^{16} - 3u^{14} + \dots + 5u + 1 \end{pmatrix} \\ a_3 &= \begin{pmatrix} -10u^{16} - 11u^{15} + \dots - 5u + 23 \\ u^{16} + u^{15} + \dots + 13u^2 - 6 \end{pmatrix} \\ a_2 &= \begin{pmatrix} -16u^{16} - 10u^{15} + \dots + 13u + 29 \\ -23u^{16} - 10u^{15} + \dots + 22u + 18 \end{pmatrix} \\ a_7 &= \begin{pmatrix} 9u^{16} + 13u^{15} + \dots + 12u - 21 \\ u^{16} + 2u^{15} + \dots + 3u + 4 \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = 1

$$\text{(iii) Cusp Shapes} = 45u^{16} + 45u^{15} + 162u^{14} + 117u^{13} - 128u^{12} - 110u^{11} - 304u^{10} - 68u^9 + 533u^8 + 181u^7 - 36u^6 - 69u^5 - 426u^4 + u^3 + 315u^2 - u - 77$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_6	$u^{17} - 6u^{16} + \dots - 18u^2 + 1$
c_2	$u^{17} + 3u^{15} + \dots + 2u - 1$
c_3, c_{10}	$u^{17} - u^{16} + \dots + 3u^2 + 1$
c_4, c_9	$u^{17} + 3u^{15} + \dots - u + 1$
c_5, c_{11}	$u^{17} + u^{16} + \dots - 3u^2 - 1$
c_7	$u^{17} + 3u^{15} + \dots + 2u + 1$
c_8	$u^{17} - u^{15} + \dots + 2u + 1$
c_{12}	$u^{17} - 6u^{16} + \dots + 2u + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_6	$y^{17} + 14y^{16} + \dots + 36y - 1$
c_2, c_7	$y^{17} + 6y^{16} + \dots + 18y^2 - 1$
c_3, c_5, c_{10} c_{11}	$y^{17} - 11y^{16} + \dots - 6y - 1$
c_4, c_9	$y^{17} + 6y^{16} + \dots + 11y - 1$
c_8	$y^{17} - 2y^{16} + \dots - 2y - 1$
c_{12}	$y^{17} - 14y^{16} + \dots + 8y - 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.944344 + 0.346033I$ $a = 0.447652 - 0.512355I$ $b = 0.944344 - 0.346033I$	$-1.03956 - 1.63645I$	$-7.74530 + 1.55839I$
$u = -0.944344 - 0.346033I$ $a = 0.447652 + 0.512355I$ $b = 0.944344 + 0.346033I$	$-1.03956 + 1.63645I$	$-7.74530 - 1.55839I$
$u = 0.864585 + 0.379164I$ $a = -0.583910 - 0.698557I$ $b = -0.864585 - 0.379164I$	$-2.10829 + 7.37198I$	$-8.93029 - 5.91248I$
$u = 0.864585 - 0.379164I$ $a = -0.583910 + 0.698557I$ $b = -0.864585 + 0.379164I$	$-2.10829 - 7.37198I$	$-8.93029 + 5.91248I$
$u = -0.037669 + 1.084830I$ $a = 0.11324 + 1.69832I$ $b = 0.037669 - 1.084830I$	$5.48501 - 2.53801I$	$5.54821 + 3.82197I$
$u = -0.037669 - 1.084830I$ $a = 0.11324 - 1.69832I$ $b = 0.037669 + 1.084830I$	$5.48501 + 2.53801I$	$5.54821 - 3.82197I$
$u = -0.878247$ $a = -0.337583$ $b = 0.878247$	-3.63803	-10.5770
$u = 0.791817 + 0.212409I$ $a = 0.104106 - 0.968399I$ $b = -0.791817 - 0.212409I$	$-7.18587 + 1.52748I$	$-14.7986 - 1.1989I$
$u = 0.791817 - 0.212409I$ $a = 0.104106 + 0.968399I$ $b = -0.791817 + 0.212409I$	$-7.18587 - 1.52748I$	$-14.7986 + 1.1989I$
$u = -0.714015 + 0.074924I$ $a = -1.185250 - 0.688415I$ $b = 0.714015 - 0.074924I$	$-3.09442 - 0.79036I$	$-5.12753 - 0.89506I$

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.714015 - 0.074924I$		
$a = -1.185250 + 0.688415I$	$-3.09442 + 0.79036I$	$-5.12753 + 0.89506I$
$b = 0.714015 + 0.074924I$		
$u = 0.704468 + 0.121688I$		
$a = 1.00603 - 1.10552I$	$-3.93940 - 4.16488I$	$-8.44169 + 5.58918I$
$b = -0.704468 - 0.121688I$		
$u = 0.704468 - 0.121688I$		
$a = 1.00603 + 1.10552I$	$-3.93940 + 4.16488I$	$-8.44169 - 5.58918I$
$b = -0.704468 + 0.121688I$		
$u = -0.145937 + 1.315070I$		
$a = 0.223886 + 1.159350I$	$2.99124 + 0.77564I$	$-5.07814 + 0.88442I$
$b = 0.145937 - 1.315070I$		
$u = -0.145937 - 1.315070I$		
$a = 0.223886 - 1.159350I$	$2.99124 - 0.77564I$	$-5.07814 - 0.88442I$
$b = 0.145937 + 1.315070I$		
$u = -0.07978 + 1.85790I$		
$a = 0.043035 + 0.691466I$	$9.06537 + 3.10101I$	$-4.13830 - 3.69699I$
$b = 0.07978 - 1.85790I$		
$u = -0.07978 - 1.85790I$		
$a = 0.043035 - 0.691466I$	$9.06537 - 3.10101I$	$-4.13830 + 3.69699I$
$b = 0.07978 + 1.85790I$		

$$\text{III. } I_3^u = \langle 1.91 \times 10^{76} u^{35} + 2.50 \times 10^{76} u^{34} + \dots + 5.35 \times 10^{78} b - 7.63 \times 10^{78}, 4.20 \times 10^{66} u^{35} + 2.74 \times 10^{66} u^{34} + \dots + 1.04 \times 10^{69} a - 1.38 \times 10^{70}, u^{36} + u^{35} + \dots - 2560u + 121 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_1 &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_4 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_5 &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_{10} &= \begin{pmatrix} -0.00403497u^{35} - 0.00263353u^{34} + \dots - 21.5564u + 13.2691 \\ -0.00356648u^{35} - 0.00468188u^{34} + \dots - 4.09637u + 1.42748 \end{pmatrix} \\ a_6 &= \begin{pmatrix} -0.00793734u^{35} - 0.00974330u^{34} + \dots - 11.1664u + 16.8486 \\ 0.00140145u^{35} + 0.000977635u^{34} + \dots + 2.93952u + 1.48823 \end{pmatrix} \\ a_9 &= \begin{pmatrix} -0.00760145u^{35} - 0.00731541u^{34} + \dots - 25.6527u + 14.6965 \\ -0.00356648u^{35} - 0.00468188u^{34} + \dots - 4.09637u + 1.42748 \end{pmatrix} \\ a_8 &= \begin{pmatrix} -0.00760145u^{35} - 0.00731541u^{34} + \dots - 25.6527u + 14.6965 \\ -0.00295648u^{35} - 0.00407097u^{34} + \dots - 2.44432u + 1.39287 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} 0.00422949u^{35} + 0.00563094u^{34} + \dots + 7.10480u - 7.88797 \\ 0.00138215u^{35} + 0.000407795u^{34} + \dots + 8.54688u - 1.12999 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 0.00561164u^{35} + 0.00603873u^{34} + \dots + 15.6517u - 9.01796 \\ 0.00138215u^{35} + 0.000407795u^{34} + \dots + 8.54688u - 1.12999 \end{pmatrix} \\ a_3 &= \begin{pmatrix} 0.00402228u^{35} + 0.00505346u^{34} + \dots + 1.24783u - 5.21219 \\ -0.00145650u^{35} - 0.0000469624u^{34} + \dots - 9.38769u - 0.107969 \end{pmatrix} \\ a_2 &= \begin{pmatrix} -0.00308517u^{35} - 0.00518618u^{34} + \dots + 5.15203u - 1.65668 \\ 0.00104331u^{35} + 0.000744287u^{34} + \dots + 5.57828u - 0.357559 \end{pmatrix} \\ a_7 &= \begin{pmatrix} -0.00764445u^{35} - 0.00742781u^{34} + \dots - 19.3477u + 13.7010 \\ 0.00208970u^{35} + 0.00142347u^{34} + \dots + 4.84956u + 1.05281 \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = -1

$$\text{(iii) Cusp Shapes} = -0.0117322u^{35} - 0.0177224u^{34} + \dots + 5.37839u + 0.170272$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_6	$(u^9 + 3u^8 + 8u^7 + 13u^6 + 17u^5 + 17u^4 + 12u^3 + 6u^2 + u - 1)^4$
c_2, c_7	$(u^9 - u^8 + 2u^7 - u^6 + 3u^5 - u^4 + 2u^3 + u + 1)^4$
c_3, c_5, c_{10} c_{11}	$u^{36} + u^{35} + \dots - 356u - 31$
c_4, c_9	$u^{36} - u^{35} + \dots + 2560u + 121$
c_8	$(u^9 + 5u^8 + 12u^7 + 15u^6 + 9u^5 - u^4 - 4u^3 - 2u^2 + u + 1)^4$
c_{12}	$(u^2 - u - 1)^{18}$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_6	$(y^9 + 7y^8 + 20y^7 + 25y^6 + 5y^5 - 15y^4 + 22y^2 + 13y - 1)^4$
c_2, c_7	$(y^9 + 3y^8 + 8y^7 + 13y^6 + 17y^5 + 17y^4 + 12y^3 + 6y^2 + y - 1)^4$
c_3, c_5, c_{10} c_{11}	$y^{36} - 13y^{35} + \dots - 49856y + 961$
c_4, c_9	$y^{36} + 23y^{35} + \dots - 5714344y + 14641$
c_8	$(y^9 - y^8 + 12y^7 - 7y^6 + 37y^5 + y^4 - 10y^2 + 5y - 1)^4$
c_{12}	$(y^2 - 3y + 1)^{18}$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.780953 + 0.638783I$		
$a = 0.474153 - 0.387835I$	$-3.33090 + 2.45442I$	$-5.67208 - 2.91298I$
$b = -1.009880 + 0.558025I$		
$u = 0.780953 - 0.638783I$		
$a = 0.474153 + 0.387835I$	$-3.33090 - 2.45442I$	$-5.67208 + 2.91298I$
$b = -1.009880 - 0.558025I$		
$u = 0.132026 + 0.843379I$		
$a = -0.29315 + 1.87262I$	$4.56478 + 2.45442I$	$-5.67208 - 2.91298I$
$b = -0.044584 - 1.300520I$		
$u = 0.132026 - 0.843379I$		
$a = -0.29315 - 1.87262I$	$4.56478 - 2.45442I$	$-5.67208 + 2.91298I$
$b = -0.044584 + 1.300520I$		
$u = -1.009880 + 0.558025I$		
$a = -0.468838 - 0.259064I$	$-3.33090 + 2.45442I$	$-5.67208 - 2.91298I$
$b = 0.780953 + 0.638783I$		
$u = -1.009880 - 0.558025I$		
$a = -0.468838 + 0.259064I$	$-3.33090 - 2.45442I$	$-5.67208 + 2.91298I$
$b = 0.780953 - 0.638783I$		
$u = -0.358853 + 0.748546I$		
$a = -0.321847 - 0.671353I$	$-0.34972 - 7.08493I$	$-2.42320 + 5.91335I$
$b = -1.53825 + 0.06109I$		
$u = -0.358853 - 0.748546I$		
$a = -0.321847 + 0.671353I$	$-0.34972 + 7.08493I$	$-2.42320 - 5.91335I$
$b = -1.53825 - 0.06109I$		
$u = 0.417017 + 0.614890I$		
$a = 0.466909 - 0.688456I$	$0.423507 + 1.336170I$	$-0.715907 - 0.701750I$
$b = 1.48241 + 0.01866I$		
$u = 0.417017 - 0.614890I$		
$a = 0.466909 + 0.688456I$	$0.423507 - 1.336170I$	$-0.715907 + 0.701750I$
$b = 1.48241 - 0.01866I$		

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.28654$ $a = 0.480385$ $b = 0.0498474$	-2.74940	0.652350
$u = -0.044584 + 1.300520I$ $a = 0.042601 + 1.242690I$ $b = 0.132026 - 0.843379I$	$4.56478 - 2.45442I$	$-5.67208 + 2.91298I$
$u = -0.044584 - 1.300520I$ $a = 0.042601 - 1.242690I$ $b = 0.132026 + 0.843379I$	$4.56478 + 2.45442I$	$-5.67208 - 2.91298I$
$u = 0.491968 + 1.251860I$ $a = -0.439989 + 1.119590I$ $b = -0.01428 - 1.56727I$	$2.16441 - 2.09337I$	$-8.51499 + 4.16283I$
$u = 0.491968 - 1.251860I$ $a = -0.439989 - 1.119590I$ $b = -0.01428 + 1.56727I$	$2.16441 + 2.09337I$	$-8.51499 - 4.16283I$
$u = -1.357600 + 0.226779I$ $a = -0.442882 - 0.073981I$ $b = 0.106990 + 0.598986I$	$-5.73128 - 2.09337I$	$-8.51499 + 4.16283I$
$u = -1.357600 - 0.226779I$ $a = -0.442882 + 0.073981I$ $b = 0.106990 - 0.598986I$	$-5.73128 + 2.09337I$	$-8.51499 - 4.16283I$
$u = 0.106990 + 0.598986I$ $a = 0.178600 - 0.999899I$ $b = -1.357600 + 0.226779I$	$-5.73128 - 2.09337I$	$-8.51499 + 4.16283I$
$u = 0.106990 - 0.598986I$ $a = 0.178600 + 0.999899I$ $b = -1.357600 - 0.226779I$	$-5.73128 + 2.09337I$	$-8.51499 - 4.16283I$
$u = 1.48241 + 0.01866I$ $a = 0.416845 - 0.005247I$ $b = 0.417017 + 0.614890I$	$0.423507 + 1.336170I$	$-0.715907 - 0.701750I$

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.48241 - 0.01866I$ $a = 0.416845 + 0.005247I$ $b = 0.417017 - 0.614890I$	$0.423507 - 1.336170I$	$-0.715907 + 0.701750I$
$u = -0.25523 + 1.49501I$ $a = 0.179535 + 1.051640I$ $b = -0.25523 - 1.49501I$	5.14629	$-60.652349 + 0.10I$
$u = -0.25523 - 1.49501I$ $a = 0.179535 - 1.051640I$ $b = -0.25523 + 1.49501I$	5.14629	$-60.652349 + 0.10I$
$u = -1.53825 + 0.06109I$ $a = -0.401145 - 0.015931I$ $b = -0.358853 + 0.748546I$	$-0.34972 - 7.08493I$	$-2.42320 + 5.91335I$
$u = -1.53825 - 0.06109I$ $a = -0.401145 + 0.015931I$ $b = -0.358853 - 0.748546I$	$-0.34972 + 7.08493I$	$-2.42320 - 5.91335I$
$u = -0.01428 + 1.56727I$ $a = 0.009404 + 1.032300I$ $b = 0.491968 - 1.251860I$	$2.16441 + 2.09337I$	$-8.51499 - 4.16283I$
$u = -0.01428 - 1.56727I$ $a = 0.009404 - 1.032300I$ $b = 0.491968 + 1.251860I$	$2.16441 - 2.09337I$	$-8.51499 + 4.16283I$
$u = 0.68303 + 1.50001I$ $a = -0.406822 + 0.893434I$ $b = 0.04160 - 1.80927I$	$7.54597 - 7.08493I$	$-2.42320 + 5.91335I$
$u = 0.68303 - 1.50001I$ $a = -0.406822 - 0.893434I$ $b = 0.04160 + 1.80927I$	$7.54597 + 7.08493I$	$-2.42320 - 5.91335I$
$u = -0.61176 + 1.54868I$ $a = 0.357003 + 0.903756I$ $b = -0.11375 - 1.79068I$	$8.31919 + 1.33617I$	0

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.61176 - 1.54868I$ $a = 0.357003 - 0.903756I$ $b = -0.11375 + 1.79068I$	$8.31919 - 1.33617I$	0
$u = -0.11375 + 1.79068I$ $a = 0.057170 + 0.899955I$ $b = -0.61176 - 1.54868I$	$8.31919 - 1.33617I$	0
$u = -0.11375 - 1.79068I$ $a = 0.057170 - 0.899955I$ $b = -0.61176 + 1.54868I$	$8.31919 + 1.33617I$	0
$u = 0.04160 + 1.80927I$ $a = -0.020553 + 0.893830I$ $b = 0.68303 - 1.50001I$	$7.54597 + 7.08493I$	$-4.00000 - 5.91335I$
$u = 0.04160 - 1.80927I$ $a = -0.020553 - 0.893830I$ $b = 0.68303 + 1.50001I$	$7.54597 - 7.08493I$	$-4.00000 + 5.91335I$
$u = 0.0498474$ $a = 12.3985$ $b = 1.28654$	-2.74940	0.652350

IV. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1, c_6	$(u^9 + 3u^8 + 8u^7 + 13u^6 + 17u^5 + 17u^4 + 12u^3 + 6u^2 + u - 1)^4$ $\cdot (u^{17} - 6u^{16} + \dots - 18u^2 + 1)(u^{28} + 9u^{27} + \dots + 84u + 16)$
c_2	$(u^9 - u^8 + 2u^7 - u^6 + 3u^5 - u^4 + 2u^3 + u + 1)^4$ $\cdot (u^{17} + 3u^{15} + \dots + 2u - 1)(u^{28} + 5u^{27} + \dots + 2u + 4)$
c_3, c_{10}	$(u^{17} - u^{16} + \dots + 3u^2 + 1)(u^{28} + u^{27} + \dots + 4u + 1)$ $\cdot (u^{36} + u^{35} + \dots - 356u - 31)$
c_4, c_9	$(u^{17} + 3u^{15} + \dots - u + 1)(u^{28} + 23u^{26} + \dots + 3u + 1)$ $\cdot (u^{36} - u^{35} + \dots + 2560u + 121)$
c_5, c_{11}	$(u^{17} + u^{16} + \dots - 3u^2 - 1)(u^{28} + u^{27} + \dots + 4u + 1)$ $\cdot (u^{36} + u^{35} + \dots - 356u - 31)$
c_7	$(u^9 - u^8 + 2u^7 - u^6 + 3u^5 - u^4 + 2u^3 + u + 1)^4$ $\cdot (u^{17} + 3u^{15} + \dots + 2u + 1)(u^{28} + 5u^{27} + \dots + 2u + 4)$
c_8	$(u^9 + 5u^8 + 12u^7 + 15u^6 + 9u^5 - u^4 - 4u^3 - 2u^2 + u + 1)^4$ $\cdot (u^{17} - u^{15} + \dots + 2u + 1)(u^{28} - 25u^{27} + \dots - 98862u + 9028)$
c_{12}	$((u^2 - u - 1)^{18})(u^{17} - 6u^{16} + \dots + 2u + 1)$ $\cdot (u^{28} + 21u^{27} + \dots + 3584u + 512)$

V. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1, c_6	$(y^9 + 7y^8 + 20y^7 + 25y^6 + 5y^5 - 15y^4 + 22y^2 + 13y - 1)^4$ $\cdot (y^{17} + 14y^{16} + \dots + 36y - 1)(y^{28} + 21y^{27} + \dots + 2064y + 256)$
c_2, c_7	$(y^9 + 3y^8 + 8y^7 + 13y^6 + 17y^5 + 17y^4 + 12y^3 + 6y^2 + y - 1)^4$ $\cdot (y^{17} + 6y^{16} + \dots + 18y^2 - 1)(y^{28} + 9y^{27} + \dots + 84y + 16)$
c_3, c_5, c_{10} c_{11}	$(y^{17} - 11y^{16} + \dots - 6y - 1)(y^{28} - 11y^{27} + \dots - 6y + 1)$ $\cdot (y^{36} - 13y^{35} + \dots - 49856y + 961)$
c_4, c_9	$(y^{17} + 6y^{16} + \dots + 11y - 1)(y^{28} + 46y^{27} + \dots + 49y + 1)$ $\cdot (y^{36} + 23y^{35} + \dots - 5714344y + 14641)$
c_8	$(y^9 - y^8 + 12y^7 - 7y^6 + 37y^5 + y^4 - 10y^2 + 5y - 1)^4$ $\cdot (y^{17} - 2y^{16} + \dots - 2y - 1)$ $\cdot (y^{28} - 3y^{27} + \dots + 785869044y + 81504784)$
c_{12}	$((y^2 - 3y + 1)^{18})(y^{17} - 14y^{16} + \dots + 8y - 1)$ $\cdot (y^{28} - 17y^{27} + \dots + 4456448y + 262144)$