



11a₃₁₆ (K11a₃₁₆)



A knot diagram¹

Linearized knot diagram



Solving Sequence

$$4,10 \xrightarrow{c_{10}} 11 \xrightarrow{c_4} 5,8 \xrightarrow{c_5} 6 \xrightarrow{c_{11}} 1 \xrightarrow{c_3} 3 \xrightarrow{c_2} 2 \xrightarrow{c_7} 7 \xrightarrow{c_9} 9 \longrightarrow c_1, c_6, c_8$$

Ideals for irreducible components² of X_{par}

$$I_1^u = \langle 1.80623 \times 10^{32} u^{61} - 5.14761 \times 10^{32} u^{60} + \dots + 4.61017 \times 10^{32} b - 1.10416 \times 10^{32}, \\ - 6.91332 \times 10^{32} u^{61} - 7.53652 \times 10^{32} u^{60} + \dots + 2.30509 \times 10^{33} a - 2.31132 \times 10^{33}, u^{62} - 2u^{61} + \dots + 3u \rangle$$

$$I_2^u = \langle b - 1, 5a + u - 2, u^2 + u - 1 \rangle$$

* 2 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 64 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/maths/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

I.

$$I_1^u = \langle 1.81 \times 10^{32} u^{61} - 5.15 \times 10^{32} u^{60} + \dots + 4.61 \times 10^{32} b - 1.10 \times 10^{32}, -6.91 \times 10^{32} u^{61} - 7.54 \times 10^{32} u^{60} + \dots + 2.31 \times 10^{33} a - 2.31 \times 10^{33}, u^{62} - 2u^{61} + \dots + 3u + 1 \rangle$$

(i) Arc colorings

$$a_4 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 0.299916u^{61} + 0.326952u^{60} + \dots - 3.67552u + 1.00270 \\ -0.391793u^{61} + 1.11658u^{60} + \dots - 0.782541u + 0.239505 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -1.37364u^{61} + 0.955447u^{60} + \dots + 15.8721u + 3.35880 \\ 0.226500u^{61} - 0.438184u^{60} + \dots + 5.26406u + 0.574836 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -u^2 + 1 \\ -u^4 + 2u^2 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} u^5 - 2u^3 + u \\ u^7 - 3u^5 + 2u^3 + u \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 0.403303u^{61} + 1.11034u^{60} + \dots + 3.59385u - 1.62323 \\ 0.435823u^{61} - 0.0545467u^{60} + \dots - 0.0437745u - 1.01330 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} 0.691708u^{61} - 0.789625u^{60} + \dots - 2.89298u + 0.763197 \\ -0.391793u^{61} + 1.11658u^{60} + \dots - 0.782541u + 0.239505 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 0.210043u^{61} + 0.259417u^{60} + \dots - 3.20318u + 1.25041 \\ -0.348888u^{61} + 0.656606u^{60} + \dots + 0.331553u + 0.825406 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 0.210043u^{61} + 0.259417u^{60} + \dots - 3.20318u + 1.25041 \\ -0.348888u^{61} + 0.656606u^{60} + \dots + 0.331553u + 0.825406 \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes = $3.89183u^{61} - 9.68250u^{60} + \dots - 4.68341u + 13.0328$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$5(5u^{62} + 7u^{61} + \dots + 100038u - 28017)$
c_2, c_6	$u^{62} + 2u^{61} + \dots - u + 1$
c_3	$u^{62} - 6u^{61} + \dots + 795u - 117$
c_4, c_{10}, c_{11}	$u^{62} + 2u^{61} + \dots - 3u + 1$
c_5	$5(5u^{62} - 14u^{61} + \dots + 61755u + 8377)$
c_7, c_9	$u^{62} + 3u^{61} + \dots + 214u - 25$
c_8	$u^{62} + u^{61} + \dots + 280u - 100$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$25(25y^{62} - 1309y^{61} + \dots - 1.54576 \times 10^{10}y + 7.84952 \times 10^8)$
c_2, c_6	$y^{62} + 42y^{61} + \dots - 25y + 1$
c_3	$y^{62} + 18y^{61} + \dots - 440613y + 13689$
c_4, c_{10}, c_{11}	$y^{62} - 54y^{61} + \dots - 25y + 1$
c_5	$25(25y^{62} - 286y^{61} + \dots - 1.29244 \times 10^9y + 7.01741 \times 10^7)$
c_7, c_9	$y^{62} - 49y^{61} + \dots - 22596y + 625$
c_8	$y^{62} - 15y^{61} + \dots - 413000y + 10000$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.887619 + 0.514849I$ $a = -0.168834 - 0.676664I$ $b = -1.351070 - 0.186432I$	$6.81266 + 5.56177I$	$4.66163 - 5.86539I$
$u = -0.887619 - 0.514849I$ $a = -0.168834 + 0.676664I$ $b = -1.351070 + 0.186432I$	$6.81266 - 5.56177I$	$4.66163 + 5.86539I$
$u = -0.956988 + 0.457534I$ $a = -0.411794 - 0.142908I$ $b = -1.44265 + 0.39962I$	$7.18631 - 7.01551I$	0
$u = -0.956988 - 0.457534I$ $a = -0.411794 + 0.142908I$ $b = -1.44265 - 0.39962I$	$7.18631 + 7.01551I$	0
$u = -0.268927 + 0.858943I$ $a = 0.487450 - 0.791896I$ $b = -1.326860 + 0.042366I$	$8.77447 - 0.73219I$	$8.26642 + 0.24787I$
$u = -0.268927 - 0.858943I$ $a = 0.487450 + 0.791896I$ $b = -1.326860 - 0.042366I$	$8.77447 + 0.73219I$	$8.26642 - 0.24787I$
$u = 0.981110 + 0.511433I$ $a = -0.156001 + 0.318273I$ $b = -1.245360 - 0.167604I$	$2.26101 + 1.03760I$	0
$u = 0.981110 - 0.511433I$ $a = -0.156001 - 0.318273I$ $b = -1.245360 + 0.167604I$	$2.26101 - 1.03760I$	0
$u = 0.216176 + 0.854810I$ $a = 0.879762 + 0.839368I$ $b = -1.284170 + 0.305083I$	$4.62237 - 5.84398I$	$3.48358 + 6.22366I$
$u = 0.216176 - 0.854810I$ $a = 0.879762 - 0.839368I$ $b = -1.284170 - 0.305083I$	$4.62237 + 5.84398I$	$3.48358 - 6.22366I$

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.222629 + 0.830394I$ $a = 0.97517 - 1.10506I$ $b = -1.47365 - 0.48988I$	$9.4690 + 11.6120I$	$5.02297 - 7.15875I$
$u = -0.222629 - 0.830394I$ $a = 0.97517 + 1.10506I$ $b = -1.47365 + 0.48988I$	$9.4690 - 11.6120I$	$5.02297 + 7.15875I$
$u = -1.133930 + 0.168026I$ $a = 1.26618 - 0.92935I$ $b = 0.586572 - 1.058950I$	$1.22471 - 2.22328I$	0
$u = -1.133930 - 0.168026I$ $a = 1.26618 + 0.92935I$ $b = 0.586572 + 1.058950I$	$1.22471 + 2.22328I$	0
$u = 1.233700 + 0.140829I$ $a = 1.11356 + 0.95442I$ $b = 0.498573 + 0.517622I$	$-2.34840 - 0.48442I$	0
$u = 1.233700 - 0.140829I$ $a = 1.11356 - 0.95442I$ $b = 0.498573 - 0.517622I$	$-2.34840 + 0.48442I$	0
$u = 1.228560 + 0.275128I$ $a = 1.155710 - 0.633770I$ $b = 1.77170 + 0.45104I$	$4.49072 - 1.35832I$	0
$u = 1.228560 - 0.275128I$ $a = 1.155710 + 0.633770I$ $b = 1.77170 - 0.45104I$	$4.49072 + 1.35832I$	0
$u = -0.154976 + 0.717888I$ $a = -0.542160 + 0.195736I$ $b = 0.278710 + 1.257500I$	$3.96361 + 5.62230I$	$4.59110 - 7.16807I$
$u = -0.154976 - 0.717888I$ $a = -0.542160 - 0.195736I$ $b = 0.278710 - 1.257500I$	$3.96361 - 5.62230I$	$4.59110 + 7.16807I$

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.046542 + 0.724748I$ $a = -1.208990 - 0.366912I$ $b = 1.66460 - 0.64188I$	$8.09797 - 2.26515I$	$10.09980 + 3.27421I$
$u = 0.046542 - 0.724748I$ $a = -1.208990 + 0.366912I$ $b = 1.66460 + 0.64188I$	$8.09797 + 2.26515I$	$10.09980 - 3.27421I$
$u = -1.263000 + 0.236260I$ $a = -0.487348 + 0.065309I$ $b = 1.210290 - 0.144045I$	$-0.11388 + 2.17095I$	0
$u = -1.263000 - 0.236260I$ $a = -0.487348 - 0.065309I$ $b = 1.210290 + 0.144045I$	$-0.11388 - 2.17095I$	0
$u = -1.288710 + 0.185153I$ $a = 2.28058 - 2.74209I$ $b = 0.864085 + 0.005366I$	$0.03983 + 2.79266I$	0
$u = -1.288710 - 0.185153I$ $a = 2.28058 + 2.74209I$ $b = 0.864085 - 0.005366I$	$0.03983 - 2.79266I$	0
$u = 0.179463 + 0.661396I$ $a = -0.055951 - 0.244113I$ $b = 0.050410 - 0.678849I$	$0.49280 - 2.22863I$	$-1.88873 + 4.27642I$
$u = 0.179463 - 0.661396I$ $a = -0.055951 + 0.244113I$ $b = 0.050410 + 0.678849I$	$0.49280 + 2.22863I$	$-1.88873 - 4.27642I$
$u = -1.293980 + 0.298379I$ $a = 0.30932 + 2.31869I$ $b = 1.60267 + 0.81560I$	$3.91657 + 5.96640I$	0
$u = -1.293980 - 0.298379I$ $a = 0.30932 - 2.31869I$ $b = 1.60267 - 0.81560I$	$3.91657 - 5.96640I$	0

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.301670 + 0.265672I$ $a = -0.14106 - 2.23111I$ $b = 1.118260 - 0.376395I$	$-0.57816 - 4.41700I$	0
$u = 1.301670 - 0.265672I$ $a = -0.14106 + 2.23111I$ $b = 1.118260 + 0.376395I$	$-0.57816 + 4.41700I$	0
$u = -0.042843 + 0.662429I$ $a = -1.81923 + 1.14582I$ $b = 1.139070 + 0.251694I$	$3.63336 + 1.04794I$	$2.34672 - 0.35139I$
$u = -0.042843 - 0.662429I$ $a = -1.81923 - 1.14582I$ $b = 1.139070 - 0.251694I$	$3.63336 - 1.04794I$	$2.34672 + 0.35139I$
$u = 1.334310 + 0.246440I$ $a = 1.069770 - 0.893080I$ $b = 0.231740 - 0.094385I$	$-1.00664 - 2.96825I$	0
$u = 1.334310 - 0.246440I$ $a = 1.069770 + 0.893080I$ $b = 0.231740 + 0.094385I$	$-1.00664 + 2.96825I$	0
$u = -0.115133 + 0.619794I$ $a = 1.22726 + 1.09164I$ $b = 0.413434 - 0.098972I$	$3.55015 - 0.20409I$	$5.98628 - 1.19975I$
$u = -0.115133 - 0.619794I$ $a = 1.22726 - 1.09164I$ $b = 0.413434 + 0.098972I$	$3.55015 + 0.20409I$	$5.98628 + 1.19975I$
$u = 1.372160 + 0.035845I$ $a = 0.47764 + 1.70860I$ $b = -0.237289 + 0.862746I$	$-4.26849 + 2.33228I$	0
$u = 1.372160 - 0.035845I$ $a = 0.47764 - 1.70860I$ $b = -0.237289 - 0.862746I$	$-4.26849 - 2.33228I$	0

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.354390 + 0.299440I$ $a = -0.88402 - 1.77945I$ $b = 0.150670 - 1.364490I$	$-0.80126 - 9.32026I$	0
$u = 1.354390 - 0.299440I$ $a = -0.88402 + 1.77945I$ $b = 0.150670 + 1.364490I$	$-0.80126 + 9.32026I$	0
$u = -1.364120 + 0.277183I$ $a = -0.491592 + 1.171990I$ $b = -0.043760 + 0.847969I$	$-4.39333 + 5.67407I$	0
$u = -1.364120 - 0.277183I$ $a = -0.491592 - 1.171990I$ $b = -0.043760 - 0.847969I$	$-4.39333 - 5.67407I$	0
$u = -1.41802 + 0.08070I$ $a = -0.165441 - 1.216540I$ $b = -0.566676 - 0.567301I$	$-7.01139 + 2.01626I$	0
$u = -1.41802 - 0.08070I$ $a = -0.165441 + 1.216540I$ $b = -0.566676 + 0.567301I$	$-7.01139 - 2.01626I$	0
$u = 1.39937 + 0.34656I$ $a = -0.28769 + 2.13431I$ $b = -1.47028 + 0.55396I$	$4.3271 - 15.8582I$	0
$u = 1.39937 - 0.34656I$ $a = -0.28769 - 2.13431I$ $b = -1.47028 - 0.55396I$	$4.3271 + 15.8582I$	0
$u = -1.39672 + 0.35868I$ $a = -0.16352 - 1.69873I$ $b = -1.286260 - 0.404733I$	$-0.48145 + 10.21060I$	0
$u = -1.39672 - 0.35868I$ $a = -0.16352 + 1.69873I$ $b = -1.286260 + 0.404733I$	$-0.48145 - 10.21060I$	0

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.463287 + 0.297906I$		
$a = 0.788454 + 0.539829I$	$-1.008560 - 0.666322I$	$-7.37810 + 3.95165I$
$b = -0.250407 + 0.311008I$		
$u = 0.463287 - 0.297906I$		
$a = 0.788454 - 0.539829I$	$-1.008560 + 0.666322I$	$-7.37810 - 3.95165I$
$b = -0.250407 - 0.311008I$		
$u = -0.550158 + 0.016824I$		
$a = 1.33228 + 1.00368I$	$1.45921 + 2.58594I$	$-1.55168 - 3.98149I$
$b = 0.256229 + 0.735207I$		
$u = -0.550158 - 0.016824I$		
$a = 1.33228 - 1.00368I$	$1.45921 - 2.58594I$	$-1.55168 + 3.98149I$
$b = 0.256229 - 0.735207I$		
$u = 1.42814 + 0.36734I$		
$a = -0.564087 + 1.146080I$	$3.38499 - 3.70672I$	0
$b = -1.270380 + 0.067319I$		
$u = 1.42814 - 0.36734I$		
$a = -0.564087 - 1.146080I$	$3.38499 + 3.70672I$	0
$b = -1.270380 - 0.067319I$		
$u = 1.50001 + 0.04160I$		
$a = -1.29195 + 0.81714I$	$-1.16526 - 6.77623I$	0
$b = -1.205330 + 0.382950I$		
$u = 1.50001 - 0.04160I$		
$a = -1.29195 - 0.81714I$	$-1.16526 + 6.77623I$	0
$b = -1.205330 - 0.382950I$		
$u = 0.239018 + 0.279647I$		
$a = 3.12607 - 0.97399I$	$4.30759 - 0.97665I$	$-0.516718 + 1.197110I$
$b = 1.190070 - 0.194532I$		
$u = 0.239018 - 0.279647I$		
$a = 3.12607 + 0.97399I$	$4.30759 + 0.97665I$	$-0.516718 - 1.197110I$
$b = 1.190070 + 0.194532I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.63921$ $a = -0.771484$ $b = -1.04713$	-7.12923	0
$u = -0.201088$ $a = 2.67242$ $b = 0.901231$	1.30954	9.93960

$$\text{II. } I_2^u = \langle b - 1, 5a + u - 2, u^2 + u - 1 \rangle$$

(i) Arc colorings

$$a_4 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 1 \\ -u + 1 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -u \\ -u + 1 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} -\frac{1}{5}u + \frac{2}{5} \\ 1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -u + \frac{1}{5} \\ -\frac{4}{5}u + \frac{2}{5} \end{pmatrix}$$

$$a_1 = \begin{pmatrix} u \\ u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} 2u - 1 \\ 3u - 1 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} \frac{4}{5}u \\ \frac{3}{5}u - \frac{1}{5} \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -\frac{1}{5}u - \frac{3}{5} \\ 1 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -\frac{1}{5}u + \frac{2}{5} \\ 1 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -\frac{1}{5}u + \frac{2}{5} \\ 1 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = $\frac{72}{5}u - 13$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$5(5u^2 - 1)$
c_2, c_{10}, c_{11}	$u^2 + u - 1$
c_3	$u^2 + 3u + 1$
c_4, c_6	$u^2 - u - 1$
c_5	$5(5u^2 + 5u + 1)$
c_7	$(u + 1)^2$
c_8	u^2
c_9	$(u - 1)^2$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$25(5y - 1)^2$
c_2, c_4, c_6 c_{10}, c_{11}	$y^2 - 3y + 1$
c_3	$y^2 - 7y + 1$
c_5	$25(25y^2 - 15y + 1)$
c_7, c_9	$(y - 1)^2$
c_8	y^2

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.618034$ $a = 0.276393$ $b = 1.00000$	0.657974	-4.10030
$u = -1.61803$ $a = 0.723607$ $b = 1.00000$	-7.23771	-36.3000

III. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1	$25(5u^2 - 1)(5u^{62} + 7u^{61} + \dots + 100038u - 28017)$
c_2	$(u^2 + u - 1)(u^{62} + 2u^{61} + \dots - u + 1)$
c_3	$(u^2 + 3u + 1)(u^{62} - 6u^{61} + \dots + 795u - 117)$
c_4	$(u^2 - u - 1)(u^{62} + 2u^{61} + \dots - 3u + 1)$
c_5	$25(5u^2 + 5u + 1)(5u^{62} - 14u^{61} + \dots + 61755u + 8377)$
c_6	$(u^2 - u - 1)(u^{62} + 2u^{61} + \dots - u + 1)$
c_7	$((u + 1)^2)(u^{62} + 3u^{61} + \dots + 214u - 25)$
c_8	$u^2(u^{62} + u^{61} + \dots + 280u - 100)$
c_9	$((u - 1)^2)(u^{62} + 3u^{61} + \dots + 214u - 25)$
c_{10}, c_{11}	$(u^2 + u - 1)(u^{62} + 2u^{61} + \dots - 3u + 1)$

IV. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1	$625(5y - 1)^2 \cdot (25y^{62} - 1309y^{61} + \dots - 15457580352y + 784952289)$
c_2, c_6	$(y^2 - 3y + 1)(y^{62} + 42y^{61} + \dots - 25y + 1)$
c_3	$(y^2 - 7y + 1)(y^{62} + 18y^{61} + \dots - 440613y + 13689)$
c_4, c_{10}, c_{11}	$(y^2 - 3y + 1)(y^{62} - 54y^{61} + \dots - 25y + 1)$
c_5	$625(25y^2 - 15y + 1) \cdot (25y^{62} - 286y^{61} + \dots - 1292437581y + 70174129)$
c_7, c_9	$((y - 1)^2)(y^{62} - 49y^{61} + \dots - 22596y + 625)$
c_8	$y^2(y^{62} - 15y^{61} + \dots - 413000y + 10000)$