



10<sub>120</sub> (K10a<sub>102</sub>)



A knot diagram<sup>1</sup>

**Linearized knot diagram**



**Solving Sequence**

$$1,5 \xrightarrow{c_1} 2,8 \xrightarrow{c_5} 6 \xrightarrow{c_8} 9 \xrightarrow{c_4} 4 \xrightarrow{c_3} 3 \xrightarrow{c_7} 7 \xrightarrow{c_{10}} 10 \longrightarrow c_2, c_6, c_9$$

**Ideals for irreducible components<sup>2</sup> of  $X_{\text{par}}$**

$$\begin{aligned}
 I_1^u &= \langle -u^6 - u^5 - 2u^4 - u^3 - u^2 + b - u + 1, \\
 &\quad -u^{10} - 3u^9 - 8u^8 - 13u^7 - 19u^6 - 20u^5 - 19u^4 - 13u^3 - 7u^2 + 2a - 2u + 2, \\
 &\quad u^{11} + 3u^{10} + 8u^9 + 13u^8 + 19u^7 + 22u^6 + 21u^5 + 17u^4 + 9u^3 + 4u^2 - 2 \rangle \\
 I_2^u &= \langle -u^{17} - 7u^{16} + \dots + b - 13, -13u^{17} - 60u^{16} + \dots + 5a + 4, u^{18} + 5u^{17} + \dots + 27u + 5 \rangle \\
 I_3^u &= \langle -3u^{11} + 11u^{10} - 29u^9 + 50u^8 - 66u^7 + 71u^6 - 59u^5 + 43u^4 - 22u^3 - au + 7u^2 + b - 4u, \\
 &\quad 3u^{10}a + u^{11} + \dots + 4a - 4, u^{12} - 3u^{11} + 8u^{10} - 13u^9 + 18u^8 - 21u^7 + 19u^6 - 17u^5 + 10u^4 - 6u^3 + 4u^2 + 1 \rangle \\
 I_4^u &= \langle -au + u^2 + b - u + 1, -u^2a + a^2 + u^2 - a + 1, u^3 - u^2 + 2u - 1 \rangle \\
 I_5^u &= \langle u^2 + b - u + 1, u^2 + a + 1, u^3 - u^2 + 2u - 1 \rangle \\
 I_6^u &= \langle u^6 - 4u^5 + 8u^4 - 11u^3 + 9u^2 + b - 5u + 3, -3u^7 + 12u^6 - 25u^5 + 34u^4 - 29u^3 + 17u^2 + 2a - 9u + 2, \\
 &\quad u^8 - 4u^7 + 9u^6 - 14u^5 + 15u^4 - 13u^3 + 9u^2 - 4u + 2 \rangle \\
 I_7^u &= \langle -au + b + u - 1, a^2 - au - 1, u^2 - u + 1 \rangle
 \end{aligned}$$

\* 7 irreducible components of  $\dim_{\mathbb{C}} = 0$ , with total 74 representations.

<sup>1</sup>The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/maths/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

<sup>2</sup>All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$\mathbf{I. } I_1^u = \langle -u^6 - u^5 - 2u^4 - u^3 - u^2 + b - u + 1, -u^{10} - 3u^9 + \dots + 2a + 2, u^{11} + 3u^{10} + \dots + 4u^2 - 2 \rangle$$

(i) Arc colorings

$$a_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} \frac{1}{2}u^{10} + \frac{3}{2}u^9 + \dots + u - 1 \\ u^6 + u^5 + 2u^4 + u^3 + u^2 + u - 1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} \frac{1}{2}u^{10} + \frac{3}{2}u^9 + \dots - \frac{1}{2}u^2 - u \\ u^9 + 2u^8 + 5u^7 + 6u^6 + 8u^5 + 7u^4 + 5u^3 + 3u^2 + u - 1 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} \frac{1}{2}u^{10} + \frac{3}{2}u^9 + \dots + \frac{11}{2}u^3 + \frac{5}{2}u^2 \\ u^6 + u^5 + 2u^4 + u^3 + u^2 + u - 1 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u \\ u^3 + u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -\frac{1}{2}u^{10} - \frac{3}{2}u^9 + \dots + u + 1 \\ -u^6 - u^5 - 2u^4 - u^3 - u^2 - u + 1 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} \frac{1}{2}u^{10} + \frac{5}{2}u^9 + \dots + \frac{7}{2}u^2 - 1 \\ -u^{10} - 3u^9 - 7u^8 - 11u^7 - 14u^6 - 15u^5 - 12u^4 - 8u^3 - 3u^2 + 1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -\frac{1}{2}u^{10} - \frac{1}{2}u^9 + \dots + u - 1 \\ -u^9 - 2u^8 - 5u^7 - 6u^6 - 8u^5 - 7u^4 - 5u^3 - 3u^2 + 1 \end{pmatrix}$$

(ii) Obstruction class = -1

$$\mathbf{(iii) } \text{Cusp Shapes} = -2u^8 - 6u^7 - 14u^6 - 18u^5 - 24u^4 - 18u^3 - 16u^2 - 6u - 6$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_4, c_6$ $c_9$	$u^{11} - 3u^{10} + 8u^9 - 13u^8 + 19u^7 - 22u^6 + 21u^5 - 17u^4 + 9u^3 - 4u^2 + 2$
$c_2, c_7$	$u^{11} + 5u^{10} + \dots + 10u + 4$
$c_3, c_5, c_8$ $c_{10}$	$u^{11} + 4u^9 + u^8 + 11u^7 + 4u^6 + 15u^5 + 3u^4 + 9u^3 - u^2 + 2u + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_4, c_6$ $c_9$	$y^{11} + 7y^{10} + \dots + 16y - 4$
$c_2, c_7$	$y^{11} - 5y^{10} + \dots + 60y - 16$
$c_3, c_5, c_8$ $c_{10}$	$y^{11} + 8y^{10} + \dots + 6y - 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.955959 + 0.181916I$ $a = -0.517203 - 1.103780I$ $b = -0.695220 - 0.961079I$	$-0.51987 - 4.74721I$	$-10.74299 + 5.17166I$
$u = -0.955959 - 0.181916I$ $a = -0.517203 + 1.103780I$ $b = -0.695220 + 0.961079I$	$-0.51987 + 4.74721I$	$-10.74299 - 5.17166I$
$u = 0.104833 + 1.064770I$ $a = -1.051120 - 0.609326I$ $b = -0.538602 + 1.183080I$	$5.39093 - 0.52336I$	$-2.07555 - 0.88155I$
$u = 0.104833 - 1.064770I$ $a = -1.051120 + 0.609326I$ $b = -0.538602 - 1.183080I$	$5.39093 + 0.52336I$	$-2.07555 + 0.88155I$
$u = 0.375570 + 1.042270I$ $a = 0.289036 + 0.451507I$ $b = 0.362037 - 0.470824I$	$2.05520 - 3.23878I$	$-8.62571 + 3.68812I$
$u = 0.375570 - 1.042270I$ $a = 0.289036 - 0.451507I$ $b = 0.362037 + 0.470824I$	$2.05520 + 3.23878I$	$-8.62571 - 3.68812I$
$u = -0.641442 + 1.159660I$ $a = -0.736546 + 0.484569I$ $b = 0.089483 + 1.164960I$	$6.47745 + 4.30838I$	$-1.34168 - 3.93056I$
$u = -0.641442 - 1.159660I$ $a = -0.736546 - 0.484569I$ $b = 0.089483 - 1.164960I$	$6.47745 - 4.30838I$	$-1.34168 + 3.93056I$
$u = -0.58305 + 1.34141I$ $a = 1.128180 + 0.208445I$ $b = 0.93740 - 1.39182I$	$6.7235 + 16.2714I$	$-5.72276 - 8.85281I$
$u = -0.58305 - 1.34141I$ $a = 1.128180 - 0.208445I$ $b = 0.93740 + 1.39182I$	$6.7235 - 16.2714I$	$-5.72276 + 8.85281I$

Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.400093$		
$a = 0.775290$	$-0.775978$	$-12.9830$
$b = -0.310188$		

$$\langle -u^{17} - 7u^{16} + \dots + b - 13, -13u^{17} - 60u^{16} + \dots + 5a + 4, u^{18} + 5u^{17} + \dots + 27u + 5 \rangle$$

II.  $I_2^u =$

(i) Arc colorings

$$a_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} \frac{13}{5}u^{17} + 12u^{16} + \dots + \frac{77}{5}u - \frac{4}{5} \\ u^{17} + 7u^{16} + \dots + 71u + 13 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -\frac{11}{5}u^{17} - 11u^{16} + \dots - \frac{229}{5}u - \frac{42}{5} \\ -3u^{15} - 13u^{14} + \dots - 50u - 11 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} \frac{8}{5}u^{17} + 5u^{16} + \dots - \frac{278}{5}u - \frac{69}{5} \\ u^{17} + 7u^{16} + \dots + 71u + 13 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u \\ u^3 + u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} \frac{4}{5}u^{17} + 8u^{16} + \dots + \frac{391}{5}u + \frac{73}{5} \\ -3u^{17} - 16u^{16} + \dots - 62u - 11 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} \frac{13}{5}u^{17} + 13u^{16} + \dots + \frac{67}{5}u - \frac{4}{5} \\ u^{16} + 5u^{15} + \dots + 42u + 8 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} \frac{9}{5}u^{17} + 6u^{16} + \dots - \frac{189}{5}u - \frac{32}{5} \\ -u^{17} - 4u^{16} + \dots + 16u + 4 \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes

$$= 26u^{17} + 130u^{16} + 415u^{15} + 871u^{14} + 1236u^{13} + 1002u^{12} - 275u^{11} - 2187u^{10} - 3562u^9 - 3091u^8 - 569u^7 + 2567u^6 + 4603u^5 + 4614u^4 + 3181u^3 + 1560u^2 + 486u + 63$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_4, c_6$ $c_9$	$u^{18} - 5u^{17} + \dots - 27u + 5$
$c_2, c_7$	$(u^9 - 2u^8 + 4u^7 - 5u^6 + 7u^5 - 5u^4 + 3u^3 - 2u^2 + u - 1)^2$
$c_3, c_5, c_8$ $c_{10}$	$u^{18} - u^{17} + \dots + 3u + 1$



(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_4, c_6$ $c_9$	$y^{18} + 9y^{17} + \dots + 61y + 25$
$c_2, c_7$	$(y^9 + 4y^8 + 10y^7 + 17y^6 + 17y^5 + y^4 - 7y^3 - 8y^2 - 3y - 1)^2$
$c_3, c_5, c_8$ $c_{10}$	$y^{18} + 9y^{17} + \dots + 43y + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_2^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.198527 + 0.827118I$ $a = -0.81586 - 1.28949I$ $b = -1.228530 + 0.418813I$	$-1.04793 + 1.11007I$	$-17.5535 - 4.7867I$
$u = -0.198527 - 0.827118I$ $a = -0.81586 + 1.28949I$ $b = -1.228530 - 0.418813I$	$-1.04793 - 1.11007I$	$-17.5535 + 4.7867I$
$u = -0.079308 + 0.836177I$ $a = 1.38823 + 0.30834I$ $b = 0.367922 - 1.136350I$	4.23983	$-2.85705 + 0.I$
$u = -0.079308 - 0.836177I$ $a = 1.38823 - 0.30834I$ $b = 0.367922 + 1.136350I$	4.23983	$-2.85705 + 0.I$
$u = -1.156420 + 0.102576I$ $a = 0.521985 + 0.890705I$ $b = 0.694998 + 0.976484I$	$2.81259 - 10.16840I$	$-7.74812 + 7.64867I$
$u = -1.156420 - 0.102576I$ $a = 0.521985 - 0.890705I$ $b = 0.694998 - 0.976484I$	$2.81259 + 10.16840I$	$-7.74812 - 7.64867I$
$u = 1.160130 + 0.229157I$ $a = 0.035605 - 0.158300I$ $b = -0.077582 + 0.175489I$	$-1.04793 - 1.11007I$	$-17.5535 + 4.7867I$
$u = 1.160130 - 0.229157I$ $a = 0.035605 + 0.158300I$ $b = -0.077582 - 0.175489I$	$-1.04793 + 1.11007I$	$-17.5535 - 4.7867I$
$u = -0.311796 + 1.205210I$ $a = 0.814403 - 0.074315I$ $b = 0.164362 - 1.004700I$	$4.26456 - 0.69984I$	$-4.65022 + 1.89978I$
$u = -0.311796 - 1.205210I$ $a = 0.814403 + 0.074315I$ $b = 0.164362 + 1.004700I$	$4.26456 + 0.69984I$	$-4.65022 - 1.89978I$

Solutions to $I_2^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.369880 + 1.229190I$ $a = 1.216830 + 0.459753I$ $b = 1.01520 - 1.32566I$	$8.30021 + 4.38855I$	$-1.11965 - 3.68700I$
$u = -0.369880 - 1.229190I$ $a = 1.216830 - 0.459753I$ $b = 1.01520 + 1.32566I$	$8.30021 - 4.38855I$	$-1.11965 + 3.68700I$
$u = -0.642487 + 0.199869I$ $a = 1.21195 - 1.14348I$ $b = 0.550112 - 0.976904I$	$4.26456 + 0.69984I$	$-4.65022 - 1.89978I$
$u = -0.642487 - 0.199869I$ $a = 1.21195 + 1.14348I$ $b = 0.550112 + 0.976904I$	$4.26456 - 0.69984I$	$-4.65022 + 1.89978I$
$u = -0.545158 + 1.253180I$ $a = -1.229130 - 0.230487I$ $b = -0.95891 + 1.41466I$	$2.81259 + 10.16840I$	$-7.74812 - 7.64867I$
$u = -0.545158 - 1.253180I$ $a = -1.229130 + 0.230487I$ $b = -0.95891 - 1.41466I$	$2.81259 - 10.16840I$	$-7.74812 + 7.64867I$
$u = -0.35655 + 1.50992I$ $a = -0.544015 + 0.110198I$ $b = -0.027580 + 0.860709I$	$8.30021 - 4.38855I$	$-1.11965 + 3.68700I$
$u = -0.35655 - 1.50992I$ $a = -0.544015 - 0.110198I$ $b = -0.027580 - 0.860709I$	$8.30021 + 4.38855I$	$-1.11965 - 3.68700I$

$$\langle -3u^{11} + 11u^{10} + \dots + b - 4u, 3u^{10}a + u^{11} + \dots + 4a - 4, u^{12} - 3u^{11} + \dots + 4u^2 + 1 \rangle$$

III.  $I_3^u =$

(i) Arc colorings

$$a_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} a \\ 3u^{11} - 11u^{10} + \dots + au + 4u \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 3u^{11}a - u^{11} + \dots - 4u - 1 \\ 1 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -3u^{11} + 11u^{10} + \dots + a - 4u \\ 3u^{11} - 11u^{10} + \dots + au + 4u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u \\ u^3 + u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -2u^{11}a + u^{11} + \dots - a + 2 \\ -u^{10}a + 4u^9a + \dots - 2a + 1 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -u^{11} + 5u^{10} + \dots + a + 2 \\ 2u^{11} - 7u^{10} + \dots + au + 2u \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -u^{10}a - u^{11} + \dots + a - 2 \\ -u^7a + u^6a + \dots + a + 1 \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes

$$= 4u^{11} - 16u^{10} + 44u^9 - 80u^8 + 112u^7 - 124u^6 + 116u^5 - 92u^4 + 60u^3 - 28u^2 + 8u - 18$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_4, c_6$ $c_9$	$(u^{12} + 3u^{11} + \dots + 4u^2 + 1)^2$
$c_2, c_7$	$(u^{12} + u^{10} - 6u^9 + 10u^8 - 2u^7 + 2u^6 - 2u^4 + 2u^2 + 1)^2$
$c_3, c_5, c_8$ $c_{10}$	$u^{24} - 3u^{23} + \dots - 4u^2 + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_4, c_6$ $c_9$	$(y^{12} + 7y^{11} + \dots + 8y + 1)^2$
$c_2, c_7$	$(y^{12} + 2y^{11} + \dots + 4y + 1)^2$
$c_3, c_5, c_8$ $c_{10}$	$y^{24} - 11y^{23} + \dots - 8y + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_3^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.234552 + 1.002020I$ $a = 0.647681 + 0.298955I$ $b = 2.11054 + 0.48664I$	$3.72285 + 6.96551I$	$-5.97171 - 10.57440I$
$u = -0.234552 + 1.002020I$ $a = 0.00700 + 2.10465I$ $b = 0.451474 - 0.578868I$	$3.72285 + 6.96551I$	$-5.97171 - 10.57440I$
$u = -0.234552 - 1.002020I$ $a = 0.647681 - 0.298955I$ $b = 2.11054 - 0.48664I$	$3.72285 - 6.96551I$	$-5.97171 + 10.57440I$
$u = -0.234552 - 1.002020I$ $a = 0.00700 - 2.10465I$ $b = 0.451474 + 0.578868I$	$3.72285 - 6.96551I$	$-5.97171 + 10.57440I$
$u = 1.090290 + 0.140460I$ $a = -0.240626 - 0.291610I$ $b = -0.401743 + 0.003834I$	$-1.05784 - 1.08263I$	$-14.2815 + 5.6276I$
$u = 1.090290 + 0.140460I$ $a = 0.362012 - 0.050153I$ $b = 0.221393 + 0.351739I$	$-1.05784 - 1.08263I$	$-14.2815 + 5.6276I$
$u = 1.090290 - 0.140460I$ $a = -0.240626 + 0.291610I$ $b = -0.401743 - 0.003834I$	$-1.05784 + 1.08263I$	$-14.2815 - 5.6276I$
$u = 1.090290 - 0.140460I$ $a = 0.362012 + 0.050153I$ $b = 0.221393 - 0.351739I$	$-1.05784 + 1.08263I$	$-14.2815 - 5.6276I$
$u = -0.185688 + 0.817666I$ $a = -0.762192 - 0.903819I$ $b = -1.44128 + 0.18321I$	$-1.05784 + 1.08263I$	$-14.2815 - 5.6276I$
$u = -0.185688 + 0.817666I$ $a = -0.59374 - 1.62783I$ $b = -0.880553 + 0.455390I$	$-1.05784 + 1.08263I$	$-14.2815 - 5.6276I$

Solutions to $I_3^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.185688 - 0.817666I$		
$a = -0.762192 + 0.903819I$	$-1.05784 - 1.08263I$	$-14.2815 + 5.6276I$
$b = -1.44128 - 0.18321I$		
$u = -0.185688 - 0.817666I$		
$a = -0.59374 + 1.62783I$	$-1.05784 - 1.08263I$	$-14.2815 + 5.6276I$
$b = -0.880553 - 0.455390I$		
$u = 0.529049 + 1.245360I$		
$a = 0.970902 - 0.051810I$	$2.26979 - 4.55813I$	$-9.74676 + 1.77049I$
$b = 0.692981 + 0.737589I$		
$u = 0.529049 + 1.245360I$		
$a = -0.701975 + 0.258240I$	$2.26979 - 4.55813I$	$-9.74676 + 1.77049I$
$b = -0.578176 - 1.181710I$		
$u = 0.529049 - 1.245360I$		
$a = 0.970902 + 0.051810I$	$2.26979 + 4.55813I$	$-9.74676 - 1.77049I$
$b = 0.692981 - 0.737589I$		
$u = 0.529049 - 1.245360I$		
$a = -0.701975 - 0.258240I$	$2.26979 + 4.55813I$	$-9.74676 - 1.77049I$
$b = -0.578176 + 1.181710I$		
$u = -0.251512 + 0.449740I$		
$a = 0.04323 + 2.16308I$	$2.26979 - 4.55813I$	$-9.74676 + 1.77049I$
$b = 0.800711 - 0.884208I$		
$u = -0.251512 + 0.449740I$		
$a = 2.25611 + 0.51868I$	$2.26979 - 4.55813I$	$-9.74676 + 1.77049I$
$b = 0.983696 + 0.524600I$		
$u = -0.251512 - 0.449740I$		
$a = 0.04323 - 2.16308I$	$2.26979 + 4.55813I$	$-9.74676 - 1.77049I$
$b = 0.800711 + 0.884208I$		
$u = -0.251512 - 0.449740I$		
$a = 2.25611 - 0.51868I$	$2.26979 + 4.55813I$	$-9.74676 - 1.77049I$
$b = 0.983696 - 0.524600I$		



Solutions to $I_3^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.55241 + 1.40748I$	$3.72285 - 6.96551I$	$-5.97171 + 10.57440I$
$a = 0.767735 - 0.370784I$		
$b = 0.486934 + 1.037080I$		
$u = 0.55241 + 1.40748I$	$3.72285 - 6.96551I$	$-5.97171 + 10.57440I$
$a = -0.756136 + 0.049190I$		
$b = -0.945979 - 0.875748I$		
$u = 0.55241 - 1.40748I$	$3.72285 + 6.96551I$	$-5.97171 - 10.57440I$
$a = 0.767735 + 0.370784I$		
$b = 0.486934 - 1.037080I$		
$u = 0.55241 - 1.40748I$	$3.72285 + 6.96551I$	$-5.97171 - 10.57440I$
$a = -0.756136 - 0.049190I$		
$b = -0.945979 + 0.875748I$		

$$\text{IV. } I_4^u = \langle -au + u^2 + b - u + 1, -u^2a + a^2 + u^2 - a + 1, u^3 - u^2 + 2u - 1 \rangle$$

(i) Arc colorings

$$a_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} a \\ au - u^2 + u - 1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -u^2a + au + u^2 - a - u + 1 \\ 1 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -au + u^2 + a - u + 1 \\ au - u^2 + u - 1 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u \\ u^2 - u + 1 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} 2au - a \\ -au + u^2 - u + 1 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} u^2a - au + u^2 + a - u \\ -u^2a + au - 2u^2 + u - 1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} u^2 + u \\ -au + a - 2u + 1 \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes =  $-8u^2 + 8u - 14$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_4, c_6$ $c_9$	$(u^3 + u^2 + 2u + 1)^2$
$c_2, c_7$	$u^6 + 5u^5 + 16u^4 + 28u^3 + 30u^2 + 18u + 5$
$c_3, c_5, c_8$ $c_{10}$	$u^6 - u^5 + 2u^4 + 2u^3 + 4u^2 + 2u + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_4, c_6$ $c_9$	$(y^3 + 3y^2 + 2y - 1)^2$
$c_2, c_7$	$y^6 + 7y^5 + 36y^4 + 6y^3 + 52y^2 - 24y + 25$
$c_3, c_5, c_8$ $c_{10}$	$y^6 + 3y^5 + 16y^4 + 18y^3 + 12y^2 + 4y + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_4^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.215080 + 1.307140I$ $a = 0.594305 - 0.123240I$ $b = 1.16635 + 1.49520I$	$7.69319 - 5.65624I$	$1.01951 + 5.95889I$
$u = 0.215080 + 1.307140I$ $a = -1.25666 + 0.68552I$ $b = -0.288915 - 0.750335I$	$7.69319 - 5.65624I$	$1.01951 + 5.95889I$
$u = 0.215080 - 1.307140I$ $a = 0.594305 + 0.123240I$ $b = 1.16635 - 1.49520I$	$7.69319 + 5.65624I$	$1.01951 - 5.95889I$
$u = 0.215080 - 1.307140I$ $a = -1.25666 - 0.68552I$ $b = -0.288915 + 0.750335I$	$7.69319 + 5.65624I$	$1.01951 - 5.95889I$
$u = 0.569840$ $a = 0.662359 + 0.941275I$ $b = -0.377439 + 0.536376I$	$-0.581975$	$-12.0390$
$u = 0.569840$ $a = 0.662359 - 0.941275I$ $b = -0.377439 - 0.536376I$	$-0.581975$	$-12.0390$

$$\mathbf{V. } I_5^u = \langle u^2 + b - u + 1, u^2 + a + 1, u^3 - u^2 + 2u - 1 \rangle$$

**(i) Arc colorings**

$$a_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} -u^2 - 1 \\ -u^2 + u - 1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -u \\ -u^2 + u - 1 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u \\ u^2 - u + 1 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} u^2 + 1 \\ u^2 - u + 1 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -1 \\ -u^2 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 0 \\ -u \end{pmatrix}$$

**(ii) Obstruction class = 1**

**(iii) Cusp Shapes =  $-8u^2 + 8u - 20$**

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_6$	$u^3 - u^2 + 2u - 1$
$c_2, c_4, c_7$ $c_9$	$u^3 + u^2 + 2u + 1$
$c_3, c_5, c_8$ $c_{10}$	$u^3 + u^2 - 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_2, c_4$ $c_6, c_7, c_9$	$y^3 + 3y^2 + 2y - 1$
$c_3, c_5, c_8$ $c_{10}$	$y^3 - y^2 + 2y - 1$



(vi) Complex Volumes and Cusp Shapes

Solutions to $I_5^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.215080 + 1.307140I$ $a = 0.662359 - 0.562280I$ $b = 0.877439 + 0.744862I$	$6.04826 - 5.65624I$	$-4.98049 + 5.95889I$
$u = 0.215080 - 1.307140I$ $a = 0.662359 + 0.562280I$ $b = 0.877439 - 0.744862I$	$6.04826 + 5.65624I$	$-4.98049 - 5.95889I$
$u = 0.569840$ $a = -1.32472$ $b = -0.754878$	$-2.22691$	$-18.0390$

$$\text{VI. } I_6^u = \langle u^6 - 4u^5 + 8u^4 - 11u^3 + 9u^2 + b - 5u + 3, -3u^7 + 12u^6 + \dots + 2a + 2, u^8 - 4u^7 + \dots - 4u + 2 \rangle$$

(i) Arc colorings

$$a_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} \frac{3}{2}u^7 - 6u^6 + \dots + \frac{9}{2}u - 1 \\ -u^6 + 4u^5 - 8u^4 + 11u^3 - 9u^2 + 5u - 3 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} \frac{1}{2}u^7 - 3u^6 + \dots + \frac{15}{2}u - 4 \\ -u^7 + 3u^6 - 5u^5 + 6u^4 - 4u^3 + 3u^2 - u - 1 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} \frac{3}{2}u^7 - 5u^6 + \dots - \frac{1}{2}u + 2 \\ -u^6 + 4u^5 - 8u^4 + 11u^3 - 9u^2 + 5u - 3 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u \\ u^3 + u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -\frac{1}{2}u^7 + 2u^6 + \dots - \frac{7}{2}u + 1 \\ u^2 - u + 1 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} \frac{3}{2}u^7 - 6u^6 + \dots + \frac{3}{2}u + 1 \\ -u^7 + 2u^6 - 2u^5 + 3u^3 - 2u^2 + 2u - 1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} \frac{1}{2}u^7 - 2u^6 + \dots + \frac{3}{2}u + 1 \\ -u^4 + 2u^3 - 3u^2 + 2u - 1 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes =  $-3u^7 + 16u^6 - 43u^5 + 71u^4 - 82u^3 + 68u^2 - 40u + 20$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_6$	$u^8 - 4u^7 + 9u^6 - 14u^5 + 15u^4 - 13u^3 + 9u^2 - 4u + 2$
$c_2, c_7$	$(u^4 - u^3 + u^2 + 1)^2$
$c_3, c_5, c_8$ $c_{10}$	$u^8 - 2u^7 + 3u^5 - 3u^4 + 3u^2 - u + 1$
$c_4, c_9$	$u^8 + 4u^7 + 9u^6 + 14u^5 + 15u^4 + 13u^3 + 9u^2 + 4u + 2$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_4, c_6$ $c_9$	$y^8 + 2y^7 - y^6 - 12y^5 - 5y^4 + 25y^3 + 37y^2 + 20y + 4$
$c_2, c_7$	$(y^4 + y^3 + 3y^2 + 2y + 1)^2$
$c_3, c_5, c_8$ $c_{10}$	$y^8 - 4y^7 + 6y^6 - 3y^5 + 7y^4 - 12y^3 + 3y^2 + 5y + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_6^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.192965 + 0.870342I$ $a = -0.81301 + 1.44822I$ $b = -1.41733 - 0.42814I$	$-0.732875 - 0.991478I$	$5.28161 - 3.59996I$
$u = 0.192965 - 0.870342I$ $a = -0.81301 - 1.44822I$ $b = -1.41733 + 0.42814I$	$-0.732875 + 0.991478I$	$5.28161 + 3.59996I$
$u = -0.138557 + 0.767522I$ $a = 0.066843 - 1.409780I$ $b = 1.072770 + 0.246639I$	$3.20028 + 5.62938I$	$-5.78161 - 5.27851I$
$u = -0.138557 - 0.767522I$ $a = 0.066843 + 1.409780I$ $b = 1.072770 - 0.246639I$	$3.20028 - 5.62938I$	$-5.78161 + 5.27851I$
$u = 1.354460 + 0.250532I$ $a = 0.008624 + 0.392991I$ $b = -0.086775 + 0.534450I$	$-0.732875 - 0.991478I$	$5.28161 - 3.59996I$
$u = 1.354460 - 0.250532I$ $a = 0.008624 - 0.392991I$ $b = -0.086775 - 0.534450I$	$-0.732875 + 0.991478I$	$5.28161 + 3.59996I$
$u = 0.59113 + 1.35317I$ $a = -0.762459 + 0.087166I$ $b = -0.568666 - 0.980213I$	$3.20028 - 5.62938I$	$-5.78161 + 5.27851I$
$u = 0.59113 - 1.35317I$ $a = -0.762459 - 0.087166I$ $b = -0.568666 + 0.980213I$	$3.20028 + 5.62938I$	$-5.78161 - 5.27851I$

$$\text{VII. } I_7^u = \langle -au + b + u - 1, a^2 - au - 1, u^2 - u + 1 \rangle$$

(i) Arc colorings

$$a_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 1 \\ u - 1 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} a \\ au - u + 1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -au + a - u \\ 1 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -au + a + u - 1 \\ au - u + 1 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u \\ u - 1 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -au + a + u \\ au \end{pmatrix}$$

$$a_7 = \begin{pmatrix} a + u \\ au \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -2au + a \\ a - u + 1 \end{pmatrix}$$

(ii) Obstruction class =  $-1$

(iii) Cusp Shapes =  $8u - 14$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_4, c_6$ $c_9$	$(u^2 + u + 1)^2$
$c_2, c_7$	$(u - 1)^4$
$c_3, c_5, c_8$ $c_{10}$	$u^4 - u^3 + 2u^2 - 2u + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_4, c_6$ $c_9$	$(y^2 + y + 1)^2$
$c_2, c_7$	$(y - 1)^4$
$c_3, c_5, c_8$ $c_{10}$	$y^4 + 3y^3 + 2y^2 + 1$



(vi) Complex Volumes and Cusp Shapes

Solutions to $I_7^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.500000 + 0.866025I$ $a = -0.692440 + 0.318148I$ $b = -0.121744 - 1.306620I$	$1.64493 - 4.05977I$	$-10.00000 + 6.92820I$
$u = 0.500000 + 0.866025I$ $a = 1.192440 + 0.547877I$ $b = 0.621744 + 0.440597I$	$1.64493 - 4.05977I$	$-10.00000 + 6.92820I$
$u = 0.500000 - 0.866025I$ $a = -0.692440 - 0.318148I$ $b = -0.121744 + 1.306620I$	$1.64493 + 4.05977I$	$-10.00000 - 6.92820I$
$u = 0.500000 - 0.866025I$ $a = 1.192440 - 0.547877I$ $b = 0.621744 - 0.440597I$	$1.64493 + 4.05977I$	$-10.00000 - 6.92820I$

### VIII. u-Polynomials

Crossings	u-Polynomials at each crossing
$c_1, c_6$	$(u^2 + u + 1)^2(u^3 - u^2 + 2u - 1)(u^3 + u^2 + 2u + 1)^2$ $\cdot (u^8 - 4u^7 + 9u^6 - 14u^5 + 15u^4 - 13u^3 + 9u^2 - 4u + 2)$ $\cdot (u^{11} - 3u^{10} + 8u^9 - 13u^8 + 19u^7 - 22u^6 + 21u^5 - 17u^4 + 9u^3 - 4u^2 + 2)$ $\cdot ((u^{12} + 3u^{11} + \dots + 4u^2 + 1)^2)(u^{18} - 5u^{17} + \dots - 27u + 5)$
$c_2, c_7$	$(u - 1)^4(u^3 + u^2 + 2u + 1)(u^4 - u^3 + u^2 + 1)^2$ $\cdot (u^6 + 5u^5 + 16u^4 + 28u^3 + 30u^2 + 18u + 5)$ $\cdot (u^9 - 2u^8 + 4u^7 - 5u^6 + 7u^5 - 5u^4 + 3u^3 - 2u^2 + u - 1)^2$ $\cdot (u^{11} + 5u^{10} + \dots + 10u + 4)$ $\cdot (u^{12} + u^{10} - 6u^9 + 10u^8 - 2u^7 + 2u^6 - 2u^4 + 2u^2 + 1)^2$
$c_3, c_5, c_8$ $c_{10}$	$(u^3 + u^2 - 1)(u^4 - u^3 + 2u^2 - 2u + 1)(u^6 - u^5 + \dots + 2u + 1)$ $\cdot (u^8 - 2u^7 + 3u^5 - 3u^4 + 3u^2 - u + 1)$ $\cdot (u^{11} + 4u^9 + u^8 + 11u^7 + 4u^6 + 15u^5 + 3u^4 + 9u^3 - u^2 + 2u + 1)$ $\cdot (u^{18} - u^{17} + \dots + 3u + 1)(u^{24} - 3u^{23} + \dots - 4u^2 + 1)$
$c_4, c_9$	$(u^2 + u + 1)^2(u^3 + u^2 + 2u + 1)^3$ $\cdot (u^8 + 4u^7 + 9u^6 + 14u^5 + 15u^4 + 13u^3 + 9u^2 + 4u + 2)$ $\cdot (u^{11} - 3u^{10} + 8u^9 - 13u^8 + 19u^7 - 22u^6 + 21u^5 - 17u^4 + 9u^3 - 4u^2 + 2)$ $\cdot ((u^{12} + 3u^{11} + \dots + 4u^2 + 1)^2)(u^{18} - 5u^{17} + \dots - 27u + 5)$

### IX. Riley Polynomials

Crossings	Riley Polynomials at each crossing
$c_1, c_4, c_6$ $c_9$	$(y^2 + y + 1)^2(y^3 + 3y^2 + 2y - 1)^3$ $\cdot (y^8 + 2y^7 - y^6 - 12y^5 - 5y^4 + 25y^3 + 37y^2 + 20y + 4)$ $\cdot (y^{11} + 7y^{10} + \dots + 16y - 4)(y^{12} + 7y^{11} + \dots + 8y + 1)^2$ $\cdot (y^{18} + 9y^{17} + \dots + 61y + 25)$
$c_2, c_7$	$(y - 1)^4(y^3 + 3y^2 + 2y - 1)(y^4 + y^3 + 3y^2 + 2y + 1)^2$ $\cdot (y^6 + 7y^5 + 36y^4 + 6y^3 + 52y^2 - 24y + 25)$ $\cdot (y^9 + 4y^8 + 10y^7 + 17y^6 + 17y^5 + y^4 - 7y^3 - 8y^2 - 3y - 1)^2$ $\cdot (y^{11} - 5y^{10} + \dots + 60y - 16)(y^{12} + 2y^{11} + \dots + 4y + 1)^2$
$c_3, c_5, c_8$ $c_{10}$	$(y^3 - y^2 + 2y - 1)(y^4 + 3y^3 + 2y^2 + 1)$ $\cdot (y^6 + 3y^5 + 16y^4 + 18y^3 + 12y^2 + 4y + 1)$ $\cdot (y^8 - 4y^7 + 6y^6 - 3y^5 + 7y^4 - 12y^3 + 3y^2 + 5y + 1)$ $\cdot (y^{11} + 8y^{10} + \dots + 6y - 1)(y^{18} + 9y^{17} + \dots + 43y + 1)$ $\cdot (y^{24} - 11y^{23} + \dots - 8y + 1)$