



12a<sub>0426</sub> (K12a<sub>0426</sub>)



A knot diagram<sup>1</sup>

**Linearized knot diagram**



**Solving Sequence**

$$7, 11 \xrightarrow{c_7} 5, 8 \xrightarrow{c_8} 9 \xrightarrow{c_{11}} 12 \xrightarrow{c_4} 4 \xrightarrow{c_{12}} 1 \xrightarrow{c_6} 6 \xrightarrow{c_{10}} 10 \xrightarrow{c_3} 3 \xrightarrow{c_2} 2 \rightsquigarrow c_1, c_5, c_9$$

**Ideals for irreducible components<sup>2</sup> of  $X_{\text{par}}$**

$$I_1^u = \langle 5u^{17} - 11u^{16} + \dots + 3b - 11, u^{17} + u^{16} + \dots + 2a + 4, u^{18} - 3u^{17} + \dots - 6u + 2 \rangle$$

$$I_2^u = \langle 2.93297 \times 10^{19}u^{49} - 3.08421 \times 10^{20}u^{48} + \dots + 4.46875 \times 10^{17}b - 5.17754 \times 10^{20}, \\ - 2.95147 \times 10^{19}u^{49} + 6.16285 \times 10^{20}u^{48} + \dots + 5.80937 \times 10^{18}a + 6.41388 \times 10^{21}, \\ u^{50} - 11u^{49} + \dots - 165u + 13 \rangle$$

$$I_3^u = \langle -u^5a - u^5 + 2u^3a + u^3 - 2au + b - u + 1, 2u^4a + u^3a - 3u^4 - 2u^2a - 2u^3 + a^2 - 2au + u^2 + 3u + 2, \\ u^6 + u^5 - u^4 - 2u^3 + u + 1 \rangle$$

$$I_4^u = \langle u^5 - u^3 + b + u, u^3 + a, u^6 + u^5 - u^4 - 2u^3 + u + 1 \rangle$$

$$I_5^u = \langle -2u^{43}a - 4u^{42}a + \dots + 4b - 65, 50u^{43}a - 95u^{43} + \dots - 468a + 943, u^{44} + 3u^{43} + \dots - 14u - 1 \rangle$$

$$I_6^u = \langle 3u^{15} + 5u^{14} - 4u^{13} - 18u^{12} - 11u^{11} + 4u^{10} + 4u^9 + u^8 + 21u^7 + 32u^6 + 22u^5 + 9u^4 + 10u^3 + 5u^2 + b - u - \\ - 7u^{15} - 28u^{14} + \dots + 2a - 22, u^{16} + 4u^{15} + \dots + 8u + 2 \rangle$$

$$I_7^u = \langle -u^2a + b - 1, a^2 + 2au + 2u^2 + a + 3u + 2, u^3 + u^2 - 1 \rangle$$

$$I_8^u = \langle -u^3a - u^2a - u^3 + au + 2b + 1, -u^2a + u^3 + a^2 + au - 2u^2 + a - u + 2, u^4 - u^2 + 1 \rangle$$

$$I_9^u = \langle -71757a^7 + 7931089b + \dots - 9811314a + 3141666, \\ a^8 + 6a^6 - 6a^5 + 26a^4 - 18a^3 + 87a^2 - 60a + 73, u - 1 \rangle$$

\* 9 irreducible components of  $\dim_{\mathbb{C}} = 0$ , with total 212 representations.

<sup>1</sup>The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew (<http://www.layer8.co.uk/maths/draw/index.htm#Running-draw>), where we modified some parts for our purpose (<https://github.com/CATsTAILs/LinksPainter>).

<sup>2</sup>All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated

I.

$$I_1^u = \langle 5u^{17} - 11u^{16} + \dots + 3b - 11, u^{17} + u^{16} + \dots + 2a + 4, u^{18} - 3u^{17} + \dots - 6u + 2 \rangle$$

(i) Arc colorings

$$a_7 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -\frac{1}{2}u^{17} - \frac{1}{2}u^{16} + \dots - \frac{3}{2}u^2 - 2 \\ -\frac{5}{3}u^{17} + \frac{11}{3}u^{16} + \dots - \frac{23}{3}u + \frac{11}{3} \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} \frac{1}{2}u^{17} - \frac{5}{2}u^{16} + \dots + 5u - 3 \\ -\frac{5}{3}u^{17} + \frac{11}{3}u^{16} + \dots - \frac{23}{3}u + \frac{11}{3} \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} \frac{7}{6}u^{17} - \frac{25}{6}u^{16} + \dots + \frac{23}{3}u - \frac{17}{3} \\ -\frac{5}{3}u^{17} + \frac{11}{3}u^{16} + \dots - \frac{23}{3}u + \frac{11}{3} \end{pmatrix}$$

$$a_1 = \begin{pmatrix} \frac{19}{6}u^{17} - \frac{37}{6}u^{16} + \dots + \frac{32}{3}u - \frac{14}{3} \\ -\frac{2}{3}u^{17} + \frac{2}{3}u^{16} + \dots + \frac{1}{3}u - \frac{1}{3} \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -\frac{7}{6}u^{17} + \frac{25}{6}u^{16} + \dots - \frac{23}{3}u + \frac{17}{3} \\ \frac{2}{3}u^{17} - \frac{2}{3}u^{16} + \dots + \frac{2}{3}u + \frac{1}{3} \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} u^3 \\ u^5 - u^3 + u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -\frac{5}{2}u^{17} + \frac{7}{2}u^{16} + \dots - 7u + 1 \\ -3.33333u^{17} + 7.33333u^{16} + \dots - 13.3333u + 6.33333 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -1.83333u^{17} + 3.83333u^{16} + \dots - 8.33333u + 3.33333 \\ -2u^{17} + 5u^{16} + \dots - 9u + 5 \end{pmatrix}$$

(ii) Obstruction class = -1

$$(iii) \text{ Cusp Shapes} = -8u^{17} + 14u^{16} + 12u^{15} - 50u^{14} - 2u^{13} + 94u^{12} - 40u^{11} - 130u^{10} + 112u^9 + 86u^8 - 140u^7 - 22u^6 + 106u^5 - 20u^4 - 36u^3 + 24u^2 - 14u + 6$$

---

in decimal forms when there is not enough margin.

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_{10}$	$u^{18} + 7u^{17} + \dots - 4u + 4$
$c_2, c_5, c_7$ $c_{11}$	$u^{18} + 3u^{17} + \dots + 6u + 2$
$c_3, c_8$	$u^{18} - 9u^{17} + \dots - 16u + 8$
$c_4, c_6, c_9$ $c_{12}$	$u^{18} + 5u^{16} + \dots + u + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_{10}$	$y^{18} + 13y^{17} + \dots - 272y + 16$
$c_2, c_5, c_7$ $c_{11}$	$y^{18} - 7y^{17} + \dots + 4y + 4$
$c_3, c_8$	$y^{18} - 9y^{17} + \dots - 480y + 64$
$c_4, c_6, c_9$ $c_{12}$	$y^{18} + 10y^{17} + \dots + y + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.929188 + 0.427848I$ $a = 1.45891 + 0.96005I$ $b = -0.612958 + 0.694927I$	$-2.52393 - 4.60971I$	$-3.54545 + 10.13031I$
$u = 0.929188 - 0.427848I$ $a = 1.45891 - 0.96005I$ $b = -0.612958 - 0.694927I$	$-2.52393 + 4.60971I$	$-3.54545 - 10.13031I$
$u = -0.880851 + 0.560750I$ $a = -0.83772 + 1.88459I$ $b = -0.21871 + 1.48927I$	$-1.22474 + 4.46780I$	$-2.01264 - 5.91058I$
$u = -0.880851 - 0.560750I$ $a = -0.83772 - 1.88459I$ $b = -0.21871 - 1.48927I$	$-1.22474 - 4.46780I$	$-2.01264 + 5.91058I$
$u = 0.582973 + 0.940039I$ $a = -0.268767 - 0.004991I$ $b = -1.09278 - 1.05818I$	$5.24090 + 8.44951I$	$3.89928 - 4.02582I$
$u = 0.582973 - 0.940039I$ $a = -0.268767 + 0.004991I$ $b = -1.09278 + 1.05818I$	$5.24090 - 8.44951I$	$3.89928 + 4.02582I$
$u = 0.829747 + 0.331402I$ $a = -0.543141 - 1.062640I$ $b = 0.506778 + 0.049880I$	$-1.42597 - 1.89978I$	$-0.27303 + 2.78097I$
$u = 0.829747 - 0.331402I$ $a = -0.543141 + 1.062640I$ $b = 0.506778 - 0.049880I$	$-1.42597 + 1.89978I$	$-0.27303 - 2.78097I$
$u = -1.174310 + 0.030732I$ $a = 0.73424 + 1.60100I$ $b = 0.540765 + 1.144710I$	$-8.44772 + 5.23335I$	$-9.78688 - 5.32435I$
$u = -1.174310 - 0.030732I$ $a = 0.73424 - 1.60100I$ $b = 0.540765 - 1.144710I$	$-8.44772 - 5.23335I$	$-9.78688 + 5.32435I$

Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.933002 + 0.761436I$		
$a = 0.373106 - 0.772716I$	$4.32570 + 5.86053I$	$5.28493 - 4.51200I$
$b = 0.113921 - 0.679581I$		
$u = -0.933002 - 0.761436I$		
$a = 0.373106 + 0.772716I$	$4.32570 - 5.86053I$	$5.28493 + 4.51200I$
$b = 0.113921 + 0.679581I$		
$u = 0.969925 + 0.853431I$		
$a = 0.717351 + 0.406552I$	$1.61741 - 6.44223I$	$-7.21121 + 5.93194I$
$b = 0.005204 + 1.085600I$		
$u = 0.969925 - 0.853431I$		
$a = 0.717351 - 0.406552I$	$1.61741 + 6.44223I$	$-7.21121 - 5.93194I$
$b = 0.005204 - 1.085600I$		
$u = 1.120310 + 0.718189I$		
$a = -0.54198 - 1.83396I$	$1.8659 - 20.6729I$	$-0.25702 + 11.54258I$
$b = 1.13133 - 1.27977I$		
$u = 1.120310 - 0.718189I$		
$a = -0.54198 + 1.83396I$	$1.8659 + 20.6729I$	$-0.25702 - 11.54258I$
$b = 1.13133 + 1.27977I$		
$u = 0.056018 + 0.547940I$		
$a = -0.591992 - 0.492924I$	$0.572438 - 1.277810I$	$3.90202 + 5.68169I$
$b = -0.373557 + 0.433371I$		
$u = 0.056018 - 0.547940I$		
$a = -0.591992 + 0.492924I$	$0.572438 + 1.277810I$	$3.90202 - 5.68169I$
$b = -0.373557 - 0.433371I$		

$$\text{II. } I_2^u = \langle 2.93 \times 10^{19} u^{49} - 3.08 \times 10^{20} u^{48} + \dots + 4.47 \times 10^{17} b - 5.18 \times 10^{20}, -2.95 \times 10^{19} u^{49} + 6.16 \times 10^{20} u^{48} + \dots + 5.81 \times 10^{18} a + 6.41 \times 10^{21}, u^{50} - 11u^{49} + \dots - 165u + 13 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_7 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_5 &= \begin{pmatrix} 5.08053u^{49} - 106.085u^{48} + \dots + 11433.4u - 1104.06 \\ -65.6329u^{49} + 690.174u^{48} + \dots - 13362.0u + 1158.61 \end{pmatrix} \\ a_8 &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_9 &= \begin{pmatrix} 3.20250u^{49} - 44.5872u^{48} + \dots + 545.377u - 12.0177 \\ -22.0074u^{49} + 211.526u^{48} + \dots - 924.850u + 52.1829 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix} \\ a_4 &= \begin{pmatrix} 70.7134u^{49} - 796.258u^{48} + \dots + 24795.5u - 2262.67 \\ -65.6329u^{49} + 690.174u^{48} + \dots - 13362.0u + 1158.61 \end{pmatrix} \\ a_1 &= \begin{pmatrix} 6.69943u^{49} - 28.5149u^{48} + \dots - 5493.60u + 527.935 \\ 20.4159u^{49} - 227.260u^{48} + \dots + 5735.08u - 500.233 \end{pmatrix} \\ a_6 &= \begin{pmatrix} -71.2237u^{49} + 730.489u^{48} + \dots - 12363.3u + 1079.94 \\ 3.20274u^{49} - 17.3842u^{48} + \dots - 795.115u + 58.9383 \end{pmatrix} \\ a_{10} &= \begin{pmatrix} u^3 \\ u^5 - u^3 + u \end{pmatrix} \\ a_3 &= \begin{pmatrix} 7.76879u^{49} - 154.326u^{48} + \dots + 14845.3u - 1429.50 \\ -84.9603u^{49} + 868.882u^{48} + \dots - 13592.6u + 1152.16 \end{pmatrix} \\ a_2 &= \begin{pmatrix} 36.0253u^{49} - 376.781u^{48} + \dots + 8948.18u - 832.007 \\ -29.2879u^{49} + 282.012u^{48} + \dots - 2240.06u + 176.845 \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = -1

$$\text{(iii) Cusp Shapes} = \frac{9871134219706577536}{446874893374714199} u^{49} - \frac{123797457722876187653}{446874893374714199} u^{48} + \dots + \frac{6910675158351158980446}{446874893374714199} u - \frac{601472827700807026450}{446874893374714199}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_{10}$	$u^{50} + 19u^{49} + \dots + 5099u + 169$
$c_2, c_5, c_7$ $c_{11}$	$u^{50} + 11u^{49} + \dots + 165u + 13$
$c_3, c_8$	$(u^{25} + 4u^{24} + \dots + 3u + 1)^2$
$c_4, c_6, c_9$ $c_{12}$	$u^{50} + 5u^{49} + \dots + 4u + 1$



(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_{10}$	$y^{50} + 13y^{49} + \dots - 3709039y + 28561$
$c_2, c_5, c_7$ $c_{11}$	$y^{50} - 19y^{49} + \dots - 5099y + 169$
$c_3, c_8$	$(y^{25} - 8y^{24} + \dots + y - 1)^2$
$c_4, c_6, c_9$ $c_{12}$	$y^{50} + 21y^{49} + \dots + 52y + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_2^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.553707 + 0.832806I$ $a = 0.043757 - 0.161962I$ $b = 1.07602 + 1.01172I$	$-2.22763 + 6.76528I$	$0. - 5.55124I$
$u = 0.553707 - 0.832806I$ $a = 0.043757 + 0.161962I$ $b = 1.07602 - 1.01172I$	$-2.22763 - 6.76528I$	$0. + 5.55124I$
$u = -0.837139 + 0.557385I$ $a = 0.79159 - 1.91622I$ $b = 0.14846 - 1.48629I$	$-1.07852$	$0$
$u = -0.837139 - 0.557385I$ $a = 0.79159 + 1.91622I$ $b = 0.14846 + 1.48629I$	$-1.07852$	$0$
$u = 0.558430 + 0.809571I$ $a = -0.1049370 - 0.0051426I$ $b = 0.475534 - 0.512990I$	$-2.54415 - 3.36836I$	$0. + 11.53583I$
$u = 0.558430 - 0.809571I$ $a = -0.1049370 + 0.0051426I$ $b = 0.475534 + 0.512990I$	$-2.54415 + 3.36836I$	$0. - 11.53583I$
$u = -0.917168 + 0.480093I$ $a = -0.86443 + 1.71315I$ $b = -0.316452 + 1.330430I$	$-2.28095 + 0.36934I$	$0$
$u = -0.917168 - 0.480093I$ $a = -0.86443 - 1.71315I$ $b = -0.316452 - 1.330430I$	$-2.28095 - 0.36934I$	$0$
$u = 0.933200 + 0.451504I$ $a = -0.748966 - 0.796862I$ $b = 0.496615 - 0.374938I$	$-1.46923 - 1.71043I$	$0$
$u = 0.933200 - 0.451504I$ $a = -0.748966 + 0.796862I$ $b = 0.496615 + 0.374938I$	$-1.46923 + 1.71043I$	$0$

Solutions to $I_2^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.855986 + 0.425974I$		
$a = 0.52160 - 1.79372I$	$-1.96570 + 3.34746I$	$0. - 8.41971I$
$b = 0.054829 - 1.272590I$		
$u = -0.855986 - 0.425974I$		
$a = 0.52160 + 1.79372I$	$-1.96570 - 3.34746I$	$0. + 8.41971I$
$b = 0.054829 + 1.272590I$		
$u = -0.947983 + 0.107928I$		
$a = -0.39264 - 2.05984I$	$-2.54415 + 3.36836I$	$-3.18716 - 11.53583I$
$b = -0.334947 - 1.326910I$		
$u = -0.947983 - 0.107928I$		
$a = -0.39264 + 2.05984I$	$-2.54415 - 3.36836I$	$-3.18716 + 11.53583I$
$b = -0.334947 + 1.326910I$		
$u = -0.932177 + 0.127667I$		
$a = 1.17876 - 2.00119I$	$-4.36972 + 2.52616I$	$-17.6949 - 8.8137I$
$b = 0.66990 - 1.34941I$		
$u = -0.932177 - 0.127667I$		
$a = 1.17876 + 2.00119I$	$-4.36972 - 2.52616I$	$-17.6949 + 8.8137I$
$b = 0.66990 + 1.34941I$		
$u = 0.705776 + 0.790233I$		
$a = -0.379152 + 0.570875I$	$3.42082 + 2.87486I$	$0$
$b = -1.29001 - 1.06513I$		
$u = 0.705776 - 0.790233I$		
$a = -0.379152 - 0.570875I$	$3.42082 - 2.87486I$	$0$
$b = -1.29001 + 1.06513I$		
$u = 0.848218 + 0.375573I$		
$a = 1.68167 + 0.17265I$	$-2.28095 + 0.36934I$	$-1.79998 + 1.57281I$
$b = -0.434516 + 0.694240I$		
$u = 0.848218 - 0.375573I$		
$a = 1.68167 - 0.17265I$	$-2.28095 - 0.36934I$	$-1.79998 - 1.57281I$
$b = -0.434516 - 0.694240I$		

Solutions to $I_2^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.903202 + 0.595706I$ $a = -0.66474 - 1.60449I$ $b = 1.44211 - 0.54229I$	$-1.79493 - 2.26748I$	0
$u = 0.903202 - 0.595706I$ $a = -0.66474 + 1.60449I$ $b = 1.44211 + 0.54229I$	$-1.79493 + 2.26748I$	0
$u = 0.553438 + 0.956996I$ $a = 0.240338 + 0.074404I$ $b = 1.09277 + 1.06584I$	$3.6258 + 14.5513I$	0
$u = 0.553438 - 0.956996I$ $a = 0.240338 - 0.074404I$ $b = 1.09277 - 1.06584I$	$3.6258 - 14.5513I$	0
$u = -0.805706 + 0.801297I$ $a = -0.487390 + 0.656086I$ $b = -0.193006 + 0.633975I$	4.71625	0
$u = -0.805706 - 0.801297I$ $a = -0.487390 - 0.656086I$ $b = -0.193006 - 0.633975I$	4.71625	0
$u = 0.251795 + 1.124930I$ $a = -0.254137 + 0.049386I$ $b = -0.423621 + 0.366735I$	$3.42082 - 2.87486I$	0
$u = 0.251795 - 1.124930I$ $a = -0.254137 - 0.049386I$ $b = -0.423621 - 0.366735I$	$3.42082 + 2.87486I$	0
$u = 0.731346 + 0.379553I$ $a = -1.245790 + 0.394727I$ $b = 0.369373 - 0.633955I$	$-1.96570 - 3.34746I$	$0. + 8.41971I$
$u = 0.731346 - 0.379553I$ $a = -1.245790 - 0.394727I$ $b = 0.369373 + 0.633955I$	$-1.96570 + 3.34746I$	$0. - 8.41971I$

Solutions to $I_2^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.370215 + 1.168900I$		
$a = 0.223126 - 0.084389I$	$2.53033 - 8.55334I$	0
$b = 0.462705 - 0.346414I$		
$u = 0.370215 - 1.168900I$		
$a = 0.223126 + 0.084389I$	$2.53033 + 8.55334I$	0
$b = 0.462705 + 0.346414I$		
$u = 0.997780 + 0.713374I$		
$a = 0.85559 + 1.68166I$	$2.53033 - 8.55334I$	0
$b = -1.31365 + 1.39313I$		
$u = 0.997780 - 0.713374I$		
$a = 0.85559 - 1.68166I$	$2.53033 + 8.55334I$	0
$b = -1.31365 - 1.39313I$		
$u = -1.249550 + 0.152177I$		
$a = -0.58334 - 1.39167I$	$-2.22763 + 6.76528I$	0
$b = -0.482964 - 1.010150I$		
$u = -1.249550 - 0.152177I$		
$a = -0.58334 + 1.39167I$	$-2.22763 - 6.76528I$	0
$b = -0.482964 + 1.010150I$		
$u = 0.948687 + 0.830787I$		
$a = -0.602613 - 0.618497I$	1.63367	0
$b = 0.388518 - 1.079670I$		
$u = 0.948687 - 0.830787I$		
$a = -0.602613 + 0.618497I$	1.63367	0
$b = 0.388518 + 1.079670I$		
$u = 1.072930 + 0.680011I$		
$a = -0.71338 - 1.81412I$	$-3.78504 - 12.42080I$	0
$b = 1.16504 - 1.24505I$		
$u = 1.072930 - 0.680011I$		
$a = -0.71338 + 1.81412I$	$-3.78504 + 12.42080I$	0
$b = 1.16504 + 1.24505I$		

Solutions to $I_2^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.296060 + 0.117362I$ $a = 0.67390 + 1.34621I$ $b = 0.541462 + 0.992346I$	$-3.78504 + 12.42080I$	0
$u = -1.296060 - 0.117362I$ $a = 0.67390 - 1.34621I$ $b = 0.541462 - 0.992346I$	$-3.78504 - 12.42080I$	0
$u = 1.102950 + 0.724130I$ $a = 0.56636 + 1.78230I$ $b = -1.13689 + 1.28166I$	$3.6258 - 14.5513I$	0
$u = 1.102950 - 0.724130I$ $a = 0.56636 - 1.78230I$ $b = -1.13689 - 1.28166I$	$3.6258 + 14.5513I$	0
$u = 1.304940 + 0.232629I$ $a = 0.177583 + 0.626392I$ $b = -0.065583 + 0.228032I$	$-1.79493 + 2.26748I$	0
$u = 1.304940 - 0.232629I$ $a = 0.177583 - 0.626392I$ $b = -0.065583 - 0.228032I$	$-1.79493 - 2.26748I$	0
$u = 1.166690 + 0.634912I$ $a = 0.454441 + 0.458068I$ $b = -0.102112 + 0.474616I$	$-4.36972 - 2.52616I$	0
$u = 1.166690 - 0.634912I$ $a = 0.454441 - 0.458068I$ $b = -0.102112 - 0.474616I$	$-4.36972 + 2.52616I$	0
$u = 0.338458 + 0.034083I$ $a = -0.59797 + 2.01326I$ $b = 0.210423 - 0.573227I$	$-1.46923 + 1.71043I$	$-1.72966 - 1.33913I$
$u = 0.338458 - 0.034083I$ $a = -0.59797 - 2.01326I$ $b = 0.210423 + 0.573227I$	$-1.46923 - 1.71043I$	$-1.72966 + 1.33913I$

$$\text{III. } I_3^u = \langle -u^5a - u^5 + 2u^3a + u^3 - 2au + b - u + 1, 2u^4a - 3u^4 + \dots + a^2 + 2, u^6 + u^5 - u^4 - 2u^3 + u + 1 \rangle$$

(i) Arc colorings

$$a_7 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_5 = \begin{pmatrix} a \\ u^5a + u^5 - 2u^3a - u^3 + 2au + u - 1 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} u^5a - u^5 - 2u^3a + u^3 + 2au + a - u - 1 \\ u^5a + u^5 - 2u^3a - u^3 + 2au + u - 1 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -u^5a - u^5 + 2u^3a + u^3 - 2au + a - u + 1 \\ u^5a + u^5 - 2u^3a - u^3 + 2au + u - 1 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -u^5a + u^3a + 2u^4 + u^3 - au - 2u^2 + a - 2u \\ -1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} u^5a + u^5 - 2u^3a + 2u^4 - u^2a + u^3 + 2au - 2u^2 + a - u - 1 \\ -u^5a - u^5 + 2u^3a + u^2a + 3u^3 - 2au - 3u - 1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} u^3 \\ u^5 - u^3 + u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -u^5a + u^3a + u^3 - au + a + 1 \\ -u^5a - u^4a + u^5 + u^3a + 2u^2a + u^2 - a - 1 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -2u^5a - u^4a + u^5 + 2u^3a + 2u^4 + 2u^2a + u^3 - au - u^2 - 2u \\ 2u^5 + 2u^4 - 2u^3 + au - 2u^2 \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes =  $8u^4 - 8u^2 - 8u - 2$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_{10}$	$(u^6 + 3u^5 + 5u^4 + 4u^3 + 2u^2 + u + 1)^2$
$c_2, c_5, c_7$ $c_{11}$	$(u^6 - u^5 - u^4 + 2u^3 - u + 1)^2$
$c_3, c_8$	$u^{12} - 9u^{11} + \dots - 104u + 17$
$c_4, c_6, c_9$ $c_{12}$	$u^{12} + u^{11} + 2u^{10} - 2u^9 + 3u^8 - 3u^7 + 17u^6 - 9u^5 + 19u^4 - 5u^3 + 6u^2 + 1$



(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_{10}$	$(y^6 + y^5 + 5y^4 + 6y^2 + 3y + 1)^2$
$c_2, c_5, c_7$ $c_{11}$	$(y^6 - 3y^5 + 5y^4 - 4y^3 + 2y^2 - y + 1)^2$
$c_3, c_8$	$y^{12} + 3y^{11} + \dots + 64y + 289$
$c_4, c_6, c_9$ $c_{12}$	$y^{12} + 3y^{11} + \dots + 12y + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_3^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.002190 + 0.295542I$		
$a = 1.72095 + 1.00813I$	$-5.42615 - 1.84861I$	$-13.43343 + 1.58845I$
$b = 0.51345 + 1.52069I$		
$u = 1.002190 + 0.295542I$		
$a = 0.39343 - 2.26995I$	$-5.42615 - 1.84861I$	$-13.43343 + 1.58845I$
$b = -0.085204 - 0.856161I$		
$u = 1.002190 - 0.295542I$		
$a = 1.72095 - 1.00813I$	$-5.42615 + 1.84861I$	$-13.43343 - 1.58845I$
$b = 0.51345 - 1.52069I$		
$u = 1.002190 - 0.295542I$		
$a = 0.39343 + 2.26995I$	$-5.42615 + 1.84861I$	$-13.43343 - 1.58845I$
$b = -0.085204 + 0.856161I$		
$u = -0.428243 + 0.664531I$		
$a = -1.071180 - 0.131182I$	$2.13628 - 1.84861I$	$1.43343 + 1.58845I$
$b = 0.170133 - 0.403810I$		
$u = -0.428243 + 0.664531I$		
$a = -0.275983 - 0.338083I$	$2.13628 - 1.84861I$	$1.43343 + 1.58845I$
$b = -1.172330 + 0.699352I$		
$u = -0.428243 - 0.664531I$		
$a = -1.071180 + 0.131182I$	$2.13628 + 1.84861I$	$1.43343 - 1.58845I$
$b = 0.170133 + 0.403810I$		
$u = -0.428243 - 0.664531I$		
$a = -0.275983 + 0.338083I$	$2.13628 + 1.84861I$	$1.43343 - 1.58845I$
$b = -1.172330 - 0.699352I$		
$u = -1.073950 + 0.558752I$		
$a = 1.34873 - 1.14126I$	$-1.64493 + 11.38600I$	$-6.00000 - 11.02114I$
$b = -0.344080 - 0.571978I$		
$u = -1.073950 + 0.558752I$		
$a = -0.11594 + 2.13765I$	$-1.64493 + 11.38600I$	$-6.00000 - 11.02114I$
$b = 1.41803 + 1.13073I$		

Solutions to $I_3^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.073950 - 0.558752I$		
$a = 1.34873 + 1.14126I$	$-1.64493 - 11.38600I$	$-6.00000 + 11.02114I$
$b = -0.344080 + 0.571978I$		
$u = -1.073950 - 0.558752I$		
$a = -0.11594 - 2.13765I$	$-1.64493 - 11.38600I$	$-6.00000 + 11.02114I$
$b = 1.41803 - 1.13073I$		

$$\text{IV. } I_4^u = \langle u^5 - u^3 + b + u, u^3 + a, u^6 + u^5 - u^4 - 2u^3 + u + 1 \rangle$$

(i) Arc colorings

$$a_7 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -u^3 \\ -u^5 + u^3 - u \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -u^5 + 2u^3 - u \\ u^5 - u^3 + u \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u^5 - 2u^3 + u \\ -u^5 + u^3 - u \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} u \\ u^3 - u \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} u^3 \\ u^5 - u^3 + u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -1 \\ -u^2 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -u^2 - 1 \\ -u^4 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes =  $8u^4 - 8u^2 - 8u + 4$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_3, c_8$ $c_{10}$	$u^6 - 3u^5 + 5u^4 - 4u^3 + 2u^2 - u + 1$
$c_2, c_7$	$u^6 + u^5 - u^4 - 2u^3 + u + 1$
$c_4, c_5, c_6$ $c_9, c_{11}, c_{12}$	$u^6 - u^5 - u^4 + 2u^3 - u + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_3, c_8$ $c_{10}$	$y^6 + y^5 + 5y^4 + 6y^2 + 3y + 1$
$c_2, c_4, c_5$ $c_6, c_7, c_9$ $c_{11}, c_{12}$	$y^6 - 3y^5 + 5y^4 - 4y^3 + 2y^2 - y + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_4^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.002190 + 0.295542I$ $a = -0.743983 - 0.864704I$ $b = -0.428243 - 0.664531I$	$-3.78121 - 1.84861I$	$-7.43343 + 1.58845I$
$u = 1.002190 - 0.295542I$ $a = -0.743983 + 0.864704I$ $b = -0.428243 + 0.664531I$	$-3.78121 + 1.84861I$	$-7.43343 - 1.58845I$
$u = -0.428243 + 0.664531I$ $a = -0.488802 - 0.072152I$ $b = 1.002190 - 0.295542I$	$3.78121 - 1.84861I$	$7.43343 + 1.58845I$
$u = -0.428243 - 0.664531I$ $a = -0.488802 + 0.072152I$ $b = 1.002190 + 0.295542I$	$3.78121 + 1.84861I$	$7.43343 - 1.58845I$
$u = -1.073950 + 0.558752I$ $a = 0.23279 - 1.75889I$ $b = -1.073950 - 0.558752I$	$11.3860I$	$0. - 11.02114I$
$u = -1.073950 - 0.558752I$ $a = 0.23279 + 1.75889I$ $b = -1.073950 + 0.558752I$	$- 11.3860I$	$0. + 11.02114I$

$$\mathbf{V. } I_5^u = \langle -2u^{43}a - 4u^{42}a + \dots + 4b - 65, 50u^{43}a - 95u^{43} + \dots - 468a + 943, u^{44} + 3u^{43} + \dots - 14u - 1 \rangle$$

(i) Arc colorings

$$a_7 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_5 = \begin{pmatrix} a \\ \frac{1}{2}u^{43}a + u^{42}a + \dots + 163u + \frac{65}{4} \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 7u^{43}a + \frac{7}{8}u^{43} + \dots - \frac{5}{2}a - \frac{91}{8} \\ \frac{9}{4}u^{43}a + \frac{9}{4}u^{42}a + \dots + \frac{7}{2}a - 1 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -\frac{1}{2}u^{43}a - u^{42}a + \dots + a - \frac{65}{4} \\ \frac{1}{2}u^{43}a + u^{42}a + \dots + 163u + \frac{65}{4} \end{pmatrix}$$

$$a_1 = \begin{pmatrix} \frac{21}{4}u^{42}a + 2u^{43} + \dots - \frac{65}{4}a + \frac{23}{2} \\ -1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} \frac{49}{2}u^{43}a + 2u^{43} + \dots - \frac{139}{4}a + \frac{21}{2} \\ -\frac{5}{2}u^{43}a - \frac{1}{8}u^{43} + \dots + \frac{1}{4}a - \frac{7}{8} \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} u^3 \\ u^5 - u^3 + u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -\frac{9}{2}u^{43} - \frac{19}{4}u^{42} + \dots + a - \frac{41}{4} \\ -\frac{1}{2}u^{43}a - 6u^{43} + \dots + 226u + \frac{83}{4} \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -\frac{95}{4}u^{43}a + \frac{13}{8}u^{43} + \dots + 29a + \frac{79}{8} \\ \frac{7}{4}u^{43}a + \frac{11}{8}u^{43} + \dots - \frac{5}{2}a - \frac{11}{8} \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes =  $-\frac{97}{4}u^{43} - 54u^{42} + \dots + \frac{1089}{4}u + \frac{51}{4}$



(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_{10}$	$(u^{44} + 19u^{43} + \dots + 70u + 1)^2$
$c_2, c_5, c_7$ $c_{11}$	$(u^{44} - 3u^{43} + \dots + 14u - 1)^2$
$c_3, c_8$	$(u^{44} + u^{43} + \dots + 840u + 271)^2$
$c_4, c_6, c_9$ $c_{12}$	$u^{88} + 11u^{87} + \dots + 20u + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_{10}$	$(y^{44} + 17y^{43} + \dots - 3330y + 1)^2$
$c_2, c_5, c_7$ $c_{11}$	$(y^{44} - 19y^{43} + \dots - 70y + 1)^2$
$c_3, c_8$	$(y^{44} - 7y^{43} + \dots - 1090962y + 73441)^2$
$c_4, c_6, c_9$ $c_{12}$	$y^{88} - 39y^{87} + \dots - 250y + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_5^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.767481 + 0.649572I$ $a = -0.209749 + 0.455086I$ $b = 1.050190 + 0.737585I$	$3.61742 + 6.06223I$	$4.09868 - 3.56729I$
$u = 0.767481 + 0.649572I$ $a = -0.69676 - 2.08757I$ $b = 1.35788 - 1.16717I$	$3.61742 + 6.06223I$	$4.09868 - 3.56729I$
$u = 0.767481 - 0.649572I$ $a = -0.209749 - 0.455086I$ $b = 1.050190 - 0.737585I$	$3.61742 - 6.06223I$	$4.09868 + 3.56729I$
$u = 0.767481 - 0.649572I$ $a = -0.69676 + 2.08757I$ $b = 1.35788 + 1.16717I$	$3.61742 - 6.06223I$	$4.09868 + 3.56729I$
$u = 0.803659 + 0.647731I$ $a = 0.091655 - 0.132248I$ $b = -1.193190 - 0.680438I$	$5.35656 + 0.02685I$	$6.61609 + 1.70053I$
$u = 0.803659 + 0.647731I$ $a = 0.66010 + 2.10096I$ $b = -1.33111 + 1.04031I$	$5.35656 + 0.02685I$	$6.61609 + 1.70053I$
$u = 0.803659 - 0.647731I$ $a = 0.091655 + 0.132248I$ $b = -1.193190 + 0.680438I$	$5.35656 - 0.02685I$	$6.61609 - 1.70053I$
$u = 0.803659 - 0.647731I$ $a = 0.66010 - 2.10096I$ $b = -1.33111 - 1.04031I$	$5.35656 - 0.02685I$	$6.61609 - 1.70053I$
$u = 0.851089 + 0.590714I$ $a = 0.07497 - 1.50979I$ $b = 1.84927 - 0.00718I$	$-1.70735 - 2.34151I$	$6.14112 + 4.86696I$
$u = 0.851089 + 0.590714I$ $a = -0.93452 - 1.56224I$ $b = 1.150410 - 0.258778I$	$-1.70735 - 2.34151I$	$6.14112 + 4.86696I$

Solutions to $I_5^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.851089 - 0.590714I$ $a = 0.07497 + 1.50979I$ $b = 1.84927 + 0.00718I$	$-1.70735 + 2.34151I$	$6.14112 - 4.86696I$
$u = 0.851089 - 0.590714I$ $a = -0.93452 + 1.56224I$ $b = 1.150410 + 0.258778I$	$-1.70735 + 2.34151I$	$6.14112 - 4.86696I$
$u = -0.473872 + 0.833820I$ $a = -0.325630 - 0.110420I$ $b = 0.598560 - 0.485867I$	$2.48681 - 1.78904I$	$-1.77660 + 1.34347I$
$u = -0.473872 + 0.833820I$ $a = -0.181027 - 0.089265I$ $b = -1.054730 + 0.566967I$	$2.48681 - 1.78904I$	$-1.77660 + 1.34347I$
$u = -0.473872 - 0.833820I$ $a = -0.325630 + 0.110420I$ $b = 0.598560 + 0.485867I$	$2.48681 + 1.78904I$	$-1.77660 - 1.34347I$
$u = -0.473872 - 0.833820I$ $a = -0.181027 + 0.089265I$ $b = -1.054730 - 0.566967I$	$2.48681 + 1.78904I$	$-1.77660 - 1.34347I$
$u = -0.503143 + 0.946145I$ $a = 0.120712 - 0.412859I$ $b = 0.808803 - 0.791262I$	$5.08140 - 5.03772I$	$5.56121 + 5.67665I$
$u = -0.503143 + 0.946145I$ $a = -0.128920 - 0.292183I$ $b = -1.123420 + 0.229540I$	$5.08140 - 5.03772I$	$5.56121 + 5.67665I$
$u = -0.503143 - 0.946145I$ $a = 0.120712 + 0.412859I$ $b = 0.808803 + 0.791262I$	$5.08140 + 5.03772I$	$5.56121 - 5.67665I$
$u = -0.503143 - 0.946145I$ $a = -0.128920 + 0.292183I$ $b = -1.123420 - 0.229540I$	$5.08140 + 5.03772I$	$5.56121 - 5.67665I$

Solutions to $I_5^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.542619 + 0.934381I$ $a = -0.217567 + 0.342060I$ $b = -0.878871 + 0.768623I$	$5.35656 - 0.02685I$	$6.61609 - 1.70053I$
$u = -0.542619 + 0.934381I$ $a = 0.055966 + 0.360352I$ $b = 1.039850 - 0.142869I$	$5.35656 - 0.02685I$	$6.61609 - 1.70053I$
$u = -0.542619 - 0.934381I$ $a = -0.217567 - 0.342060I$ $b = -0.878871 - 0.768623I$	$5.35656 + 0.02685I$	$6.61609 + 1.70053I$
$u = -0.542619 - 0.934381I$ $a = 0.055966 - 0.360352I$ $b = 1.039850 + 0.142869I$	$5.35656 + 0.02685I$	$6.61609 + 1.70053I$
$u = 0.892273 + 0.637275I$ $a = -0.798485 + 0.407722I$ $b = -1.70312 - 0.82985I$	$5.08140 - 5.03772I$	$5.56121 + 5.67665I$
$u = 0.892273 + 0.637275I$ $a = 0.79044 + 2.21552I$ $b = -0.931912 + 0.821648I$	$5.08140 - 5.03772I$	$5.56121 + 5.67665I$
$u = 0.892273 - 0.637275I$ $a = -0.798485 - 0.407722I$ $b = -1.70312 + 0.82985I$	$5.08140 + 5.03772I$	$5.56121 - 5.67665I$
$u = 0.892273 - 0.637275I$ $a = 0.79044 - 2.21552I$ $b = -0.931912 - 0.821648I$	$5.08140 + 5.03772I$	$5.56121 - 5.67665I$
$u = -0.802626 + 0.757447I$ $a = -0.492685 + 1.009980I$ $b = 0.379182 + 0.670238I$	4.62114	$5.54727 + 0.I$
$u = -0.802626 + 0.757447I$ $a = -0.458276 + 0.324516I$ $b = -0.733776 + 0.580087I$	4.62114	$5.54727 + 0.I$

Solutions to $I_5^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.802626 - 0.757447I$ $a = -0.492685 - 1.009980I$ $b = 0.379182 - 0.670238I$	4.62114	$5.54727 + 0.I$
$u = -0.802626 - 0.757447I$ $a = -0.458276 - 0.324516I$ $b = -0.733776 - 0.580087I$	4.62114	$5.54727 + 0.I$
$u = 0.918693 + 0.631895I$ $a = 1.086560 - 0.368999I$ $b = 1.77703 + 0.97307I$	$3.15038 - 11.06350I$	$2.31684 + 10.51224I$
$u = 0.918693 + 0.631895I$ $a = -0.79220 - 2.32690I$ $b = 0.801722 - 0.841787I$	$3.15038 - 11.06350I$	$2.31684 + 10.51224I$
$u = 0.918693 - 0.631895I$ $a = 1.086560 + 0.368999I$ $b = 1.77703 - 0.97307I$	$3.15038 + 11.06350I$	$2.31684 - 10.51224I$
$u = 0.918693 - 0.631895I$ $a = -0.79220 + 2.32690I$ $b = 0.801722 + 0.841787I$	$3.15038 + 11.06350I$	$2.31684 - 10.51224I$
$u = 1.11703$ $a = -0.559527 + 1.206560I$ $b = -0.064688 + 0.847977I$	-3.19687	-5.88290
$u = 1.11703$ $a = -0.559527 - 1.206560I$ $b = -0.064688 - 0.847977I$	-3.19687	-5.88290
$u = -0.972088 + 0.562819I$ $a = 1.63782 - 0.13643I$ $b = -0.300509 - 0.254450I$	$-3.73793 + 3.81466I$	$-6.30765 - 8.57961I$
$u = -0.972088 + 0.562819I$ $a = -0.71054 + 2.33938I$ $b = 1.35813 + 1.69326I$	$-3.73793 + 3.81466I$	$-6.30765 - 8.57961I$

Solutions to $I_5^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.972088 - 0.562819I$ $a = 1.63782 + 0.13643I$ $b = -0.300509 + 0.254450I$	$-3.73793 - 3.81466I$	$-6.30765 + 8.57961I$
$u = -0.972088 - 0.562819I$ $a = -0.71054 - 2.33938I$ $b = 1.35813 - 1.69326I$	$-3.73793 - 3.81466I$	$-6.30765 + 8.57961I$
$u = -0.714060 + 0.501949I$ $a = 1.34919 + 0.68585I$ $b = 1.59813 - 1.14955I$	$-2.85723 + 0.57531I$	$-2.18655 + 5.07930I$
$u = -0.714060 + 0.501949I$ $a = 1.59862 - 1.48482I$ $b = 0.0868291 - 0.0247619I$	$-2.85723 + 0.57531I$	$-2.18655 + 5.07930I$
$u = -0.714060 - 0.501949I$ $a = 1.34919 - 0.68585I$ $b = 1.59813 + 1.14955I$	$-2.85723 - 0.57531I$	$-2.18655 - 5.07930I$
$u = -0.714060 - 0.501949I$ $a = 1.59862 + 1.48482I$ $b = 0.0868291 + 0.0247619I$	$-2.85723 - 0.57531I$	$-2.18655 - 5.07930I$
$u = 1.182370 + 0.249072I$ $a = 1.33410 + 0.52761I$ $b = 0.716253 + 0.936843I$	$-3.73793 + 3.81466I$	$-6.30765 - 8.57961I$
$u = 1.182370 + 0.249072I$ $a = -0.04691 - 1.87311I$ $b = -0.139509 - 0.906647I$	$-3.73793 + 3.81466I$	$-6.30765 - 8.57961I$
$u = 1.182370 - 0.249072I$ $a = 1.33410 - 0.52761I$ $b = 0.716253 - 0.936843I$	$-3.73793 - 3.81466I$	$-6.30765 + 8.57961I$
$u = 1.182370 - 0.249072I$ $a = -0.04691 + 1.87311I$ $b = -0.139509 + 0.906647I$	$-3.73793 - 3.81466I$	$-6.30765 + 8.57961I$

Solutions to $I_5^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.044760 + 0.613681I$		
$a = -0.947346 + 0.792517I$	$0.48539 + 6.82448I$	$0. - 6.22925I$
$b = 0.530131 + 0.447375I$		
$u = -1.044760 + 0.613681I$		
$a = 0.34576 - 1.93178I$	$0.48539 + 6.82448I$	$0. - 6.22925I$
$b = -1.17630 - 1.20444I$		
$u = -1.044760 - 0.613681I$		
$a = -0.947346 - 0.792517I$	$0.48539 - 6.82448I$	$0. + 6.22925I$
$b = 0.530131 - 0.447375I$		
$u = -1.044760 - 0.613681I$		
$a = 0.34576 + 1.93178I$	$0.48539 - 6.82448I$	$0. + 6.22925I$
$b = -1.17630 + 1.20444I$		
$u = 1.215330 + 0.165372I$		
$a = -1.025870 - 0.466462I$	$-2.85723 - 0.57531I$	$0$
$b = -0.541188 - 0.729277I$		
$u = 1.215330 + 0.165372I$		
$a = 0.04140 + 1.59807I$	$-2.85723 - 0.57531I$	$0$
$b = 0.156527 + 0.829153I$		
$u = 1.215330 - 0.165372I$		
$a = -1.025870 + 0.466462I$	$-2.85723 + 0.57531I$	$0$
$b = -0.541188 + 0.729277I$		
$u = 1.215330 - 0.165372I$		
$a = 0.04140 - 1.59807I$	$-2.85723 + 0.57531I$	$0$
$b = 0.156527 - 0.829153I$		
$u = -0.695081 + 0.217967I$		
$a = 0.305437 + 1.080750I$	$0.57189 - 7.31268I$	$-5.07151 + 6.33709I$
$b = 1.56163 - 0.21982I$		
$u = -0.695081 + 0.217967I$		
$a = 3.07338 - 0.83141I$	$0.57189 - 7.31268I$	$-5.07151 + 6.33709I$
$b = 0.347865 - 0.007131I$		



Solutions to $I_5^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.695081 - 0.217967I$ $a = 0.305437 - 1.080750I$ $b = 1.56163 + 0.21982I$	$0.57189 + 7.31268I$	$-5.07151 - 6.33709I$
$u = -0.695081 - 0.217967I$ $a = 3.07338 + 0.83141I$ $b = 0.347865 + 0.007131I$	$0.57189 + 7.31268I$	$-5.07151 - 6.33709I$
$u = -1.109330 + 0.644832I$ $a = -0.65014 + 1.36464I$ $b = 0.683547 + 0.755743I$	$0.57189 + 7.31268I$	0
$u = -1.109330 + 0.644832I$ $a = 0.09902 - 1.63155I$ $b = -1.15242 - 0.87299I$	$0.57189 + 7.31268I$	0
$u = -1.109330 - 0.644832I$ $a = -0.65014 - 1.36464I$ $b = 0.683547 - 0.755743I$	$0.57189 - 7.31268I$	0
$u = -1.109330 - 0.644832I$ $a = 0.09902 + 1.63155I$ $b = -1.15242 + 0.87299I$	$0.57189 - 7.31268I$	0
$u = 1.318490 + 0.024433I$ $a = 0.150294 + 0.994451I$ $b = 0.127511 + 0.470493I$	$-1.70735 + 2.34151I$	0
$u = 1.318490 + 0.024433I$ $a = -0.274504 + 0.224235I$ $b = -0.188454 - 0.112226I$	$-1.70735 + 2.34151I$	0
$u = 1.318490 - 0.024433I$ $a = 0.150294 - 0.994451I$ $b = 0.127511 - 0.470493I$	$-1.70735 - 2.34151I$	0
$u = 1.318490 - 0.024433I$ $a = -0.274504 - 0.224235I$ $b = -0.188454 + 0.112226I$	$-1.70735 - 2.34151I$	0

Solutions to $I_5^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.110670 + 0.713391I$ $a = 0.113297 + 1.048100I$ $b = 1.135860 + 0.456326I$	$3.61742 + 6.06223I$	0
$u = -1.110670 + 0.713391I$ $a = 0.44380 - 1.68331I$ $b = -0.806935 - 1.052440I$	$3.61742 + 6.06223I$	0
$u = -1.110670 - 0.713391I$ $a = 0.113297 - 1.048100I$ $b = 1.135860 - 0.456326I$	$3.61742 - 6.06223I$	0
$u = -1.110670 - 0.713391I$ $a = 0.44380 + 1.68331I$ $b = -0.806935 + 1.052440I$	$3.61742 - 6.06223I$	0
$u = -0.235182 + 0.635023I$ $a = 1.24503 + 0.86650I$ $b = 0.016785 + 0.660631I$	$0.48539 - 6.82448I$	$-1.22382 + 6.22925I$
$u = -0.235182 + 0.635023I$ $a = -0.118272 + 0.377558I$ $b = 1.207550 - 0.716047I$	$0.48539 - 6.82448I$	$-1.22382 + 6.22925I$
$u = -0.235182 - 0.635023I$ $a = 1.24503 - 0.86650I$ $b = 0.016785 - 0.660631I$	$0.48539 + 6.82448I$	$-1.22382 - 6.22925I$
$u = -0.235182 - 0.635023I$ $a = -0.118272 - 0.377558I$ $b = 1.207550 + 0.716047I$	$0.48539 + 6.82448I$	$-1.22382 - 6.22925I$
$u = -1.133930 + 0.701691I$ $a = -0.187501 - 1.213320I$ $b = -1.214170 - 0.539939I$	$3.15038 + 11.06350I$	0
$u = -1.133930 + 0.701691I$ $a = -0.50414 + 1.69437I$ $b = 0.748726 + 1.033840I$	$3.15038 + 11.06350I$	0

Solutions to $I_5^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.133930 - 0.701691I$ $a = -0.187501 + 1.213320I$ $b = -1.214170 + 0.539939I$	$3.15038 - 11.06350I$	0
$u = -1.133930 - 0.701691I$ $a = -0.50414 - 1.69437I$ $b = 0.748726 - 1.033840I$	$3.15038 - 11.06350I$	0
$u = -0.604720 + 0.161152I$ $a = -0.485397 - 0.758391I$ $b = -1.382110 + 0.218862I$	$2.48681 - 1.78904I$	$-1.77660 + 1.34347I$
$u = -0.604720 + 0.161152I$ $a = -3.01193 + 0.44916I$ $b = -0.403113 - 0.055546I$	$2.48681 - 1.78904I$	$-1.77660 + 1.34347I$
$u = -0.604720 - 0.161152I$ $a = -0.485397 + 0.758391I$ $b = -1.382110 - 0.218862I$	$2.48681 + 1.78904I$	$-1.77660 - 1.34347I$
$u = -0.604720 - 0.161152I$ $a = -3.01193 - 0.44916I$ $b = -0.403113 + 0.055546I$	$2.48681 + 1.78904I$	$-1.77660 - 1.34347I$
$u = -0.131617$ $a = 5.64036 + 4.49230I$ $b = 0.731173 - 0.579698I$	-3.19687	-5.88290
$u = -0.131617$ $a = 5.64036 - 4.49230I$ $b = 0.731173 + 0.579698I$	-3.19687	-5.88290

$$\text{VI. } J_6^u = \langle 3u^{15} + 5u^{14} + \dots + b - 1, -7u^{15} - 28u^{14} + \dots + 2a - 22, u^{16} + 4u^{15} + \dots + 8u + 2 \rangle$$

(i) Arc colorings

$$a_7 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_5 = \begin{pmatrix} \frac{7}{2}u^{15} + 14u^{14} + \dots + \frac{59}{2}u + 11 \\ -3u^{15} - 5u^{14} + \dots + u + 1 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -\frac{11}{2}u^{15} - 17u^{14} + \dots - \frac{67}{2}u - 8 \\ u^{15} + 3u^{14} + \dots + 7u + 1 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} \frac{13}{2}u^{15} + 19u^{14} + \dots + \frac{57}{2}u + 10 \\ -3u^{15} - 5u^{14} + \dots + u + 1 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -\frac{13}{2}u^{15} - 20u^{14} + \dots - \frac{67}{2}u - 11 \\ 4u^{15} + 9u^{14} + \dots + 7u + 1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} \frac{17}{2}u^{15} + 30u^{14} + \dots + \frac{143}{2}u + 24 \\ -4u^{15} - 13u^{14} + \dots - 30u - 9 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} u^3 \\ u^5 - u^3 + u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} \frac{3}{2}u^{15} + 4u^{14} + \dots + \frac{5}{2}u + 2 \\ -4u^{15} - 11u^{14} + \dots - 18u - 5 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} \frac{13}{2}u^{15} + 26u^{14} + \dots + \frac{139}{2}u + 25 \\ -7u^{15} - 21u^{14} + \dots - 41u - 11 \end{pmatrix}$$

(ii) Obstruction class = 1

$$\text{(iii) Cusp Shapes} = -38u^{15} - 69u^{14} + 46u^{13} + 251u^{12} + 175u^{11} - 71u^{10} - 114u^9 - 26u^8 - 246u^7 - 430u^6 - 310u^5 - 60u^4 - 64u^3 - 35u^2 + 26u + 42$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_{10}$	$u^{16} - 6u^{15} + \dots + 4u + 4$
$c_2, c_7$	$u^{16} + 4u^{15} + \dots + 8u + 2$
$c_3, c_8$	$(u^8 + 2u^7 + 2u^6 + u^2 + u + 1)^2$
$c_4, c_6, c_9$ $c_{12}$	$u^{16} - 6u^{14} + \dots - u^2 + 1$
$c_5, c_{11}$	$u^{16} - 4u^{15} + \dots - 8u + 2$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_{10}$	$y^{16} + 2y^{15} + \dots + 152y + 16$
$c_2, c_5, c_7$ $c_{11}$	$y^{16} - 6y^{15} + \dots + 4y + 4$
$c_3, c_8$	$(y^8 + 4y^6 + 2y^5 + 2y^4 + 4y^3 + y^2 + y + 1)^2$
$c_4, c_6, c_9$ $c_{12}$	$y^{16} - 12y^{15} + \dots - 2y + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_6^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.484640 + 0.846797I$ $a = -0.680045 + 0.269791I$ $b = -0.618812 - 0.101203I$	$2.47558 - 7.94608I$	$0.92481 + 5.79314I$
$u = 0.484640 - 0.846797I$ $a = -0.680045 - 0.269791I$ $b = -0.618812 + 0.101203I$	$2.47558 + 7.94608I$	$0.92481 - 5.79314I$
$u = -0.620876 + 0.837136I$ $a = 0.129335 + 0.230266I$ $b = 0.980946 - 0.677334I$	$3.68697 - 2.14875I$	$5.80448 + 2.27366I$
$u = -0.620876 - 0.837136I$ $a = 0.129335 - 0.230266I$ $b = 0.980946 + 0.677334I$	$3.68697 + 2.14875I$	$5.80448 - 2.27366I$
$u = -0.868732 + 0.620313I$ $a = 0.43706 - 1.78935I$ $b = -1.81641 - 0.27594I$	$-2.07294 + 2.43245I$	$-21.6315 - 8.0683I$
$u = -0.868732 - 0.620313I$ $a = 0.43706 + 1.78935I$ $b = -1.81641 + 0.27594I$	$-2.07294 - 2.43245I$	$-21.6315 + 8.0683I$
$u = -0.597300 + 0.462977I$ $a = 1.26482 - 0.64500I$ $b = -1.184110 + 0.133480I$	$1.66766 - 7.05131I$	$3.40216 + 4.97138I$
$u = -0.597300 - 0.462977I$ $a = 1.26482 + 0.64500I$ $b = -1.184110 - 0.133480I$	$1.66766 + 7.05131I$	$3.40216 - 4.97138I$
$u = -1.034390 + 0.719349I$ $a = -0.62644 + 1.38546I$ $b = 0.99042 + 1.11066I$	$2.47558 + 7.94608I$	$0.92481 - 5.79314I$
$u = -1.034390 - 0.719349I$ $a = -0.62644 - 1.38546I$ $b = 0.99042 - 1.11066I$	$2.47558 - 7.94608I$	$0.92481 + 5.79314I$

Solutions to $I_6^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.125140 + 0.624650I$		
$a = -0.20300 + 1.50281I$	$1.66766 + 7.05131I$	$3.40216 - 4.97138I$
$b = 0.927126 + 0.676564I$		
$u = -1.125140 - 0.624650I$		
$a = -0.20300 - 1.50281I$	$1.66766 - 7.05131I$	$3.40216 + 4.97138I$
$b = 0.927126 - 0.676564I$		
$u = 0.270172 + 0.658282I$		
$a = 0.744963 - 0.703702I$	$3.68697 - 2.14875I$	$5.80448 + 2.27366I$
$b = 0.764226 + 0.048273I$		
$u = 0.270172 - 0.658282I$		
$a = 0.744963 + 0.703702I$	$3.68697 + 2.14875I$	$5.80448 - 2.27366I$
$b = 0.764226 - 0.048273I$		
$u = 1.49163 + 0.08776I$		
$a = -0.066697 - 0.588457I$	$-2.07294 + 2.43245I$	$-21.6315 - 8.0683I$
$b = -0.043384 - 0.465221I$		
$u = 1.49163 - 0.08776I$		
$a = -0.066697 + 0.588457I$	$-2.07294 - 2.43245I$	$-21.6315 + 8.0683I$
$b = -0.043384 + 0.465221I$		



$$\text{VII. } I_7^u = \langle -u^2a + b - 1, a^2 + 2au + 2u^2 + a + 3u + 2, u^3 + u^2 - 1 \rangle$$

(i) Arc colorings

$$a_7 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_5 = \begin{pmatrix} a \\ u^2a + 1 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -a + 1 \\ -u^2a + u^2 - 1 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -u \\ u^2 + u - 1 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -u^2a + a - 1 \\ u^2a + 1 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -a - 2u - 1 \\ -1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -u^2a - a - 2u - 1 \\ -u^2a - au - u^2 + a - 1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -u^2 + 1 \\ u^2 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -u^2a + u^2 + a - 2 \\ u^2a - u^2 + 1 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -au - u^2 - u - 2 \\ u^2a + au - a - u \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes =  $-8u - 2$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_{10}$	$(u^3 + u^2 + 2u + 1)^2$
$c_2, c_5, c_7$ $c_{11}$	$(u^3 - u^2 + 1)^2$
$c_3, c_8$	$(u + 1)^6$
$c_4, c_6, c_9$ $c_{12}$	$u^6 + u^5 + 2u^4 + 2u^3 + 2u^2 + 2u + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_{10}$	$(y^3 + 3y^2 + 2y - 1)^2$
$c_2, c_5, c_7$ $c_{11}$	$(y^3 - y^2 + 2y - 1)^2$
$c_3, c_8$	$(y - 1)^6$
$c_4, c_6, c_9$ $c_{12}$	$y^6 + 3y^5 + 4y^4 + 2y^3 + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_7^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.877439 + 0.744862I$		
$a = 0.562133 - 1.239140I$	$4.40332 + 5.65624I$	$5.01951 - 5.95889I$
$b = -0.498832 - 1.001300I$		
$u = -0.877439 + 0.744862I$		
$a = 0.192744 - 0.250580I$	$4.40332 + 5.65624I$	$5.01951 - 5.95889I$
$b = 0.713912 - 0.305839I$		
$u = -0.877439 - 0.744862I$		
$a = 0.562133 + 1.239140I$	$4.40332 - 5.65624I$	$5.01951 + 5.95889I$
$b = -0.498832 + 1.001300I$		
$u = -0.877439 - 0.744862I$		
$a = 0.192744 + 0.250580I$	$4.40332 - 5.65624I$	$5.01951 + 5.95889I$
$b = 0.713912 + 0.305839I$		
$u = 0.754878$		
$a = -1.25488 + 1.95694I$	$-3.87184$	$-8.03900$
$b = 0.284920 + 1.115140I$		
$u = 0.754878$		
$a = -1.25488 - 1.95694I$	$-3.87184$	$-8.03900$
$b = 0.284920 - 1.115140I$		

$$\text{VIII. } I_8^u = \langle -u^3a - u^2a - u^3 + au + 2b + 1, -u^2a + u^3 + a^2 + au - 2u^2 + a - u + 2, u^4 - u^2 + 1 \rangle$$

(i) Arc colorings

$$a_7 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_5 = \begin{pmatrix} a \\ \frac{1}{2}u^3a + \frac{1}{2}u^3 + \dots - \frac{1}{2}au - \frac{1}{2} \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -\frac{1}{2}u^3a + u^3 + \dots - \frac{1}{2}a + \frac{1}{2} \\ u^3 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -\frac{1}{2}u^3a - \frac{1}{2}u^3 + \dots + a + \frac{1}{2} \\ \frac{1}{2}u^3a + \frac{1}{2}u^3 + \dots - \frac{1}{2}au - \frac{1}{2} \end{pmatrix}$$

$$a_1 = \begin{pmatrix} \frac{1}{2}u^3a - \frac{1}{2}u^2 + \dots - \frac{1}{2}a + \frac{1}{2} \\ 1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -\frac{1}{2}u^3a + \frac{1}{2}u^2 + \dots + \frac{1}{2}a - \frac{1}{2} \\ -1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} u^3 \\ 0 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} a \\ \frac{1}{2}u^3a + \frac{1}{2}u^3 + \dots - \frac{1}{2}au - \frac{1}{2} \end{pmatrix}$$

$$a_2 = \begin{pmatrix} \frac{1}{2}u^3a - \frac{1}{2}u^2 + \dots + \frac{1}{2}a + \frac{1}{2} \\ \frac{1}{2}u^3a + \frac{1}{2}u^3 + \dots - \frac{1}{2}au + \frac{1}{2} \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes =  $4u^2 - 8$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1, c_2$	$(u - 1)^8$
$c_3, c_8$	$u^8 - 2u^6 + 2u^5 + 2u^4 - 2u^3 + 3u^2 + 4u + 1$
$c_4, c_{12}$	$u^8 - 4u^7 + 10u^6 - 16u^5 + 18u^4 - 14u^3 + 7u^2 - 2u + 1$
$c_5$	$(u + 1)^8$
$c_6, c_9$	$(u^2 + 1)^4$
$c_7, c_{11}$	$(u^4 - u^2 + 1)^2$
$c_{10}$	$(u^2 - u + 1)^4$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1, c_2, c_5$	$(y - 1)^8$
$c_3, c_8$	$y^8 - 4y^7 + 8y^6 - 6y^5 + 2y^4 - 12y^3 + 29y^2 - 10y + 1$
$c_4, c_{12}$	$y^8 + 4y^7 + 8y^6 + 6y^5 + 2y^4 + 12y^3 + 29y^2 + 10y + 1$
$c_6, c_9$	$(y + 1)^8$
$c_7, c_{11}$	$(y^2 - y + 1)^4$
$c_{10}$	$(y^2 + y + 1)^4$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_g^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.866025 + 0.500000I$		
$a = 0.131115 + 0.786143I$	$-3.28987 - 2.02988I$	$-6.00000 + 3.46410I$
$b = -1.060940 + 0.445679I$		
$u = 0.866025 + 0.500000I$		
$a = -1.49714 - 0.42012I$	$-3.28987 - 2.02988I$	$-6.00000 + 3.46410I$
$b = 0.060942 - 0.445679I$		
$u = 0.866025 - 0.500000I$		
$a = 0.131115 - 0.786143I$	$-3.28987 + 2.02988I$	$-6.00000 - 3.46410I$
$b = -1.060940 - 0.445679I$		
$u = 0.866025 - 0.500000I$		
$a = -1.49714 + 0.42012I$	$-3.28987 + 2.02988I$	$-6.00000 - 3.46410I$
$b = 0.060942 + 0.445679I$		
$u = -0.866025 + 0.500000I$		
$a = -0.553254 + 1.002550I$	$-3.28987 + 2.02988I$	$-6.00000 - 3.46410I$
$b = -0.69440 + 1.28601I$		
$u = -0.866025 + 0.500000I$		
$a = 0.91928 - 2.36858I$	$-3.28987 + 2.02988I$	$-6.00000 - 3.46410I$
$b = -0.305600 - 1.286010I$		
$u = -0.866025 - 0.500000I$		
$a = -0.553254 - 1.002550I$	$-3.28987 - 2.02988I$	$-6.00000 + 3.46410I$
$b = -0.69440 - 1.28601I$		
$u = -0.866025 - 0.500000I$		
$a = 0.91928 + 2.36858I$	$-3.28987 - 2.02988I$	$-6.00000 + 3.46410I$
$b = -0.305600 + 1.286010I$		



$$\text{IX. } I_9^u = \langle 7.93 \times 10^6 b - 7.18 \times 10^4 a^7 + \dots - 9.81 \times 10^6 a + 3.14 \times 10^6, a^8 + 6a^6 + \dots - 60a + 73, u - 1 \rangle$$

(i) Arc colorings

$$a_7 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} a \\ 0.00904756a^7 + 0.00216843a^6 + \dots + 1.23707a - 0.396120 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -0.00216843a^7 - 0.00652609a^6 + \dots - 0.146733a + 1.66047 \\ 0.00216843a^7 + 0.00652609a^6 + \dots + 0.146733a + 1.33953 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -0.00904756a^7 - 0.00216843a^6 + \dots - 0.237070a + 0.396120 \\ 0.00904756a^7 + 0.00216843a^6 + \dots + 1.23707a - 0.396120 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} 0.00216843a^7 + 0.00652609a^6 + \dots + 0.146733a - 0.660472 \\ -1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 0.000900759a^7 + 0.0124795a^6 + \dots + 0.0865369a + 3.18407 \\ 0.00867371a^7 + 0.0261044a^6 + \dots + 0.586933a + 1.35811 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -0.0271427a^7 - 0.00650529a^6 + \dots - 1.71121a + 1.18836 \\ -0.00904756a^7 - 0.00216843a^6 + \dots - 0.237070a + 0.396120 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -0.0242196a^7 - 0.0533541a^6 + \dots - 1.58772a - 0.706203 \\ -0.00994138a^7 - 0.0201510a^6 + \dots - 0.647129a - 0.513573 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes

$$= -\frac{864}{10477}a^7 - \frac{144}{10477}a^6 - \frac{5208}{10477}a^5 + \frac{4316}{10477}a^4 - \frac{14760}{10477}a^3 + \frac{13092}{10477}a^2 - \frac{52032}{10477}a - \frac{40648}{10477}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1$	$(u^2 - u + 1)^4$
$c_2, c_5$	$(u^4 - u^2 + 1)^2$
$c_3, c_8$	$u^8 - 2u^6 + 2u^5 + 2u^4 - 2u^3 + 3u^2 + 4u + 1$
$c_4, c_{12}$	$(u^2 + 1)^4$
$c_6, c_9$	$u^8 - 4u^7 + 10u^6 - 16u^5 + 18u^4 - 14u^3 + 7u^2 - 2u + 1$
$c_7, c_{10}$	$(u - 1)^8$
$c_{11}$	$(u + 1)^8$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1$	$(y^2 + y + 1)^4$
$c_2, c_5$	$(y^2 - y + 1)^4$
$c_3, c_8$	$y^8 - 4y^7 + 8y^6 - 6y^5 + 2y^4 - 12y^3 + 29y^2 - 10y + 1$
$c_4, c_{12}$	$(y + 1)^8$
$c_6, c_9$	$y^8 + 4y^7 + 8y^6 + 6y^5 + 2y^4 + 12y^3 + 29y^2 + 10y + 1$
$c_7, c_{10}, c_{11}$	$(y - 1)^8$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_9^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.00000$ $a = 0.445679 + 0.939058I$ $b = 1.000000I$	$-3.28987 + 2.02988I$	$-6.00000 - 3.46410I$
$u = 1.00000$ $a = 0.445679 - 0.939058I$ $b = -1.000000I$	$-3.28987 - 2.02988I$	$-6.00000 + 3.46410I$
$u = 1.00000$ $a = 1.28601 + 1.30560I$ $b = 1.000000I$	$-3.28987 - 2.02988I$	$-6.00000 + 3.46410I$
$u = 1.00000$ $a = 1.28601 - 1.30560I$ $b = -1.000000I$	$-3.28987 + 2.02988I$	$-6.00000 - 3.46410I$
$u = 1.00000$ $a = -0.44568 + 2.06094I$ $b = 1.000000I$	$-3.28987 + 2.02988I$	$-6.00000 - 3.46410I$
$u = 1.00000$ $a = -0.44568 - 2.06094I$ $b = -1.000000I$	$-3.28987 - 2.02988I$	$-6.00000 + 3.46410I$
$u = 1.00000$ $a = -1.28601 + 1.69440I$ $b = 1.000000I$	$-3.28987 - 2.02988I$	$-6.00000 + 3.46410I$
$u = 1.00000$ $a = -1.28601 - 1.69440I$ $b = -1.000000I$	$-3.28987 + 2.02988I$	$-6.00000 - 3.46410I$

## X. u-Polynomials

Crossings	u-Polynomials at each crossing
$c_1, c_{10}$	$(u-1)^8(u^2-u+1)^4(u^3+u^2+2u+1)^2$ $\cdot (u^6-3u^5+5u^4-4u^3+2u^2-u+1)$ $\cdot ((u^6+3u^5+5u^4+4u^3+2u^2+u+1)^2)(u^{16}-6u^{15}+\dots+4u+4)$ $\cdot (u^{18}+7u^{17}+\dots-4u+4)(u^{44}+19u^{43}+\dots+70u+1)^2$ $\cdot (u^{50}+19u^{49}+\dots+5099u+169)$
$c_2, c_7$	$(u-1)^8(u^3-u^2+1)^2(u^4-u^2+1)^2(u^6-u^5-u^4+2u^3-u+1)^2$ $\cdot (u^6+u^5-u^4-2u^3+u+1)(u^{16}+4u^{15}+\dots+8u+2)$ $\cdot (u^{18}+3u^{17}+\dots+6u+2)(u^{44}-3u^{43}+\dots+14u-1)^2$ $\cdot (u^{50}+11u^{49}+\dots+165u+13)$
$c_3, c_8$	$(u+1)^6(u^6-3u^5+5u^4-4u^3+2u^2-u+1)$ $\cdot (u^8-2u^6+2u^5+2u^4-2u^3+3u^2+4u+1)^2$ $\cdot ((u^8+2u^7+2u^6+u^2+u+1)^2)(u^{12}-9u^{11}+\dots-104u+17)$ $\cdot (u^{18}-9u^{17}+\dots-16u+8)(u^{25}+4u^{24}+\dots+3u+1)^2$ $\cdot (u^{44}+u^{43}+\dots+840u+271)^2$
$c_4, c_6, c_9$ $c_{12}$	$((u^2+1)^4)(u^6-u^5+\dots-u+1)(u^6+u^5+\dots+2u+1)$ $\cdot (u^8-4u^7+10u^6-16u^5+18u^4-14u^3+7u^2-2u+1)$ $\cdot (u^{12}+u^{11}+2u^{10}-2u^9+3u^8-3u^7+17u^6-9u^5+19u^4-5u^3+6u^2+1)$ $\cdot (u^{16}-6u^{14}+\dots-u^2+1)(u^{18}+5u^{16}+\dots+u+1)$ $\cdot (u^{50}+5u^{49}+\dots+4u+1)(u^{88}+11u^{87}+\dots+20u+1)$
$c_5, c_{11}$	$(u+1)^8(u^3-u^2+1)^2(u^4-u^2+1)^2(u^6-u^5-u^4+2u^3-u+1)^3$ $\cdot (u^{16}-4u^{15}+\dots-8u+2)(u^{18}+3u^{17}+\dots+6u+2)$ $\cdot ((u^{44}-3u^{43}+\dots+14u-1)^2)(u^{50}+11u^{49}+\dots+165u+13)$

## XI. Riley Polynomials

Crossings	Riley Polynomials at each crossing
$c_1, c_{10}$	$(y-1)^8(y^2+y+1)^4(y^3+3y^2+2y-1)^2$ $\cdot ((y^6+y^5+5y^4+6y^2+3y+1)^3)(y^{16}+2y^{15}+\dots+152y+16)$ $\cdot (y^{18}+13y^{17}+\dots-272y+16)(y^{44}+17y^{43}+\dots-3330y+1)^2$ $\cdot (y^{50}+13y^{49}+\dots-3709039y+28561)$
$c_2, c_5, c_7$ $c_{11}$	$(y-1)^8(y^2-y+1)^4(y^3-y^2+2y-1)^2$ $\cdot ((y^6-3y^5+5y^4-4y^3+2y^2-y+1)^3)(y^{16}-6y^{15}+\dots+4y+4)$ $\cdot (y^{18}-7y^{17}+\dots+4y+4)(y^{44}-19y^{43}+\dots-70y+1)^2$ $\cdot (y^{50}-19y^{49}+\dots-5099y+169)$
$c_3, c_8$	$(y-1)^6(y^6+y^5+5y^4+6y^2+3y+1)$ $\cdot (y^8+4y^6+2y^5+2y^4+4y^3+y^2+y+1)^2$ $\cdot (y^8-4y^7+8y^6-6y^5+2y^4-12y^3+29y^2-10y+1)^2$ $\cdot (y^{12}+3y^{11}+\dots+64y+289)(y^{18}-9y^{17}+\dots-480y+64)$ $\cdot (y^{25}-8y^{24}+\dots+y-1)^2$ $\cdot (y^{44}-7y^{43}+\dots-1090962y+73441)^2$
$c_4, c_6, c_9$ $c_{12}$	$((y+1)^8)(y^6-3y^5+\dots-y+1)(y^6+3y^5+\dots+2y^3+1)$ $\cdot (y^8+4y^7+8y^6+6y^5+2y^4+12y^3+29y^2+10y+1)$ $\cdot (y^{12}+3y^{11}+\dots+12y+1)(y^{16}-12y^{15}+\dots-2y+1)$ $\cdot (y^{18}+10y^{17}+\dots+y+1)(y^{50}+21y^{49}+\dots+52y+1)$ $\cdot (y^{88}-39y^{87}+\dots-250y+1)$