


12a<sub>0563</sub> (K12a<sub>0563</sub>)



**Ideals for irreducible components<sup>2</sup> of  $X_{\text{par}}$**

$$I_1^u = \langle -3.21428 \times 10^{21}u^{47} + 3.07657 \times 10^{21}u^{46} + \dots + 1.91294 \times 10^{22}b + 2.10944 \times 10^{22}, \\ 8.01226 \times 10^{20}u^{47} - 8.31592 \times 10^{20}u^{46} + \dots + 9.56472 \times 10^{20}a + 3.53130 \times 10^{21}, u^{48} - u^{47} + \dots + u + 2 \rangle$$

$$I_2^u = \langle -u^2 + b, a - 1, u^{18} + 6u^{16} + \dots + u - 1 \rangle$$

$$I_3^u = \langle b + 1, a^4 + a^3u - 4a^3 - 3a^2u + 5a^2 + 3au - 2a - u + 1, u^2 + 1 \rangle$$

\* 3 irreducible components of  $\dim_{\mathbb{C}} = 0$ , with total 74 representations.

<sup>1</sup>The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/maths/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

<sup>2</sup>All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

**I.**

$$I_1^u = \langle -3.21 \times 10^{21} u^{47} + 3.08 \times 10^{21} u^{46} + \dots + 1.91 \times 10^{22} b + 2.11 \times 10^{22}, 8.01 \times 10^{20} u^{47} - 8.32 \times 10^{20} u^{46} + \dots + 9.56 \times 10^{20} a + 3.53 \times 10^{21}, u^{48} - u^{47} + \dots + u + 2 \rangle$$

**(i) Arc colorings**

$$a_3 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -0.837689u^{47} + 0.869438u^{46} + \dots + 13.2059u - 3.69201 \\ 0.168028u^{47} - 0.160829u^{46} + \dots - 1.56252u - 1.10272 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -0.669662u^{47} + 0.708609u^{46} + \dots + 11.6434u - 4.79473 \\ 0.168028u^{47} - 0.160829u^{46} + \dots - 1.56252u - 1.10272 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -0.573968u^{47} + 0.0603230u^{46} + \dots + 9.04562u + 5.11395 \\ 0.0433537u^{47} + 0.0687272u^{46} + \dots - 4.47263u + 1.08386 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} u \\ u^3 + u \end{pmatrix}$$

$$a_5 = \begin{pmatrix} u^2 + 1 \\ u^4 + 2u^2 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -0.775286u^{47} + 0.102961u^{46} + \dots + 16.4932u + 5.06341 \\ 0.115161u^{47} - 0.190045u^{46} + \dots - 4.11240u + 0.828323 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -0.519611u^{47} + 0.601722u^{46} + \dots + 25.7485u + 2.68654 \\ -0.0513589u^{47} - 0.116669u^{46} + \dots - 1.89721u + 1.51116 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -1.34327u^{47} + 1.07781u^{46} + \dots + 27.0507u + 5.07361 \\ -0.273830u^{47} + 0.198143u^{46} + \dots + 3.26013u - 0.158606 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 1.33343u^{47} - 1.45372u^{46} + \dots - 17.4102u - 8.92856 \\ 0.00796109u^{47} + 0.0832675u^{46} + \dots - 0.966904u + 0.230758 \end{pmatrix}$$

**(ii) Obstruction class = -1**

$$\text{(iii) Cusp Shapes} = \frac{27465448057329986227}{27804404382329290780} u^{47} - \frac{1722649487279789829}{2780440438232929078} u^{46} + \dots - \frac{776992786016705604511}{27804404382329290780} u - \frac{198347232390961135567}{13902202191164645390}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1$	$u^{48} + 17u^{47} + \dots + 35u + 4$
$c_2, c_7$	$u^{48} + u^{47} + \dots - 3u + 2$
$c_3, c_4, c_8$	$u^{48} + u^{47} + \dots - u + 2$
$c_5, c_6, c_{10}$ $c_{11}$	$u^{48} - 2u^{47} + \dots + u + 2$
$c_9, c_{12}$	$u^{48} - 8u^{47} + \dots - 7639u + 1016$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1$	$y^{48} + 37y^{47} + \dots + 12799y + 16$
$c_2, c_7$	$y^{48} + 17y^{47} + \dots + 35y + 4$
$c_3, c_4, c_8$	$y^{48} + 53y^{47} + \dots - 237y + 4$
$c_5, c_6, c_{10}$ $c_{11}$	$y^{48} - 52y^{47} + \dots + 19y + 4$
$c_9, c_{12}$	$y^{48} + 32y^{47} + \dots - 318369y + 1032256$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.279405 + 0.971105I$		
$a = 0.633756 + 0.239746I$	$-4.68544 + 3.19221I$	$-6.37639 - 4.16497I$
$b = 0.165180 + 0.161172I$		
$u = -0.279405 - 0.971105I$		
$a = 0.633756 - 0.239746I$	$-4.68544 - 3.19221I$	$-6.37639 + 4.16497I$
$b = 0.165180 - 0.161172I$		
$u = 0.881286 + 0.378029I$		
$a = -0.468437 + 0.143230I$	$-5.10511 - 9.85415I$	$-11.23414 + 7.48976I$
$b = -0.67254 - 1.34172I$		
$u = 0.881286 - 0.378029I$		
$a = -0.468437 - 0.143230I$	$-5.10511 + 9.85415I$	$-11.23414 - 7.48976I$
$b = -0.67254 + 1.34172I$		
$u = -0.845379 + 0.414724I$		
$a = -0.399315 - 0.163435I$	$1.89498 + 7.08453I$	$-7.53014 - 8.63075I$
$b = -0.586845 + 1.279740I$		
$u = -0.845379 - 0.414724I$		
$a = -0.399315 + 0.163435I$	$1.89498 - 7.08453I$	$-7.53014 + 8.63075I$
$b = -0.586845 - 1.279740I$		
$u = -0.757362 + 0.544501I$		
$a = -0.177555 - 0.142317I$	$-3.80725 + 0.44416I$	$-9.21656 - 2.62442I$
$b = -0.315094 + 1.121340I$		
$u = -0.757362 - 0.544501I$		
$a = -0.177555 + 0.142317I$	$-3.80725 - 0.44416I$	$-9.21656 + 2.62442I$
$b = -0.315094 - 1.121340I$		
$u = 0.801878 + 0.461107I$		
$a = -0.308887 + 0.177382I$	$2.31942 - 3.03168I$	$-5.96588 + 2.35658I$
$b = -0.484403 - 1.202350I$		
$u = 0.801878 - 0.461107I$		
$a = -0.308887 - 0.177382I$	$2.31942 + 3.03168I$	$-5.96588 - 2.35658I$
$b = -0.484403 + 1.202350I$		

Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.094584 + 0.916633I$		
$a = 0.842564 - 0.081921I$	$1.88453 - 1.52925I$	$-1.27440 + 5.26025I$
$b = -0.0039149 - 0.0498200I$		
$u = 0.094584 - 0.916633I$		
$a = 0.842564 + 0.081921I$	$1.88453 + 1.52925I$	$-1.27440 - 5.26025I$
$b = -0.0039149 + 0.0498200I$		
$u = 0.019449 + 1.296200I$		
$a = 0.96626 + 1.66275I$	$-4.87695 + 2.29360I$	0
$b = -0.218022 - 0.686915I$		
$u = 0.019449 - 1.296200I$		
$a = 0.96626 - 1.66275I$	$-4.87695 - 2.29360I$	0
$b = -0.218022 + 0.686915I$		
$u = 0.662333 + 0.127045I$		
$a = -0.823194 + 0.695101I$	$-10.70200 - 3.57800I$	$-16.9686 + 4.2625I$
$b = -1.028930 - 0.838109I$		
$u = 0.662333 - 0.127045I$		
$a = -0.823194 - 0.695101I$	$-10.70200 + 3.57800I$	$-16.9686 - 4.2625I$
$b = -1.028930 + 0.838109I$		
$u = -0.027632 + 1.374070I$		
$a = 0.60190 - 1.62387I$	$3.01436 - 0.56771I$	0
$b = -0.134767 + 0.924503I$		
$u = -0.027632 - 1.374070I$		
$a = 0.60190 + 1.62387I$	$3.01436 + 0.56771I$	0
$b = -0.134767 - 0.924503I$		
$u = 0.218315 + 1.358270I$		
$a = 0.09353 + 2.15802I$	$-5.99306 - 6.70057I$	0
$b = -0.696041 - 1.142180I$		
$u = 0.218315 - 1.358270I$		
$a = 0.09353 - 2.15802I$	$-5.99306 + 6.70057I$	0
$b = -0.696041 + 1.142180I$		

Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.574563 + 0.232091I$ $a = -0.449839 - 0.845197I$ $b = -0.830134 + 0.745730I$	$-2.99542 + 2.92052I$	$-16.1336 - 6.6064I$
$u = -0.574563 - 0.232091I$ $a = -0.449839 + 0.845197I$ $b = -0.830134 - 0.745730I$	$-2.99542 - 2.92052I$	$-16.1336 + 6.6064I$
$u = -0.17592 + 1.40998I$ $a = 0.17221 - 1.93269I$ $b = -0.509228 + 1.232380I$	$2.29935 + 5.57635I$	0
$u = -0.17592 - 1.40998I$ $a = 0.17221 + 1.93269I$ $b = -0.509228 - 1.232380I$	$2.29935 - 5.57635I$	0
$u = 0.09301 + 1.43048I$ $a = 0.33273 + 1.72234I$ $b = -0.243029 - 1.177930I$	$5.11959 - 2.66046I$	0
$u = 0.09301 - 1.43048I$ $a = 0.33273 - 1.72234I$ $b = -0.243029 + 1.177930I$	$5.11959 + 2.66046I$	0
$u = 0.328252 + 0.355874I$ $a = 0.529736 + 1.011980I$ $b = -0.606974 - 0.412729I$	$-0.575619 - 1.189230I$	$-7.30572 + 5.21199I$
$u = 0.328252 - 0.355874I$ $a = 0.529736 - 1.011980I$ $b = -0.606974 + 0.412729I$	$-0.575619 + 1.189230I$	$-7.30572 - 5.21199I$
$u = 0.33907 + 1.49289I$ $a = -0.27071 + 1.81325I$ $b = -0.88608 - 1.67141I$	$0.9224 - 14.2869I$	0
$u = 0.33907 - 1.49289I$ $a = -0.27071 - 1.81325I$ $b = -0.88608 + 1.67141I$	$0.9224 + 14.2869I$	0

Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.31588 + 1.50354I$ $a = -0.22485 - 1.79337I$ $b = -0.80439 + 1.67797I$	$8.10317 + 11.31590I$	0
$u = -0.31588 - 1.50354I$ $a = -0.22485 + 1.79337I$ $b = -0.80439 - 1.67797I$	$8.10317 - 11.31590I$	0
$u = 0.28776 + 1.51291I$ $a = -0.17153 + 1.77103I$ $b = -0.70857 - 1.67398I$	$8.73649 - 7.00574I$	0
$u = 0.28776 - 1.51291I$ $a = -0.17153 - 1.77103I$ $b = -0.70857 + 1.67398I$	$8.73649 + 7.00574I$	0
$u = -0.23601 + 1.52628I$ $a = 0.268656 + 1.050800I$ $b = 0.809501 - 0.930896I$	$3.13587 + 8.05851I$	0
$u = -0.23601 - 1.52628I$ $a = 0.268656 - 1.050800I$ $b = 0.809501 + 0.930896I$	$3.13587 - 8.05851I$	0
$u = -0.24303 + 1.52669I$ $a = -0.09228 - 1.72690I$ $b = -0.55642 + 1.66139I$	$2.98841 + 4.04672I$	0
$u = -0.24303 - 1.52669I$ $a = -0.09228 + 1.72690I$ $b = -0.55642 - 1.66139I$	$2.98841 - 4.04672I$	0
$u = 0.20089 + 1.53586I$ $a = 0.267467 - 1.098980I$ $b = 0.735565 + 1.017990I$	$10.08960 - 5.05548I$	0
$u = 0.20089 - 1.53586I$ $a = 0.267467 + 1.098980I$ $b = 0.735565 - 1.017990I$	$10.08960 + 5.05548I$	0



Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.16329 + 1.54443I$		
$a = 0.262928 + 1.151930I$	$10.46800 + 0.72296I$	0
$b = 0.650504 - 1.106080I$		
$u = -0.16329 - 1.54443I$		
$a = 0.262928 - 1.151930I$	$10.46800 - 0.72296I$	0
$b = 0.650504 + 1.106080I$		
$u = 0.11419 + 1.55675I$		
$a = 0.245810 - 1.222640I$	$4.35322 + 2.28070I$	0
$b = 0.537126 + 1.223150I$		
$u = 0.11419 - 1.55675I$		
$a = 0.245810 + 1.222640I$	$4.35322 - 2.28070I$	0
$b = 0.537126 - 1.223150I$		
$u = 0.276434 + 0.095381I$		
$a = -2.68244 - 2.75338I$	$-8.62006 - 3.30984I$	$-17.2557 + 2.3212I$
$b = -1.136770 + 0.255170I$		
$u = 0.276434 - 0.095381I$		
$a = -2.68244 + 2.75338I$	$-8.62006 + 3.30984I$	$-17.2557 - 2.3212I$
$b = -1.136770 - 0.255170I$		
$u = -0.198964 + 0.041420I$		
$a = -0.39851 - 4.51714I$	$-1.51896 - 1.35228I$	$-14.3368 + 4.1508I$
$b = -0.975718 + 0.209762I$		
$u = -0.198964 - 0.041420I$		
$a = -0.39851 + 4.51714I$	$-1.51896 + 1.35228I$	$-14.3368 - 4.1508I$
$b = -0.975718 - 0.209762I$		

$$\text{II. } I_2^u = \langle -u^2 + b, a - 1, u^{18} + 6u^{16} + \dots + u - 1 \rangle$$

(i) Arc colorings

$$a_3 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} u^2 + 1 \\ u^2 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -u^5 - 2u^3 - u \\ -u^5 - u^3 + u \end{pmatrix}$$

$$a_8 = \begin{pmatrix} u \\ u^3 + u \end{pmatrix}$$

$$a_5 = \begin{pmatrix} u^2 + 1 \\ u^4 + 2u^2 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -u^{11} - 4u^9 - 6u^7 - 6u^5 - 5u^3 - 2u \\ -u^{13} - 5u^{11} - 9u^9 - 8u^7 - 6u^5 - 3u^3 + u \end{pmatrix}$$

$$a_7 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} u^8 + 3u^6 + 3u^4 + 2u^2 + 1 \\ u^8 + 2u^6 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} u^{17} + 6u^{15} + 15u^{13} + 22u^{11} + 23u^9 + 18u^7 + 10u^5 + 4u^3 + u \\ u^{17} + 5u^{15} + 9u^{13} + 8u^{11} + 5u^9 + 2u^7 + u \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes

$$= -4u^{12} - 16u^{10} - 4u^9 - 24u^8 - 12u^7 - 24u^6 - 12u^5 - 20u^4 - 8u^3 - 8u^2 - 4u - 10$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1$	$u^{18} + 12u^{17} + \dots - 5u + 1$
$c_2, c_3, c_4$ $c_7, c_8$	$u^{18} + 6u^{16} + \dots - u - 1$
$c_5, c_6, c_{10}$ $c_{11}$	$(u^6 + u^5 - 3u^4 - 2u^3 + 2u^2 - u - 1)^3$
$c_9, c_{12}$	$(u^6 - u^5 + 3u^4 - 2u^3 + 2u^2 - u - 1)^3$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1$	$y^{18} - 12y^{17} + \dots - 57y + 1$
$c_2, c_3, c_4$ $c_7, c_8$	$y^{18} + 12y^{17} + \dots - 5y + 1$
$c_5, c_6, c_{10}$ $c_{11}$	$(y^6 - 7y^5 + 17y^4 - 16y^3 + 6y^2 - 5y + 1)^3$
$c_9, c_{12}$	$(y^6 + 5y^5 + 9y^4 + 4y^3 - 6y^2 - 5y + 1)^3$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_2^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.637469 + 0.735789I$ $a = 1.00000$ $b = -0.135019 - 0.938086I$	$2.96024 - 1.97241I$	$-4.57572 + 3.68478I$
$u = -0.637469 - 0.735789I$ $a = 1.00000$ $b = -0.135019 + 0.938086I$	$2.96024 + 1.97241I$	$-4.57572 - 3.68478I$
$u = 0.639652 + 0.826288I$ $a = 1.00000$ $b = -0.273597 + 1.057070I$	$-3.69558 + 4.59213I$	$-8.58114 - 3.20482I$
$u = 0.639652 - 0.826288I$ $a = 1.00000$ $b = -0.273597 - 1.057070I$	$-3.69558 - 4.59213I$	$-8.58114 + 3.20482I$
$u = -0.182330 + 1.048680I$ $a = 1.00000$ $b = -1.066490 - 0.382411I$	$-0.738851$	$-13.41678 + 0.I$
$u = -0.182330 - 1.048680I$ $a = 1.00000$ $b = -1.066490 + 0.382411I$	$-0.738851$	$-13.41678 + 0.I$
$u = 0.667042 + 0.642083I$ $a = 1.00000$ $b = 0.032675 + 0.856592I$	$2.96024 - 1.97241I$	$-4.57572 + 3.68478I$
$u = 0.667042 - 0.642083I$ $a = 1.00000$ $b = 0.032675 - 0.856592I$	$2.96024 + 1.97241I$	$-4.57572 - 3.68478I$
$u = -0.724676 + 0.565991I$ $a = 1.00000$ $b = 0.204809 - 0.820320I$	$-3.69558 + 4.59213I$	$-8.58114 - 3.20482I$
$u = -0.724676 - 0.565991I$ $a = 1.00000$ $b = 0.204809 + 0.820320I$	$-3.69558 - 4.59213I$	$-8.58114 + 3.20482I$

Solutions to $I_2^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.313436 + 1.137860I$ $a = 1.00000$ $b = -1.196490 + 0.713294I$	-7.66009	$-12.26950 + 0.I$
$u = 0.313436 - 1.137860I$ $a = 1.00000$ $b = -1.196490 - 0.713294I$	-7.66009	$-12.26950 + 0.I$
$u = -0.626873$ $a = 1.00000$ $b = 0.392969$	-7.66009	-12.2690
$u = -0.029572 + 1.377870I$ $a = 1.00000$ $b = -1.89766 - 0.08149I$	$2.96024 + 1.97241I$	$-4.57572 - 3.68478I$
$u = -0.029572 - 1.377870I$ $a = 1.00000$ $b = -1.89766 + 0.08149I$	$2.96024 - 1.97241I$	$-4.57572 + 3.68478I$
$u = 0.085024 + 1.392280I$ $a = 1.00000$ $b = -1.93121 + 0.23675I$	$-3.69558 - 4.59213I$	$-8.58114 + 3.20482I$
$u = 0.085024 - 1.392280I$ $a = 1.00000$ $b = -1.93121 - 0.23675I$	$-3.69558 + 4.59213I$	$-8.58114 - 3.20482I$
$u = 0.364659$ $a = 1.00000$ $b = 0.132976$	-0.738851	-13.4170

$$\text{III. } I_3^u = \langle b + 1, a^3u - 3a^2u + \cdots - 2a + 1, u^2 + 1 \rangle$$

(i) Arc colorings

$$a_3 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_2 = \begin{pmatrix} a \\ -1 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} a - 1 \\ -1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -a^2u + 2au - u \\ au \end{pmatrix}$$

$$a_8 = \begin{pmatrix} u \\ 0 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -a^2u + 2au - u \\ -a^2u + 3au - u \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -au + u \\ u \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -a^3 + 3a^2 - 2a \\ a^2 - a - 1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} a^3u - 3a^2u - a^2 + 3au + 2a - u \\ a^3u + a^3 - 3a^2u - 3a^2 + 3au + 3a - u - 1 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes =  $-4a^2 - 4au + 8a + 4u - 12$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
$c_1$	$(u - 1)^8$
$c_2, c_3, c_4$ $c_7, c_8$	$(u^2 + 1)^4$
$c_5, c_6, c_{10}$ $c_{11}$	$u^8 - 5u^6 + 7u^4 - 2u^2 + 1$
$c_9$	$(u^4 + u^3 + u^2 + 1)^2$
$c_{12}$	$(u^4 - u^3 + u^2 + 1)^2$



(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
$c_1$	$(y - 1)^8$
$c_2, c_3, c_4$ $c_7, c_8$	$(y + 1)^8$
$c_5, c_6, c_{10}$ $c_{11}$	$(y^4 - 5y^3 + 7y^2 - 2y + 1)^2$
$c_9, c_{12}$	$(y^4 + y^3 + 3y^2 + 2y + 1)^2$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_3^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.000000I$ $a = 0.088708 - 0.851808I$ $b = -1.00000$	$-6.79074 + 3.16396I$	$-11.82674 - 2.56480I$
$u = 1.000000I$ $a = 0.279658 + 0.351808I$ $b = -1.00000$	$0.21101 - 1.41510I$	$-8.17326 + 4.90874I$
$u = 1.000000I$ $a = 1.72034 + 0.35181I$ $b = -1.00000$	$0.21101 + 1.41510I$	$-8.17326 - 4.90874I$
$u = 1.000000I$ $a = 1.91129 - 0.85181I$ $b = -1.00000$	$-6.79074 - 3.16396I$	$-11.82674 + 2.56480I$
$u = -1.000000I$ $a = 0.088708 + 0.851808I$ $b = -1.00000$	$-6.79074 - 3.16396I$	$-11.82674 + 2.56480I$
$u = -1.000000I$ $a = 0.279658 - 0.351808I$ $b = -1.00000$	$0.21101 + 1.41510I$	$-8.17326 - 4.90874I$
$u = -1.000000I$ $a = 1.72034 - 0.35181I$ $b = -1.00000$	$0.21101 - 1.41510I$	$-8.17326 + 4.90874I$
$u = -1.000000I$ $a = 1.91129 + 0.85181I$ $b = -1.00000$	$-6.79074 + 3.16396I$	$-11.82674 - 2.56480I$

#### IV. u-Polynomials

Crossings	u-Polynomials at each crossing
$c_1$	$((u-1)^8)(u^{18} + 12u^{17} + \dots - 5u + 1)(u^{48} + 17u^{47} + \dots + 35u + 4)$
$c_2, c_7$	$((u^2 + 1)^4)(u^{18} + 6u^{16} + \dots - u - 1)(u^{48} + u^{47} + \dots - 3u + 2)$
$c_3, c_4, c_8$	$((u^2 + 1)^4)(u^{18} + 6u^{16} + \dots - u - 1)(u^{48} + u^{47} + \dots - u + 2)$
$c_5, c_6, c_{10}$ $c_{11}$	$(u^6 + u^5 - 3u^4 - 2u^3 + 2u^2 - u - 1)^3(u^8 - 5u^6 + 7u^4 - 2u^2 + 1)$ $\cdot (u^{48} - 2u^{47} + \dots + u + 2)$
$c_9$	$(u^4 + u^3 + u^2 + 1)^2(u^6 - u^5 + 3u^4 - 2u^3 + 2u^2 - u - 1)^3$ $\cdot (u^{48} - 8u^{47} + \dots - 7639u + 1016)$
$c_{12}$	$(u^4 - u^3 + u^2 + 1)^2(u^6 - u^5 + 3u^4 - 2u^3 + 2u^2 - u - 1)^3$ $\cdot (u^{48} - 8u^{47} + \dots - 7639u + 1016)$

### V. Riley Polynomials

Crossings	Riley Polynomials at each crossing
$c_1$	$((y-1)^8)(y^{18} - 12y^{17} + \dots - 57y + 1)$ $\cdot (y^{48} + 37y^{47} + \dots + 12799y + 16)$
$c_2, c_7$	$((y+1)^8)(y^{18} + 12y^{17} + \dots - 5y + 1)(y^{48} + 17y^{47} + \dots + 35y + 4)$
$c_3, c_4, c_8$	$((y+1)^8)(y^{18} + 12y^{17} + \dots - 5y + 1)(y^{48} + 53y^{47} + \dots - 237y + 4)$
$c_5, c_6, c_{10}$ $c_{11}$	$(y^4 - 5y^3 + 7y^2 - 2y + 1)^2(y^6 - 7y^5 + 17y^4 - 16y^3 + 6y^2 - 5y + 1)^3$ $\cdot (y^{48} - 52y^{47} + \dots + 19y + 4)$
$c_9, c_{12}$	$(y^4 + y^3 + 3y^2 + 2y + 1)^2(y^6 + 5y^5 + 9y^4 + 4y^3 - 6y^2 - 5y + 1)^3$ $\cdot (y^{48} + 32y^{47} + \dots - 318369y + 1032256)$