


7₆ (K7a₂)



Ideals for irreducible components² of X_{par}

$$I_1^u = \langle u^9 - u^8 - 2u^7 + 3u^6 + u^5 - 3u^4 + 2u^3 - u + 1 \rangle$$

* 1 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 9 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/math/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$\text{I. } I_1^u = \langle u^9 - u^8 - 2u^7 + 3u^6 + u^5 - 3u^4 + 2u^3 - u + 1 \rangle$$

(i) Arc colorings

$$a_1 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} 1 \\ -u^2 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} u \\ -u^3 + u \end{pmatrix}$$

$$a_6 = \begin{pmatrix} u^3 \\ -u^3 + u \end{pmatrix}$$

$$a_5 = \begin{pmatrix} u^6 - u^4 + 1 \\ -u^6 + 2u^4 - u^2 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -u^3 \\ u^5 - u^3 + u \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -u^3 \\ u^5 - u^3 + u \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes = $4u^7 - 8u^5 + 4u^4 + 8u^3 - 4u^2 + 4u - 2$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^9 + 5u^8 + 12u^7 + 15u^6 + 9u^5 - u^4 - 4u^3 - 2u^2 + u + 1$
c_2, c_5	$u^9 - u^8 + 2u^7 - u^6 + 3u^5 - u^4 + 2u^3 + u + 1$
c_3, c_7	$u^9 - u^8 - 2u^7 + 3u^6 + u^5 - 3u^4 + 2u^3 - u + 1$
c_4, c_6	$u^9 - 3u^8 + 8u^7 - 13u^6 + 17u^5 - 17u^4 + 12u^3 - 6u^2 + u + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^9 - y^8 + 12y^7 - 7y^6 + 37y^5 + y^4 - 10y^2 + 5y - 1$
c_2, c_5	$y^9 + 3y^8 + 8y^7 + 13y^6 + 17y^5 + 17y^4 + 12y^3 + 6y^2 + y - 1$
c_3, c_7	$y^9 - 5y^8 + 12y^7 - 15y^6 + 9y^5 + y^4 - 4y^3 + 2y^2 + y - 1$
c_4, c_6	$y^9 + 7y^8 + 20y^7 + 25y^6 + 5y^5 - 15y^4 + 22y^2 + 13y - 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.772920 + 0.510351I$	$1.78344 - 2.09337I$	$0.51499 + 4.16283I$
$u = 0.772920 - 0.510351I$	$1.78344 + 2.09337I$	$0.51499 - 4.16283I$
$u = -0.825933$	-1.19845	-8.65230
$u = -1.173910 + 0.391555I$	$-4.37135 + 1.33617I$	$-7.28409 - 0.70175I$
$u = -1.173910 - 0.391555I$	$-4.37135 - 1.33617I$	$-7.28409 + 0.70175I$
$u = 0.141484 + 0.739668I$	$-0.61694 + 2.45442I$	$-2.32792 - 2.91298I$
$u = 0.141484 - 0.739668I$	$-0.61694 - 2.45442I$	$-2.32792 + 2.91298I$
$u = 1.172470 + 0.500383I$	$-3.59813 - 7.08493I$	$-5.57680 + 5.91335I$
$u = 1.172470 - 0.500383I$	$-3.59813 + 7.08493I$	$-5.57680 - 5.91335I$

II. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1	$u^9 + 5u^8 + 12u^7 + 15u^6 + 9u^5 - u^4 - 4u^3 - 2u^2 + u + 1$
c_2, c_5	$u^9 - u^8 + 2u^7 - u^6 + 3u^5 - u^4 + 2u^3 + u + 1$
c_3, c_7	$u^9 - u^8 - 2u^7 + 3u^6 + u^5 - 3u^4 + 2u^3 - u + 1$
c_4, c_6	$u^9 - 3u^8 + 8u^7 - 13u^6 + 17u^5 - 17u^4 + 12u^3 - 6u^2 + u + 1$

III. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1	$y^9 - y^8 + 12y^7 - 7y^6 + 37y^5 + y^4 - 10y^2 + 5y - 1$
c_2, c_5	$y^9 + 3y^8 + 8y^7 + 13y^6 + 17y^5 + 17y^4 + 12y^3 + 6y^2 + y - 1$
c_3, c_7	$y^9 - 5y^8 + 12y^7 - 15y^6 + 9y^5 + y^4 - 4y^3 + 2y^2 + y - 1$
c_4, c_6	$y^9 + 7y^8 + 20y^7 + 25y^6 + 5y^5 - 15y^4 + 22y^2 + 13y - 1$