


$12a_{0681}$ ($K12a_{0681}$)



Ideals for irreducible components² of X_{par}

$$I_1^u = \langle u^{76} - 35u^{74} + \dots + b - 1, \ u^{75} + u^{74} + \dots + a + 1, \ u^{77} + 2u^{76} + \dots - u - 1 \rangle$$

$$I_2^u = \langle b, \ u^4 - u^3 - u^2 + a + u, \ u^5 - u^4 - 2u^3 + u^2 + u + 1 \rangle$$

* 2 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 82 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/math/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$I_1^u = \langle u^{76} - 35u^{74} + \cdots + b - 1, \ u^{75} + u^{74} + \cdots + a + 1, \ u^{77} + 2u^{76} + \cdots - u - 1 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_3 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_4 &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix} \\ a_5 &= \begin{pmatrix} -u^2 + 1 \\ -u^4 + 2u^2 \end{pmatrix} \\ a_7 &= \begin{pmatrix} -u^{75} - u^{74} + \cdots - 5u - 1 \\ -u^{76} + 35u^{74} + \cdots + u + 1 \end{pmatrix} \\ a_2 &= \begin{pmatrix} u^{11} - 6u^9 + 12u^7 - 8u^5 + u^3 - 2u \\ -u^{11} + 5u^9 - 8u^7 + 3u^5 + u^3 + u \end{pmatrix} \\ a_8 &= \begin{pmatrix} -u^{75} - u^{74} + \cdots - 10u^2 - 4u \\ u^{76} - 35u^{74} + \cdots + 9u^2 - 1 \end{pmatrix} \\ a_1 &= \begin{pmatrix} -u^9 + 4u^7 - 5u^5 + 2u^3 - u \\ -u^{11} + 5u^9 - 8u^7 + 3u^5 + u^3 + u \end{pmatrix} \\ a_{10} &= \begin{pmatrix} u^5 - 2u^3 + u \\ u^7 - 3u^5 + 2u^3 + u \end{pmatrix} \\ a_9 &= \begin{pmatrix} u^{13} - 6u^{11} + 13u^9 - 12u^7 + 6u^5 - 4u^3 + u \\ u^{15} - 7u^{13} + 18u^{11} - 19u^9 + 6u^7 - 2u^5 + 4u^3 + u \end{pmatrix} \\ a_6 &= \begin{pmatrix} -u^{75} - u^{74} + \cdots - 10u^2 - 4u \\ u^{37} - 17u^{35} + \cdots + 6u^2 + u \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes = $2u^{76} + 4u^{75} + \cdots - 9u - 6$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{77} + 33u^{76} + \cdots - 7680u - 1024$
c_2, c_7	$u^{77} + u^{76} + \cdots - 96u - 32$
c_3, c_4, c_{11}	$u^{77} + 2u^{76} + \cdots - u - 1$
c_5, c_6, c_8	$u^{77} - 6u^{76} + \cdots + u - 1$
c_9, c_{12}	$u^{77} + 12u^{76} + \cdots + 1689u + 73$
c_{10}	$u^{77} - 6u^{76} + \cdots + 813u - 935$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{77} + 13y^{76} + \cdots + 11141120y - 1048576$
c_2, c_7	$y^{77} + 33y^{76} + \cdots - 7680y - 1024$
c_3, c_4, c_{11}	$y^{77} - 72y^{76} + \cdots + 7y - 1$
c_5, c_6, c_8	$y^{77} - 68y^{76} + \cdots - 7y - 1$
c_9, c_{12}	$y^{77} + 60y^{76} + \cdots + 148363y - 5329$
c_{10}	$y^{77} - 24y^{76} + \cdots + 24711039y - 874225$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.118910 + 0.179695I$	$-4.41825 - 2.77630I$	0
$a = -0.440072 + 1.213090I$		
$b = 0.513943 - 0.982821I$		
$u = -1.118910 - 0.179695I$	$-4.41825 + 2.77630I$	0
$a = -0.440072 - 1.213090I$		
$b = 0.513943 + 0.982821I$		
$u = 0.388896 + 0.697285I$	$-6.76637 - 11.69120I$	$-8.55592 + 8.80415I$
$a = 2.58400 + 0.08701I$		
$b = -0.694399 + 1.197770I$		
$u = 0.388896 - 0.697285I$	$-6.76637 + 11.69120I$	$-8.55592 - 8.80415I$
$a = 2.58400 - 0.08701I$		
$b = -0.694399 - 1.197770I$		
$u = 0.467253 + 0.642938I$	$-11.71880 - 2.13120I$	$-12.50690 + 3.22543I$
$a = -1.072070 + 0.222088I$		
$b = 0.025679 - 1.363990I$		
$u = 0.467253 - 0.642938I$	$-11.71880 + 2.13120I$	$-12.50690 - 3.22543I$
$a = -1.072070 - 0.222088I$		
$b = 0.025679 + 1.363990I$		
$u = -1.199980 + 0.164041I$	$-0.198946 + 0.491194I$	0
$a = 0.765140 - 0.964895I$		
$b = -0.662389 + 0.724093I$		
$u = -1.199980 - 0.164041I$	$-0.198946 - 0.491194I$	0
$a = 0.765140 + 0.964895I$		
$b = -0.662389 - 0.724093I$		
$u = 0.550317 + 0.554367I$	$-7.39452 + 7.48341I$	$-10.14800 - 2.91847I$
$a = -0.557825 - 0.493865I$		
$b = 0.660792 + 1.201360I$		
$u = 0.550317 - 0.554367I$	$-7.39452 - 7.48341I$	$-10.14800 + 2.91847I$
$a = -0.557825 + 0.493865I$		
$b = 0.660792 - 1.201360I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.382467 + 0.674572I$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$a = -2.43549 - 0.42583I$	$-1.41933 - 7.47214I$	$-5.05697 + 8.34642I$
$b = 0.583029 - 1.060690I$		
$u = 0.382467 - 0.674572I$		
$a = -2.43549 + 0.42583I$	$-1.41933 + 7.47214I$	$-5.05697 - 8.34642I$
$b = 0.583029 + 1.060690I$		
$u = -0.391946 + 0.660763I$		
$a = 1.07868 - 1.47152I$	$-4.41478 + 5.37995I$	$-7.77105 - 5.68914I$
$b = -1.058370 + 0.462731I$		
$u = -0.391946 - 0.660763I$		
$a = 1.07868 + 1.47152I$	$-4.41478 - 5.37995I$	$-7.77105 + 5.68914I$
$b = -1.058370 - 0.462731I$		
$u = 0.392387 + 0.638561I$		
$a = 1.89360 + 0.81444I$	$-3.46473 - 2.75353I$	$-9.00366 + 4.34907I$
$b = -0.337099 + 0.970309I$		
$u = 0.392387 - 0.638561I$		
$a = 1.89360 - 0.81444I$	$-3.46473 + 2.75353I$	$-9.00366 - 4.34907I$
$b = -0.337099 - 0.970309I$		
$u = -1.25296$		
$a = -0.489598$	-2.70327	0
$b = 0.440012$		
$u = 1.247280 + 0.146275I$		
$a = -2.07717 - 0.81765I$	$-3.52703 - 1.49966I$	0
$b = 0.556055 - 0.712579I$		
$u = 1.247280 - 0.146275I$		
$a = -2.07717 + 0.81765I$	$-3.52703 + 1.49966I$	0
$b = 0.556055 + 0.712579I$		
$u = 0.512426 + 0.536307I$		
$a = 0.551321 + 0.026348I$	$-1.97887 + 3.43479I$	$-6.68994 - 2.31052I$
$b = -0.538012 - 1.042190I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.512426 - 0.536307I$		
$a = 0.551321 - 0.026348I$	$-1.97887 - 3.43479I$	$-6.68994 + 2.31052I$
$b = -0.538012 + 1.042190I$		
$u = -0.488346 + 0.553137I$		
$a = -1.48960 + 0.38906I$	$-4.84664 - 1.37205I$	$-9.12609 - 0.79115I$
$b = 1.044170 + 0.400052I$		
$u = -0.488346 - 0.553137I$		
$a = -1.48960 - 0.38906I$	$-4.84664 + 1.37205I$	$-9.12609 + 0.79115I$
$b = 1.044170 - 0.400052I$		
$u = 1.254630 + 0.207259I$		
$a = 2.05786 + 0.27148I$	$-0.73681 - 5.52218I$	0
$b = -0.637359 + 0.904548I$		
$u = 1.254630 - 0.207259I$		
$a = 2.05786 - 0.27148I$	$-0.73681 + 5.52218I$	0
$b = -0.637359 - 0.904548I$		
$u = 0.453993 + 0.561890I$		
$a = 0.036223 + 0.618083I$	$-3.76621 - 1.14578I$	$-10.36985 + 2.95072I$
$b = 0.244719 + 0.964800I$		
$u = 0.453993 - 0.561890I$		
$a = 0.036223 - 0.618083I$	$-3.76621 + 1.14578I$	$-10.36985 - 2.95072I$
$b = 0.244719 - 0.964800I$		
$u = -0.345053 + 0.634395I$		
$a = -0.540936 + 1.304630I$	$0.27587 + 2.54213I$	$-1.22958 - 3.86119I$
$b = 0.687430 - 0.491293I$		
$u = -0.345053 - 0.634395I$		
$a = -0.540936 - 1.304630I$	$0.27587 - 2.54213I$	$-1.22958 + 3.86119I$
$b = 0.687430 + 0.491293I$		
$u = -1.266550 + 0.180413I$		
$a = -1.16776 + 0.80875I$	$-3.88660 + 3.89208I$	0
$b = 0.835295 - 0.561310I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.266550 - 0.180413I$		
$a = -1.16776 - 0.80875I$	$-3.88660 - 3.89208I$	0
$b = 0.835295 + 0.561310I$		
$u = -0.263295 + 0.667225I$		
$a = -0.30263 - 1.80362I$	$-3.32754 + 0.27451I$	$-8.04091 - 0.77276I$
$b = -0.322531 + 0.936487I$		
$u = -0.263295 - 0.667225I$		
$a = -0.30263 + 1.80362I$	$-3.32754 - 0.27451I$	$-8.04091 + 0.77276I$
$b = -0.322531 - 0.936487I$		
$u = 1.268150 + 0.243748I$		
$a = -2.03646 + 0.04790I$	$-5.48995 - 9.38623I$	0
$b = 0.637081 - 1.071140I$		
$u = 1.268150 - 0.243748I$		
$a = -2.03646 - 0.04790I$	$-5.48995 + 9.38623I$	0
$b = 0.637081 + 1.071140I$		
$u = -0.623273 + 0.290139I$		
$a = -0.841020 - 0.501914I$	$-4.77164 + 3.23057I$	$-10.85312 - 4.67428I$
$b = 0.400346 + 1.029660I$		
$u = -0.623273 - 0.290139I$		
$a = -0.841020 + 0.501914I$	$-4.77164 - 3.23057I$	$-10.85312 + 4.67428I$
$b = 0.400346 - 1.029660I$		
$u = -0.076390 + 0.675447I$		
$a = 2.15894 + 1.81721I$	$-1.33875 + 6.04370I$	$-3.74417 - 6.13811I$
$b = -0.585185 - 1.035670I$		
$u = -0.076390 - 0.675447I$		
$a = 2.15894 - 1.81721I$	$-1.33875 - 6.04370I$	$-3.74417 + 6.13811I$
$b = -0.585185 + 1.035670I$		
$u = 1.352570 + 0.042876I$		
$a = -0.466208 - 1.294710I$	$-5.20171 - 1.91816I$	0
$b = 0.174346 - 0.892035I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.352570 - 0.042876I$		
$a = -0.466208 + 1.294710I$	$-5.20171 + 1.91816I$	0
$b = 0.174346 + 0.892035I$		
$u = -1.35955$		
$a = 0.837136$	-7.10233	0
$b = -0.928982$		
$u = -0.040790 + 0.631699I$		
$a = -2.35095 - 1.38726I$	$3.22737 + 2.45907I$	$2.41719 - 4.22669I$
$b = 0.630731 + 0.823161I$		
$u = -0.040790 - 0.631699I$		
$a = -2.35095 + 1.38726I$	$3.22737 - 2.45907I$	$2.41719 + 4.22669I$
$b = 0.630731 - 0.823161I$		
$u = -0.385095 + 0.464622I$		
$a = 0.995000 - 0.148955I$	$-0.270419 + 0.973560I$	$-2.30158 - 4.17778I$
$b = -0.526774 - 0.405932I$		
$u = -0.385095 - 0.464622I$		
$a = 0.995000 + 0.148955I$	$-0.270419 - 0.973560I$	$-2.30158 + 4.17778I$
$b = -0.526774 + 0.405932I$		
$u = 0.033568 + 0.581274I$		
$a = 2.74191 + 0.87771I$	$0.094766 - 1.112510I$	$-0.414963 + 0.658190I$
$b = -0.687117 - 0.568792I$		
$u = 0.033568 - 0.581274I$		
$a = 2.74191 - 0.87771I$	$0.094766 + 1.112510I$	$-0.414963 - 0.658190I$
$b = -0.687117 + 0.568792I$		
$u = 1.39972 + 0.25277I$		
$a = 0.700734 - 0.638857I$	$-8.62598 - 3.61399I$	0
$b = 0.251532 + 0.891469I$		
$u = 1.39972 - 0.25277I$		
$a = 0.700734 + 0.638857I$	$-8.62598 + 3.61399I$	0
$b = 0.251532 - 0.891469I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.42729 + 0.06631I$	$-11.08620 - 4.31379I$	0
$a = 0.417873 + 0.696599I$		
$b = -0.374341 + 1.178740I$		
$u = 1.42729 - 0.06631I$	$-11.08620 + 4.31379I$	0
$a = 0.417873 - 0.696599I$		
$b = -0.374341 - 1.178740I$		
$u = 1.43420 + 0.20084I$	$-6.06731 - 3.57322I$	0
$a = -0.399133 - 0.452591I$		
$b = 0.643759 - 0.302963I$		
$u = 1.43420 - 0.20084I$	$-6.06731 + 3.57322I$	0
$a = -0.399133 + 0.452591I$		
$b = 0.643759 + 0.302963I$		
$u = 1.43740 + 0.24134I$	$-5.45133 - 5.75261I$	0
$a = -0.080952 + 0.770503I$		
$b = -0.743107 - 0.462573I$		
$u = 1.43740 - 0.24134I$	$-5.45133 + 5.75261I$	0
$a = -0.080952 - 0.770503I$		
$b = -0.743107 + 0.462573I$		
$u = -1.45356 + 0.23958I$	$-9.40303 + 5.97378I$	0
$a = -1.51331 + 1.62581I$		
$b = 0.362141 + 1.020760I$		
$u = -1.45356 - 0.23958I$	$-9.40303 - 5.97378I$	0
$a = -1.51331 - 1.62581I$		
$b = 0.362141 - 1.020760I$		
$u = -1.46014 + 0.20569I$	$-9.90965 + 3.96691I$	0
$a = -0.29342 + 1.41422I$		
$b = -0.220303 + 1.036640I$		
$u = -1.46014 - 0.20569I$	$-9.90965 - 3.96691I$	0
$a = -0.29342 - 1.41422I$		
$b = -0.220303 - 1.036640I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.45430 + 0.25332I$		
$a = 1.82596 - 1.39983I$	$-7.32899 + 10.86290I$	0
$b = -0.594804 - 1.088050I$		
$u = -1.45430 - 0.25332I$		
$a = 1.82596 + 1.39983I$	$-7.32899 - 10.86290I$	0
$b = -0.594804 + 1.088050I$		
$u = 1.45612 + 0.24706I$		
$a = -0.082844 - 0.995707I$	$-10.36290 - 8.70087I$	0
$b = 1.093950 + 0.480855I$		
$u = 1.45612 - 0.24706I$		
$a = -0.082844 + 0.995707I$	$-10.36290 + 8.70087I$	0
$b = 1.093950 - 0.480855I$		
$u = -1.46657 + 0.18641I$		
$a = -0.067219 - 0.960194I$	$-8.31617 - 0.82132I$	0
$b = 0.502448 - 1.083330I$		
$u = -1.46657 - 0.18641I$		
$a = -0.067219 + 0.960194I$	$-8.31617 + 0.82132I$	0
$b = 0.502448 + 1.083330I$		
$u = 1.46582 + 0.19639I$		
$a = 0.512261 + 0.674143I$	$-11.11540 - 1.35977I$	0
$b = -1.098560 + 0.367386I$		
$u = 1.46582 - 0.19639I$		
$a = 0.512261 - 0.674143I$	$-11.11540 + 1.35977I$	0
$b = -1.098560 - 0.367386I$		
$u = -1.45965 + 0.26153I$		
$a = -1.90826 + 1.20800I$	$-12.7169 + 15.1904I$	0
$b = 0.712807 + 1.210690I$		
$u = -1.45965 - 0.26153I$		
$a = -1.90826 - 1.20800I$	$-12.7169 - 15.1904I$	0
$b = 0.712807 - 1.210690I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.48110 + 0.17882I$		
$a = 0.014011 + 0.628968I$	$-13.9447 - 4.8720I$	0
$b = -0.647796 + 1.238820I$		
$u = -1.48110 - 0.17882I$		
$a = 0.014011 - 0.628968I$	$-13.9447 + 4.8720I$	0
$b = -0.647796 - 1.238820I$		
$u = -1.47901 + 0.22669I$		
$a = 0.999876 - 0.986609I$	$-18.0111 + 5.2978I$	0
$b = -0.050785 - 1.399520I$		
$u = -1.47901 - 0.22669I$		
$a = 0.999876 + 0.986609I$	$-18.0111 - 5.2978I$	0
$b = -0.050785 + 1.399520I$		
$u = -0.303521 + 0.301278I$		
$a = 1.018990 - 0.220898I$	$-0.201500 + 0.933792I$	$-4.38363 - 6.63917I$
$b = -0.312600 - 0.550494I$		
$u = -0.303521 - 0.301278I$		
$a = 1.018990 + 0.220898I$	$-0.201500 - 0.933792I$	$-4.38363 + 6.63917I$
$b = -0.312600 + 0.550494I$		
$u = 0.278524$		
$a = -2.80564$	-2.11486	-4.47090
$b = 0.551515$		

$$\text{III. } I_2^u = \langle b, u^4 - u^3 - u^2 + a + u, u^5 - u^4 - 2u^3 + u^2 + u + 1 \rangle$$

(i) Arc colorings

$$a_3 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -u^2 + 1 \\ -u^4 + 2u^2 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -u^4 + u^3 + u^2 - u \\ 0 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} -u^4 + u^3 + u^2 - u \\ 0 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} u^4 - u^2 - 1 \\ u^4 - 2u^2 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} u^2 - 1 \\ u^4 - 2u^2 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -u^4 + u^3 - u + 1 \\ -u^4 + 2u^2 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = $5u^3 - u^2 - 8u - 9$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_2, c_7	u^5
c_3, c_4	$u^5 - u^4 - 2u^3 + u^2 + u + 1$
c_5, c_6	$(u - 1)^5$
c_8	$(u + 1)^5$
c_9	$u^5 - u^4 + 2u^3 - u^2 + u - 1$
c_{10}	$u^5 - 3u^4 + 4u^3 - u^2 - u + 1$
c_{11}	$u^5 + u^4 - 2u^3 - u^2 + u - 1$
c_{12}	$u^5 + u^4 + 2u^3 + u^2 + u + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_7	y^5
c_3, c_4, c_{11}	$y^5 - 5y^4 + 8y^3 - 3y^2 - y - 1$
c_5, c_6, c_8	$(y - 1)^5$
c_9, c_{12}	$y^5 + 3y^4 + 4y^3 + y^2 - y - 1$
c_{10}	$y^5 - y^4 + 8y^3 - 3y^2 + 3y - 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.21774$		
$a = -1.30408$	-4.04602	-9.76980
$b = 0$		
$u = -0.309916 + 0.549911I$		
$a = 0.428550 - 1.039280I$	$-1.97403 + 1.53058I$	$-5.05737 - 4.09764I$
$b = 0$		
$u = -0.309916 - 0.549911I$		
$a = 0.428550 + 1.039280I$	$-1.97403 - 1.53058I$	$-5.05737 + 4.09764I$
$b = 0$		
$u = 1.41878 + 0.21917I$		
$a = -0.276511 - 0.728237I$	$-7.51750 - 4.40083I$	$-9.05774 + 4.18967I$
$b = 0$		
$u = 1.41878 - 0.21917I$		
$a = -0.276511 + 0.728237I$	$-7.51750 + 4.40083I$	$-9.05774 - 4.18967I$
$b = 0$		

III. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1	$u^5(u^{77} + 33u^{76} + \dots - 7680u - 1024)$
c_2, c_7	$u^5(u^{77} + u^{76} + \dots - 96u - 32)$
c_3, c_4	$(u^5 - u^4 - 2u^3 + u^2 + u + 1)(u^{77} + 2u^{76} + \dots - u - 1)$
c_5, c_6	$((u - 1)^5)(u^{77} - 6u^{76} + \dots + u - 1)$
c_8	$((u + 1)^5)(u^{77} - 6u^{76} + \dots + u - 1)$
c_9	$(u^5 - u^4 + 2u^3 - u^2 + u - 1)(u^{77} + 12u^{76} + \dots + 1689u + 73)$
c_{10}	$(u^5 - 3u^4 + 4u^3 - u^2 - u + 1)(u^{77} - 6u^{76} + \dots + 813u - 935)$
c_{11}	$(u^5 + u^4 - 2u^3 - u^2 + u - 1)(u^{77} + 2u^{76} + \dots - u - 1)$
c_{12}	$(u^5 + u^4 + 2u^3 + u^2 + u + 1)(u^{77} + 12u^{76} + \dots + 1689u + 73)$

IV. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1	$y^5(y^{77} + 13y^{76} + \dots + 1.11411 \times 10^7 y - 1048576)$
c_2, c_7	$y^5(y^{77} + 33y^{76} + \dots - 7680y - 1024)$
c_3, c_4, c_{11}	$(y^5 - 5y^4 + 8y^3 - 3y^2 - y - 1)(y^{77} - 72y^{76} + \dots + 7y - 1)$
c_5, c_6, c_8	$((y - 1)^5)(y^{77} - 68y^{76} + \dots - 7y - 1)$
c_9, c_{12}	$(y^5 + 3y^4 + 4y^3 + y^2 - y - 1)(y^{77} + 60y^{76} + \dots + 148363y - 5329)$
c_{10}	$(y^5 - y^4 + 8y^3 - 3y^2 + 3y - 1) \cdot (y^{77} - 24y^{76} + \dots + 24711039y - 874225)$