


$12a_{1153}$ ($K12a_{1153}$)



Ideals for irreducible components² of X_{par}

$$\begin{aligned}
 I_1^u &= \langle 7.92523 \times 10^{127} u^{89} + 3.07371 \times 10^{128} u^{88} + \dots + 5.59035 \times 10^{128} b + 1.35846 \times 10^{130}, \\
 &\quad - 2.38182 \times 10^{128} u^{89} + 6.32318 \times 10^{129} u^{88} + \dots + 6.42890 \times 10^{129} a + 1.54116 \times 10^{131}, \\
 &\quad u^{90} - u^{89} + \dots + 42u - 23 \rangle \\
 I_2^u &= \langle u^{19} - u^{18} + \dots + b - u, -u^{19} + u^{18} + \dots + a - u, u^{20} - 2u^{19} + \dots - 7u^2 + 1 \rangle \\
 I_3^u &= \langle 2b - a - 1, a^2 + 3, u + 1 \rangle
 \end{aligned}$$

* 3 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 112 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/math/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$\text{I. } I_1^u = \langle 7.93 \times 10^{127} u^{89} + 3.07 \times 10^{128} u^{88} + \dots + 5.59 \times 10^{128} b + 1.36 \times 10^{130}, -2.38 \times 10^{128} u^{89} + 6.32 \times 10^{129} u^{88} + \dots + 6.43 \times 10^{129} a + 1.54 \times 10^{131}, u^{90} - u^{89} + \dots + 42u - 23 \rangle$$

(i) **Arc colorings**

$$\begin{aligned} a_5 &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_{10} &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_6 &= \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix} \\ a_{12} &= \begin{pmatrix} -u^2 + 1 \\ -u^4 + 2u^2 \end{pmatrix} \\ a_2 &= \begin{pmatrix} 0.0370486u^{89} - 0.983555u^{88} + \dots + 46.7107u - 23.9724 \\ -0.141766u^{89} - 0.549825u^{88} + \dots + 26.6619u - 24.3001 \end{pmatrix} \\ a_4 &= \begin{pmatrix} u \\ u \end{pmatrix} \\ a_1 &= \begin{pmatrix} 0.154839u^{89} - 0.652648u^{88} + \dots + 31.8915u - 18.1094 \\ -0.0239759u^{89} - 0.218917u^{88} + \dots + 11.8427u - 18.4371 \end{pmatrix} \\ a_7 &= \begin{pmatrix} -0.246377u^{89} - 0.228324u^{88} + \dots + 12.1157u + 8.38131 \\ -0.916252u^{89} + 0.319252u^{88} + \dots - 15.5123u - 10.8053 \end{pmatrix} \\ a_3 &= \begin{pmatrix} 0.501815u^{89} + 0.758825u^{88} + \dots - 42.7975u + 32.0231 \\ -0.197294u^{89} + 1.30218u^{88} + \dots - 76.8115u + 9.02154 \end{pmatrix} \\ a_9 &= \begin{pmatrix} 0.0157211u^{89} - 0.164426u^{88} + \dots + 9.47923u - 22.2811 \\ 0.281354u^{89} - 0.146535u^{88} + \dots + 7.20437u + 20.3485 \end{pmatrix} \\ a_8 &= \begin{pmatrix} 0.0160725u^{89} - 0.401426u^{88} + \dots + 17.9232u - 9.14048 \\ -0.0306325u^{89} - 0.222591u^{88} + \dots + 6.58810u + 23.5750 \end{pmatrix} \end{aligned}$$

(ii) **Obstruction class** = -1

(iii) **Cusp Shapes** = $-1.20940u^{89} + 1.41711u^{88} + \dots - 91.7193u + 35.6522$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{90} - 5u^{89} + \cdots - 37888032u + 5971091$
c_2, c_9	$u^{90} + u^{89} + \cdots - 37544u - 17471$
c_3, c_7	$u^{90} - 3u^{89} + \cdots + 642u - 241$
c_4, c_5, c_{10} c_{11}	$u^{90} + u^{89} + \cdots - 42u - 23$
c_6	$u^{90} - u^{89} + \cdots - 383961u - 84943$
c_8, c_{12}	$u^{90} + 3u^{89} + \cdots - 4u - 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{90} - 49y^{89} + \cdots - 1076958384213138y + 35653927730281$
c_2, c_9	$y^{90} + 79y^{89} + \cdots + 6461797462y + 305235841$
c_3, c_7	$y^{90} + 59y^{89} + \cdots + 1230010y + 58081$
c_4, c_5, c_{10} c_{11}	$y^{90} - 113y^{89} + \cdots - 12666y + 529$
c_6	$y^{90} - 23y^{89} + \cdots - 105154505343y + 7215313249$
c_8, c_{12}	$y^{90} + 45y^{89} + \cdots + 76y + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.012970 + 0.095264I$		
$a = -0.077161 - 1.261250I$	$-5.25522 - 3.65690I$	0
$b = 0.71779 - 1.34685I$		
$u = 1.012970 - 0.095264I$		
$a = -0.077161 + 1.261250I$	$-5.25522 + 3.65690I$	0
$b = 0.71779 + 1.34685I$		
$u = 0.857581 + 0.461252I$		
$a = -1.45181 - 1.18703I$	$-8.90905 - 6.49876I$	0
$b = 0.328398 - 1.371680I$		
$u = 0.857581 - 0.461252I$		
$a = -1.45181 + 1.18703I$	$-8.90905 + 6.49876I$	0
$b = 0.328398 + 1.371680I$		
$u = -0.949097 + 0.212026I$		
$a = -0.61914 + 1.31760I$	$-3.71502 + 2.89112I$	0
$b = 0.478508 + 0.868615I$		
$u = -0.949097 - 0.212026I$		
$a = -0.61914 - 1.31760I$	$-3.71502 - 2.89112I$	0
$b = 0.478508 - 0.868615I$		
$u = -0.758287 + 0.530225I$		
$a = 1.031630 - 0.931605I$	$-8.33159 + 1.02539I$	0
$b = -0.00618 - 1.50531I$		
$u = -0.758287 - 0.530225I$		
$a = 1.031630 + 0.931605I$	$-8.33159 - 1.02539I$	0
$b = -0.00618 + 1.50531I$		
$u = -0.905905 + 0.175822I$		
$a = -0.552584 + 1.000800I$	$-3.65291 + 2.91212I$	0
$b = 0.560162 + 0.490336I$		
$u = -0.905905 - 0.175822I$		
$a = -0.552584 - 1.000800I$	$-3.65291 - 2.91212I$	0
$b = 0.560162 - 0.490336I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.909719 + 0.579164I$		
$a = 0.94929 - 1.26410I$	$-6.3264 + 13.2229I$	0
$b = -0.50964 - 1.44518I$		
$u = -0.909719 - 0.579164I$		
$a = 0.94929 + 1.26410I$	$-6.3264 - 13.2229I$	0
$b = -0.50964 + 1.44518I$		
$u = 0.888563 + 0.619652I$		
$a = 0.83455 + 1.14442I$	$-2.42194 - 6.81135I$	0
$b = -0.316840 + 1.346050I$		
$u = 0.888563 - 0.619652I$		
$a = 0.83455 - 1.14442I$	$-2.42194 + 6.81135I$	0
$b = -0.316840 - 1.346050I$		
$u = 0.115526 + 0.862957I$		
$a = 0.192538 + 0.083063I$	$-0.04214 + 1.86425I$	0
$b = 0.173180 + 1.221970I$		
$u = 0.115526 - 0.862957I$		
$a = 0.192538 - 0.083063I$	$-0.04214 - 1.86425I$	0
$b = 0.173180 - 1.221970I$		
$u = 0.981169 + 0.591185I$		
$a = -0.956676 - 0.747354I$	$-6.57746 + 3.74120I$	0
$b = -0.140122 - 1.272130I$		
$u = 0.981169 - 0.591185I$		
$a = -0.956676 + 0.747354I$	$-6.57746 - 3.74120I$	0
$b = -0.140122 + 1.272130I$		
$u = -1.076100 + 0.414278I$		
$a = -0.89733 + 1.13386I$	$-3.86394 + 2.62701I$	0
$b = 0.062943 + 1.084810I$		
$u = -1.076100 - 0.414278I$		
$a = -0.89733 - 1.13386I$	$-3.86394 - 2.62701I$	0
$b = 0.062943 - 1.084810I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.011701 + 0.834793I$		
$a = -0.123216 + 0.122071I$	$-3.59507 - 8.53511I$	0
$b = 0.360219 - 1.321610I$		
$u = -0.011701 - 0.834793I$		
$a = -0.123216 - 0.122071I$	$-3.59507 + 8.53511I$	0
$b = 0.360219 + 1.321610I$		
$u = 0.749413 + 0.327729I$		
$a = -0.087691 + 0.377673I$	$-1.31373 - 7.06051I$	0
$b = -1.302630 + 0.116841I$		
$u = 0.749413 - 0.327729I$		
$a = -0.087691 - 0.377673I$	$-1.31373 + 7.06051I$	0
$b = -1.302630 - 0.116841I$		
$u = -1.195240 + 0.007025I$		
$a = -0.74496 + 1.26726I$	$-3.53900 + 2.66619I$	0
$b = -0.152260 + 0.624422I$		
$u = -1.195240 - 0.007025I$		
$a = -0.74496 - 1.26726I$	$-3.53900 - 2.66619I$	0
$b = -0.152260 - 0.624422I$		
$u = -0.630929 + 0.422737I$		
$a = 0.1320450 - 0.0192536I$	$2.14970 + 2.95397I$	$0. - 5.77136I$
$b = -0.736208 - 0.094191I$		
$u = -0.630929 - 0.422737I$		
$a = 0.1320450 + 0.0192536I$	$2.14970 - 2.95397I$	$0. + 5.77136I$
$b = -0.736208 + 0.094191I$		
$u = -0.683833 + 0.159729I$		
$a = -0.30569 + 3.42355I$	$-1.89496 + 5.35806I$	$-9.62311 - 9.49203I$
$b = 0.306313 + 0.952007I$		
$u = -0.683833 - 0.159729I$		
$a = -0.30569 - 3.42355I$	$-1.89496 - 5.35806I$	$-9.62311 + 9.49203I$
$b = 0.306313 - 0.952007I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.438158 + 0.548624I$		
$a = 0.972243 + 0.724630I$	$-0.18298 + 1.88594I$	$-8.05875 - 4.24644I$
$b = -0.148755 - 1.072620I$		
$u = -0.438158 - 0.548624I$		
$a = 0.972243 - 0.724630I$	$-0.18298 - 1.88594I$	$-8.05875 + 4.24644I$
$b = -0.148755 + 1.072620I$		
$u = 0.540460 + 0.448177I$		
$a = 0.541123 - 0.872191I$	$-1.06587 - 2.16161I$	$-7.84161 + 4.07570I$
$b = 0.484861 + 0.598278I$		
$u = 0.540460 - 0.448177I$		
$a = 0.541123 + 0.872191I$	$-1.06587 + 2.16161I$	$-7.84161 - 4.07570I$
$b = 0.484861 - 0.598278I$		
$u = 0.691662$		
$a = -0.137190$	-1.21324	-7.56480
$b = 0.501893$		
$u = -0.674141 + 0.102515I$		
$a = 1.00488 - 1.94243I$	$-7.28568 + 0.43053I$	$-12.84317 + 4.45391I$
$b = -0.25194 - 1.83276I$		
$u = -0.674141 - 0.102515I$		
$a = 1.00488 + 1.94243I$	$-7.28568 - 0.43053I$	$-12.84317 - 4.45391I$
$b = -0.25194 + 1.83276I$		
$u = 0.673258 + 0.060594I$		
$a = 1.05103 - 2.74686I$	$-0.669088 - 1.009510I$	$-8.02815 + 0.51628I$
$b = 0.047191 - 1.063590I$		
$u = 0.673258 - 0.060594I$		
$a = 1.05103 + 2.74686I$	$-0.669088 + 1.009510I$	$-8.02815 - 0.51628I$
$b = 0.047191 + 1.063590I$		
$u = -0.033569 + 0.644884I$		
$a = 0.604100 + 0.465382I$	$-6.24678 + 2.78122I$	$-9.13407 - 2.15165I$
$b = -0.125543 - 1.364100I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.033569 - 0.644884I$		
$a = 0.604100 - 0.465382I$	$-6.24678 - 2.78122I$	$-9.13407 + 2.15165I$
$b = -0.125543 + 1.364100I$		
$u = -0.303612 + 0.561553I$		
$a = -0.118298 + 1.016180I$	$3.15527 + 0.45173I$	$2.37144 - 2.33678I$
$b = 0.433202 + 0.176963I$		
$u = -0.303612 - 0.561553I$		
$a = -0.118298 - 1.016180I$	$3.15527 - 0.45173I$	$2.37144 + 2.33678I$
$b = 0.433202 - 0.176963I$		
$u = 1.40345 + 0.22038I$		
$a = -0.202364 + 0.963975I$	$-2.28469 - 3.22073I$	0
$b = -0.093332 + 0.518362I$		
$u = 1.40345 - 0.22038I$		
$a = -0.202364 - 0.963975I$	$-2.28469 + 3.22073I$	0
$b = -0.093332 - 0.518362I$		
$u = 1.50828 + 0.07397I$		
$a = -0.449420 - 0.256556I$	$-6.34987 - 3.87000I$	0
$b = 0.443654 - 0.858953I$		
$u = 1.50828 - 0.07397I$		
$a = -0.449420 + 0.256556I$	$-6.34987 + 3.87000I$	0
$b = 0.443654 + 0.858953I$		
$u = 0.144300 + 0.451946I$		
$a = -0.51701 - 1.95209I$	$0.48887 + 4.31858I$	$-0.92438 - 2.28385I$
$b = 0.804357 - 0.071594I$		
$u = 0.144300 - 0.451946I$		
$a = -0.51701 + 1.95209I$	$0.48887 - 4.31858I$	$-0.92438 + 2.28385I$
$b = 0.804357 + 0.071594I$		
$u = 0.222649 + 0.417701I$		
$a = 1.208810 - 0.224028I$	$-0.305635 - 0.881854I$	$-4.92602 + 3.47601I$
$b = -0.549524 + 0.619458I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.222649 - 0.417701I$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	
$a = 1.208810 + 0.224028I$	$-0.305635 + 0.881854I$	$-4.92602 - 3.47601I$
$b = -0.549524 - 0.619458I$		
$u = -1.53355 + 0.11529I$		
$a = -0.612137 - 0.121716I$	$-7.97086 + 4.15764I$	0
$b = -0.316181 + 0.577678I$		
$u = -1.53355 - 0.11529I$		
$a = -0.612137 + 0.121716I$	$-7.97086 - 4.15764I$	0
$b = -0.316181 - 0.577678I$		
$u = 0.455234 + 0.019153I$		
$a = 1.86166 + 0.55835I$	$0.087702 - 0.656020I$	$-8.74359 - 0.87707I$
$b = -0.503361 + 0.850806I$		
$u = 0.455234 - 0.019153I$		
$a = 1.86166 - 0.55835I$	$0.087702 + 0.656020I$	$-8.74359 + 0.87707I$
$b = -0.503361 - 0.850806I$		
$u = -1.58900 + 0.02697I$		
$a = -0.263885 + 1.064150I$	$-7.18439 + 0.98071I$	0
$b = 0.805040 + 0.921807I$		
$u = -1.58900 - 0.02697I$		
$a = -0.263885 - 1.064150I$	$-7.18439 - 0.98071I$	0
$b = 0.805040 - 0.921807I$		
$u = 1.59941 + 0.09338I$		
$a = 0.354560 - 0.276286I$	$-5.48344 - 4.72712I$	0
$b = 1.000530 - 0.274393I$		
$u = 1.59941 - 0.09338I$		
$a = 0.354560 + 0.276286I$	$-5.48344 + 4.72712I$	0
$b = 1.000530 + 0.274393I$		
$u = 0.186645 + 0.338776I$		
$a = 1.176650 - 0.184939I$	$-0.293445 - 0.966895I$	$-5.72850 + 6.30392I$
$b = -0.277893 + 0.577578I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.186645 - 0.338776I$	$-0.293445 + 0.966895I$	$-5.72850 - 6.30392I$
$a = 1.176650 + 0.184939I$		
$b = -0.277893 - 0.577578I$		
$u = -0.287943 + 0.234517I$	$-0.80294 - 3.88261I$	$-4.29656 - 1.29503I$
$a = 1.82271 - 0.65644I$		
$b = -0.624740 + 0.858555I$		
$u = -0.287943 - 0.234517I$	$-0.80294 + 3.88261I$	$-4.29656 + 1.29503I$
$a = 1.82271 + 0.65644I$		
$b = -0.624740 - 0.858555I$		
$u = -1.63002 + 0.01524I$	$-8.78831 + 1.28372I$	0
$a = -0.37399 - 2.31738I$		
$b = 0.119340 - 1.308960I$		
$u = -1.63002 - 0.01524I$	$-8.78831 - 1.28372I$	0
$a = -0.37399 + 2.31738I$		
$b = 0.119340 + 1.308960I$		
$u = 1.63236 + 0.03832I$	$-10.04520 - 6.06116I$	0
$a = -0.09253 + 2.59020I$		
$b = -0.181723 + 1.182940I$		
$u = 1.63236 - 0.03832I$	$-10.04520 + 6.06116I$	0
$a = -0.09253 - 2.59020I$		
$b = -0.181723 - 1.182940I$		
$u = -1.63618$	-9.43889	0
$a = -0.235304$		
$b = -0.875768$		
$u = 1.63978 + 0.01940I$	$-15.5099 - 0.8376I$	0
$a = -0.01052 - 2.33295I$		
$b = 0.53247 - 1.99550I$		
$u = 1.63978 - 0.01940I$	$-15.5099 + 0.8376I$	0
$a = -0.01052 + 2.33295I$		
$b = 0.53247 + 1.99550I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.64318 + 0.08271I$ $a = 0.970520 + 0.369129I$ $b = 1.64897 + 0.16956I$	$-9.64971 + 8.56439I$	0
$u = -1.64318 - 0.08271I$ $a = 0.970520 - 0.369129I$ $b = 1.64897 - 0.16956I$	$-9.64971 - 8.56439I$	0
$u = 1.64607 + 0.17474I$ $a = -0.63968 - 1.88241I$ $b = 0.09702 - 1.62425I$	$-16.5366 - 3.8070I$	0
$u = 1.64607 - 0.17474I$ $a = -0.63968 + 1.88241I$ $b = 0.09702 + 1.62425I$	$-16.5366 + 3.8070I$	0
$u = 1.66541 + 0.03233I$ $a = -0.060097 + 0.537563I$ $b = -0.879840 + 0.217747I$	$-12.60920 - 3.59916I$	0
$u = 1.66541 - 0.03233I$ $a = -0.060097 - 0.537563I$ $b = -0.879840 - 0.217747I$	$-12.60920 + 3.59916I$	0
$u = -1.66382 + 0.13223I$ $a = 0.72905 - 1.80069I$ $b = -0.48612 - 1.42287I$	$-17.6040 + 8.8049I$	0
$u = -1.66382 - 0.13223I$ $a = 0.72905 + 1.80069I$ $b = -0.48612 + 1.42287I$	$-17.6040 - 8.8049I$	0
$u = -1.68038 + 0.17715I$ $a = -0.46078 + 1.91226I$ $b = 0.41651 + 1.49721I$	$-11.2168 + 9.9098I$	0
$u = -1.68038 - 0.17715I$ $a = -0.46078 - 1.91226I$ $b = 0.41651 - 1.49721I$	$-11.2168 - 9.9098I$	0

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.68472 + 0.16892I$		
$a = -0.42408 - 1.97664I$	$-15.2323 - 16.1682I$	0
$b = 0.61190 - 1.57435I$		
$u = 1.68472 - 0.16892I$		
$a = -0.42408 + 1.97664I$	$-15.2323 + 16.1682I$	0
$b = 0.61190 + 1.57435I$		
$u = 1.69447 + 0.10434I$		
$a = 0.47504 + 1.73465I$	$-13.37100 - 4.57563I$	0
$b = -0.396008 + 1.266770I$		
$u = 1.69447 - 0.10434I$		
$a = 0.47504 - 1.73465I$	$-13.37100 + 4.57563I$	0
$b = -0.396008 - 1.266770I$		
$u = 1.70372 + 0.05355I$		
$a = 0.01280 + 1.54850I$	$-13.13290 - 3.93184I$	0
$b = -0.721684 + 1.064470I$		
$u = 1.70372 - 0.05355I$		
$a = 0.01280 - 1.54850I$	$-13.13290 + 3.93184I$	0
$b = -0.721684 - 1.064470I$		
$u = -1.70643 + 0.03143I$		
$a = -0.37326 - 1.89806I$	$-14.8726 + 4.2127I$	0
$b = -0.89338 - 1.67140I$		
$u = -1.70643 - 0.03143I$		
$a = -0.37326 + 1.89806I$	$-14.8726 - 4.2127I$	0
$b = -0.89338 + 1.67140I$		
$u = -1.72856 + 0.14953I$		
$a = 0.56663 - 1.47750I$	$-16.0616 - 0.7646I$	0
$b = -0.131725 - 1.304230I$		
$u = -1.72856 - 0.14953I$		
$a = 0.56663 + 1.47750I$	$-16.0616 + 0.7646I$	0
$b = -0.131725 + 1.304230I$		

$$I_2^u = \langle u^{19} - u^{18} + \cdots + b - u, \quad -u^{19} + u^{18} + \cdots + a - u, \quad u^{20} - 2u^{19} + \cdots - 7u^2 + 1 \rangle$$

(i) **Arc colorings**

$$\begin{aligned} a_5 &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_{10} &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_6 &= \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix} \\ a_{12} &= \begin{pmatrix} -u^2 + 1 \\ -u^4 + 2u^2 \end{pmatrix} \\ a_2 &= \begin{pmatrix} u^{19} - u^{18} + \cdots - 7u^2 + u \\ -u^{19} + u^{18} + \cdots - 3u^2 + u \end{pmatrix} \\ a_4 &= \begin{pmatrix} u \\ u \end{pmatrix} \\ a_1 &= \begin{pmatrix} -2u^{19} + u^{18} + \cdots + 3u + 2 \\ -4u^{19} + 3u^{18} + \cdots + 3u + 2 \end{pmatrix} \\ a_7 &= \begin{pmatrix} -2u^{19} + 2u^{18} + \cdots + 3u + 1 \\ -2u^{19} + u^{18} + \cdots + 2u + 3 \end{pmatrix} \\ a_3 &= \begin{pmatrix} -u^{16} + u^{15} + \cdots + 6u - 1 \\ u^{19} - 2u^{18} + \cdots - u - 2 \end{pmatrix} \\ a_9 &= \begin{pmatrix} 2u^{19} - 3u^{18} + \cdots - 7u - 1 \\ -u^{18} + 13u^{16} + \cdots - 3u - 1 \end{pmatrix} \\ a_8 &= \begin{pmatrix} -2u^{18} + 25u^{16} + \cdots - 5u + 1 \\ -u^{19} - u^{18} + \cdots - 3u^2 - 2u \end{pmatrix} \end{aligned}$$

(ii) **Obstruction class = 1**

(iii) **Cusp Shapes**

$$\begin{aligned} &= 14u^{19} - 8u^{18} - 182u^{17} + 111u^{16} + 973u^{15} - 643u^{14} - 2726u^{13} + 1996u^{12} + 4164u^{11} - \\ &\quad 3523u^{10} - 3100u^9 + 3390u^8 + 456u^7 - 1408u^6 + 581u^5 - 52u^4 - 136u^3 + 102u^2 + u - 21 \end{aligned}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{20} - 7u^{19} + \cdots - 7u + 1$
c_2	$u^{20} - u^{19} + \cdots + u + 1$
c_3	$u^{20} - u^{19} + \cdots + u + 1$
c_4, c_5	$u^{20} + 2u^{19} + \cdots - 7u^2 + 1$
c_6	$u^{20} + 2u^{19} + \cdots + u + 1$
c_7	$u^{20} + u^{19} + \cdots - u + 1$
c_8	$u^{20} + u^{19} + \cdots - u + 1$
c_9	$u^{20} + u^{19} + \cdots - u + 1$
c_{10}, c_{11}	$u^{20} - 2u^{19} + \cdots - 7u^2 + 1$
c_{12}	$u^{20} - u^{19} + \cdots + u + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{20} - 3y^{19} + \cdots - 5y + 1$
c_2, c_9	$y^{20} + 21y^{19} + \cdots + 15y + 1$
c_3, c_7	$y^{20} + 13y^{19} + \cdots + 11y + 1$
c_4, c_5, c_{10} c_{11}	$y^{20} - 28y^{19} + \cdots - 14y + 1$
c_6	$y^{20} + 2y^{19} + \cdots + 9y + 1$
c_8, c_{12}	$y^{20} + 15y^{19} + \cdots + 21y + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.236780 + 0.257246I$		
$a = -0.160920 + 1.140040I$	$-2.96433 + 4.09234I$	$-8.21729 - 6.76865I$
$b = 0.304832 + 0.961803I$		
$u = -1.236780 - 0.257246I$		
$a = -0.160920 - 1.140040I$	$-2.96433 - 4.09234I$	$-8.21729 + 6.76865I$
$b = 0.304832 - 0.961803I$		
$u = 0.676350 + 0.266205I$		
$a = -0.97121 - 1.61423I$	$-7.09778 - 0.93539I$	$-6.18058 + 8.72308I$
$b = 0.16047 - 1.77044I$		
$u = 0.676350 - 0.266205I$		
$a = -0.97121 + 1.61423I$	$-7.09778 + 0.93539I$	$-6.18058 - 8.72308I$
$b = 0.16047 + 1.77044I$		
$u = 1.366380 + 0.211467I$		
$a = 0.58002 + 1.30951I$	$-3.26524 - 1.54191I$	$-7.52291 - 2.90402I$
$b = 0.344378 + 1.031880I$		
$u = 1.366380 - 0.211467I$		
$a = 0.58002 - 1.30951I$	$-3.26524 + 1.54191I$	$-7.52291 + 2.90402I$
$b = 0.344378 - 1.031880I$		
$u = 0.467911 + 0.387225I$		
$a = 1.13017 - 1.14976I$	$0.91339 - 1.46665I$	$-0.60659 + 3.36448I$
$b = -0.147056 + 0.694920I$		
$u = 0.467911 - 0.387225I$		
$a = 1.13017 + 1.14976I$	$0.91339 + 1.46665I$	$-0.60659 - 3.36448I$
$b = -0.147056 - 0.694920I$		
$u = 0.040099 + 0.531181I$		
$a = -0.496026 - 0.240428I$	$1.15814 - 0.98852I$	$-0.555070 + 1.124089I$
$b = -0.253507 + 0.945631I$		
$u = 0.040099 - 0.531181I$		
$a = -0.496026 + 0.240428I$	$1.15814 + 0.98852I$	$-0.555070 - 1.124089I$
$b = -0.253507 - 0.945631I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.53052 + 0.04941I$	$-7.53461 - 5.30221I$	$-9.13332 + 7.29862I$
$a = 0.340914 + 0.433719I$		
$b = 0.436448 - 0.482124I$		
$u = 1.53052 - 0.04941I$	$-7.53461 + 5.30221I$	$-9.13332 - 7.29862I$
$a = 0.340914 - 0.433719I$		
$b = 0.436448 + 0.482124I$		
$u = -1.55472 + 0.09338I$	$-6.02214 + 3.05677I$	$-5.71579 + 1.69066I$
$a = -0.710342 - 0.085200I$		
$b = 0.248590 + 0.542254I$		
$u = -1.55472 - 0.09338I$	$-6.02214 - 3.05677I$	$-5.71579 - 1.69066I$
$a = -0.710342 + 0.085200I$		
$b = 0.248590 - 0.542254I$		
$u = -0.336288 + 0.122318I$	$-1.00182 + 4.64059I$	$-7.03732 - 7.06030I$
$a = -1.74662 + 2.45740I$		
$b = -0.475237 - 0.606200I$		
$u = -0.336288 - 0.122318I$	$-1.00182 - 4.64059I$	$-7.03732 + 7.06030I$
$a = -1.74662 - 2.45740I$		
$b = -0.475237 + 0.606200I$		
$u = -1.66475 + 0.07750I$	$-15.4766 + 2.2859I$	$-11.50746 - 1.72176I$
$a = 0.09942 - 2.10149I$		
$b = -0.43525 - 1.80930I$		
$u = -1.66475 - 0.07750I$	$-15.4766 - 2.2859I$	$-11.50746 + 1.72176I$
$a = 0.09942 + 2.10149I$		
$b = -0.43525 + 1.80930I$		
$u = 1.71129 + 0.03601I$	$-12.99190 - 4.52813I$	$-8.52367 + 9.83951I$
$a = -0.06541 + 1.69236I$		
$b = -0.683669 + 1.135430I$		
$u = 1.71129 - 0.03601I$	$-12.99190 + 4.52813I$	$-8.52367 - 9.83951I$
$a = -0.06541 - 1.69236I$		
$b = -0.683669 - 1.135430I$		

$$\text{III. } I_3^u = \langle 2b - a - 1, a^2 + 3, u + 1 \rangle$$

(i) Arc colorings

$$a_5 = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} a \\ \frac{1}{2}a + \frac{1}{2} \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -1 \\ -1 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} \frac{1}{2}a + \frac{1}{2} \\ 1 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} \frac{1}{2}a + \frac{3}{2} \\ 1 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} a - 1 \\ \frac{1}{2}a - \frac{1}{2} \end{pmatrix}$$

$$a_9 = \begin{pmatrix} \frac{1}{2}a - \frac{1}{2} \\ \frac{1}{2}a - \frac{1}{2} \end{pmatrix}$$

$$a_8 = \begin{pmatrix} \frac{1}{2}a - \frac{1}{2} \\ 0 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = $-4a - 9$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_3, c_9 c_{12}	$u^2 - u + 1$
c_2, c_7, c_8	$u^2 + u + 1$
c_4, c_5, c_6	$(u - 1)^2$
c_{10}, c_{11}	$(u + 1)^2$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_3 c_7, c_8, c_9 c_{12}	$y^2 + y + 1$
c_4, c_5, c_6 c_{10}, c_{11}	$(y - 1)^2$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.00000$		
$a = 1.73205I$	$-3.28987 + 4.05977I$	$-9.00000 - 6.92820I$
$b = 0.500000 + 0.866025I$		
$u = -1.00000$		
$a = -1.73205I$	$-3.28987 - 4.05977I$	$-9.00000 + 6.92820I$
$b = 0.500000 - 0.866025I$		

IV. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1	$(u^2 - u + 1)(u^{20} - 7u^{19} + \dots - 7u + 1)$ $\cdot (u^{90} - 5u^{89} + \dots - 37888032u + 5971091)$
c_2	$(u^2 + u + 1)(u^{20} - u^{19} + \dots + u + 1)(u^{90} + u^{89} + \dots - 37544u - 17471)$
c_3	$(u^2 - u + 1)(u^{20} - u^{19} + \dots + u + 1)(u^{90} - 3u^{89} + \dots + 642u - 241)$
c_4, c_5	$((u - 1)^2)(u^{20} + 2u^{19} + \dots - 7u^2 + 1)(u^{90} + u^{89} + \dots - 42u - 23)$
c_6	$((u - 1)^2)(u^{20} + 2u^{19} + \dots + u + 1)(u^{90} - u^{89} + \dots - 383961u - 84943)$
c_7	$(u^2 + u + 1)(u^{20} + u^{19} + \dots - u + 1)(u^{90} - 3u^{89} + \dots + 642u - 241)$
c_8	$(u^2 + u + 1)(u^{20} + u^{19} + \dots - u + 1)(u^{90} + 3u^{89} + \dots - 4u - 1)$
c_9	$(u^2 - u + 1)(u^{20} + u^{19} + \dots - u + 1)(u^{90} + u^{89} + \dots - 37544u - 17471)$
c_{10}, c_{11}	$((u + 1)^2)(u^{20} - 2u^{19} + \dots - 7u^2 + 1)(u^{90} + u^{89} + \dots - 42u - 23)$
c_{12}	$(u^2 - u + 1)(u^{20} - u^{19} + \dots + u + 1)(u^{90} + 3u^{89} + \dots - 4u - 1)$

V. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1	$(y^2 + y + 1)(y^{20} - 3y^{19} + \cdots - 5y + 1)$ $\cdot (y^{90} - 49y^{89} + \cdots - 1076958384213138y + 35653927730281)$
c_2, c_9	$(y^2 + y + 1)(y^{20} + 21y^{19} + \cdots + 15y + 1)$ $\cdot (y^{90} + 79y^{89} + \cdots + 6461797462y + 305235841)$
c_3, c_7	$(y^2 + y + 1)(y^{20} + 13y^{19} + \cdots + 11y + 1)$ $\cdot (y^{90} + 59y^{89} + \cdots + 1230010y + 58081)$
c_4, c_5, c_{10} c_{11}	$((y - 1)^2)(y^{20} - 28y^{19} + \cdots - 14y + 1)$ $\cdot (y^{90} - 113y^{89} + \cdots - 12666y + 529)$
c_6	$((y - 1)^2)(y^{20} + 2y^{19} + \cdots + 9y + 1)$ $\cdot (y^{90} - 23y^{89} + \cdots - 105154505343y + 7215313249)$
c_8, c_{12}	$(y^2 + y + 1)(y^{20} + 15y^{19} + \cdots + 21y + 1)(y^{90} + 45y^{89} + \cdots + 76y + 1)$