



12n₀₂₆₆ (K12n₀₂₆₆)



A knot diagram¹

Linearized knot diagram



Solving Sequence

$$3,8 \xrightarrow{c_8} 9,12 \xrightarrow{c_7} 7 \xrightarrow{c_9} 10 \xrightarrow{c_{12}} 1 \xrightarrow{c_6} 6 \xrightarrow{c_{11}} 11 \xrightarrow{c_5} 5 \xrightarrow{c_2} 2 \xrightarrow{c_4} 4 \twoheadrightarrow c_1, c_3, c_{10}$$

Ideals for irreducible components² of X_{par}

$$I_1^u = \langle 1.30470 \times 10^{57} u^{29} - 8.04261 \times 10^{57} u^{28} + \dots + 3.78260 \times 10^{57} b - 1.44716 \times 10^{60}, \\ 1.67303 \times 10^{56} u^{29} - 2.36382 \times 10^{57} u^{28} + \dots + 3.02608 \times 10^{58} a - 1.98438 \times 10^{60}, \\ u^{30} - 7u^{29} + \dots - 8960u + 1024 \rangle$$

$$I_2^u = \langle 85a^5 u^3 + 81a^4 u^3 + \dots - 914a - 116, 4a^5 u^3 + 36a^4 u^3 + \dots + 70a - 27, u^4 + 3u^3 + 3u^2 + 2u + 2 \rangle$$

$$I_3^u = \langle 4147258u^{15} + 7664206u^{14} + \dots + 53503381b + 7608876, \\ - 151936201u^{15} + 44795175u^{14} + \dots + 53503381a - 367339747, \\ u^{16} - 3u^{14} - 4u^{12} + 5u^{11} + 23u^{10} - 7u^9 - 13u^8 + 6u^7 - 11u^6 - 12u^5 + 13u^4 - 4u^3 + 5u^2 + u + 1 \rangle$$

$$I_4^u = \langle 9.10023 \times 10^{19} a^{11} u + 1.90099 \times 10^{20} a^{10} u + \dots + 3.63923 \times 10^{20} a + 7.09845 \times 10^{19}, \\ - a^{11} u - 29a^{10} u + \dots - 3954a + 1387, u^2 - u - 1 \rangle$$

$$I_1^v = \langle a, 8v^3 + 12v^2 + b + 10v + 3, 8v^4 + 12v^3 + 12v^2 + 5v + 1 \rangle$$

$$I_2^v = \langle a, b^6 + b^5 + 2b^4 + 2b^3 + 2b^2 + 2b + 1, v - 1 \rangle$$

* 6 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 104 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/maths/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$\mathbf{I. } I_1^u = \langle 1.30 \times 10^{57} u^{29} - 8.04 \times 10^{57} u^{28} + \dots + 3.78 \times 10^{57} b - 1.45 \times 10^{60}, 1.67 \times 10^{56} u^{29} - 2.36 \times 10^{57} u^{28} + \dots + 3.03 \times 10^{58} a - 1.98 \times 10^{60}, u^{30} - 7u^{29} + \dots - 8960u + 1024 \rangle$$

(i) Arc colorings

$$a_3 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ -u^2 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -0.00552871u^{29} + 0.0781148u^{28} + \dots - 455.718u + 65.5759 \\ -0.344921u^{29} + 2.12621u^{28} + \dots - 3000.66u + 382.582 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -0.0965905u^{29} + 0.626567u^{28} + \dots - 1195.73u + 163.169 \\ 0.437196u^{29} - 2.77353u^{28} + \dots + 5006.91u - 693.942 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -0.199941u^{29} + 1.25377u^{28} + \dots - 2078.67u + 283.434 \\ 0.126079u^{29} - 0.746114u^{28} + \dots + 510.964u - 30.8012 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -0.237653u^{29} + 1.50770u^{28} + \dots - 2693.25u + 371.955 \\ -0.0793898u^{29} + 0.480636u^{28} + \dots - 898.423u + 142.638 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -0.393468u^{29} + 2.51203u^{28} + \dots - 4452.78u + 601.479 \\ -0.252406u^{29} + 1.63090u^{28} + \dots - 2955.26u + 391.693 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -0.350450u^{29} + 2.20433u^{28} + \dots - 3456.38u + 448.158 \\ -0.344921u^{29} + 2.12621u^{28} + \dots - 3000.66u + 382.582 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -0.0528628u^{29} + 0.365076u^{28} + \dots - 641.595u + 69.7063 \\ 0.184790u^{29} - 1.14262u^{28} + \dots + 2051.66u - 302.249 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -0.237653u^{29} + 1.50770u^{28} + \dots - 2693.25u + 371.955 \\ -0.184790u^{29} + 1.14262u^{28} + \dots - 2051.66u + 302.249 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -u \\ -u \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes = $-0.780219u^{29} + 4.90701u^{28} + \dots - 10318.3u + 1584.28$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{30} + 15u^{29} + \dots + 5376u + 4096$
c_2, c_4	$u^{30} - 5u^{29} + \dots - 144u + 64$
c_3, c_8	$u^{30} + 7u^{29} + \dots + 8960u + 1024$
c_5, c_7, c_{10} c_{11}	$u^{30} + 16u^{28} + \dots - 4u + 1$
c_6, c_9	$u^{30} + u^{29} + \dots - 3u + 1$
c_{12}	$u^{30} + 25u^{29} + \dots + 4224u + 256$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{30} + 5y^{29} + \dots + 918224896y + 16777216$
c_2, c_4	$y^{30} - 15y^{29} + \dots - 5376y + 4096$
c_3, c_8	$y^{30} - 15y^{29} + \dots - 7667712y + 1048576$
c_5, c_7, c_{10} c_{11}	$y^{30} + 32y^{29} + \dots - 14y + 1$
c_6, c_9	$y^{30} - 13y^{29} + \dots + 5y + 1$
c_{12}	$y^{30} + 7y^{29} + \dots + 180224y + 65536$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.295304 + 0.827577I$		
$a = 0.432685 + 0.180516I$	$-1.52614 - 1.54845I$	$-7.05353 + 2.25516I$
$b = 0.502251 - 0.079193I$		
$u = 0.295304 - 0.827577I$		
$a = 0.432685 - 0.180516I$	$-1.52614 + 1.54845I$	$-7.05353 - 2.25516I$
$b = 0.502251 + 0.079193I$		
$u = -1.118510 + 0.267682I$		
$a = 0.344821 + 0.251878I$	$2.84106 - 1.32814I$	$-1.11844 - 2.01669I$
$b = -0.488387 - 0.036524I$		
$u = -1.118510 - 0.267682I$		
$a = 0.344821 - 0.251878I$	$2.84106 + 1.32814I$	$-1.11844 + 2.01669I$
$b = -0.488387 + 0.036524I$		
$u = 0.790526 + 0.025536I$		
$a = 0.369046 + 0.400347I$	$-1.67774 + 2.21849I$	$-1.35323 - 5.31460I$
$b = 0.758312 - 0.591430I$		
$u = 0.790526 - 0.025536I$		
$a = 0.369046 - 0.400347I$	$-1.67774 - 2.21849I$	$-1.35323 + 5.31460I$
$b = 0.758312 + 0.591430I$		
$u = 1.150710 + 0.487372I$		
$a = 0.146682 - 0.393834I$	$1.25283 + 6.44217I$	$-7.35418 - 5.48188I$
$b = -0.578991 + 0.126474I$		
$u = 1.150710 - 0.487372I$		
$a = 0.146682 + 0.393834I$	$1.25283 - 6.44217I$	$-7.35418 + 5.48188I$
$b = -0.578991 - 0.126474I$		
$u = 0.587875 + 0.294150I$		
$a = 1.49596 - 0.54450I$	$-2.33932 - 0.29501I$	$-4.95113 - 2.59003I$
$b = -0.410397 - 0.248863I$		
$u = 0.587875 - 0.294150I$		
$a = 1.49596 + 0.54450I$	$-2.33932 + 0.29501I$	$-4.95113 + 2.59003I$
$b = -0.410397 + 0.248863I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.281814 + 0.580917I$ $a = 0.633016 - 0.523942I$ $b = 0.265637 + 0.486839I$	$-0.114599 - 1.183600I$	$-1.52357 + 6.11116I$
$u = -0.281814 - 0.580917I$ $a = 0.633016 + 0.523942I$ $b = 0.265637 - 0.486839I$	$-0.114599 + 1.183600I$	$-1.52357 - 6.11116I$
$u = 0.581579 + 0.213573I$ $a = -0.181652 - 0.352155I$ $b = -0.562214 + 1.036190I$	$1.23856 + 8.05442I$	$5.2311 - 14.4328I$
$u = 0.581579 - 0.213573I$ $a = -0.181652 + 0.352155I$ $b = -0.562214 - 1.036190I$	$1.23856 - 8.05442I$	$5.2311 + 14.4328I$
$u = -0.21710 + 1.53848I$ $a = 0.204399 + 0.699894I$ $b = 0.241334 - 0.976861I$	$-5.66421 + 0.94543I$	$-11.8910 - 17.8120I$
$u = -0.21710 - 1.53848I$ $a = 0.204399 - 0.699894I$ $b = 0.241334 + 0.976861I$	$-5.66421 - 0.94543I$	$-11.8910 + 17.8120I$
$u = 0.05187 + 1.70653I$ $a = -0.098466 + 0.394900I$ $b = -0.28890 - 1.53476I$	$7.74868 - 8.57967I$	0
$u = 0.05187 - 1.70653I$ $a = -0.098466 - 0.394900I$ $b = -0.28890 + 1.53476I$	$7.74868 + 8.57967I$	0
$u = 1.69998 + 0.25374I$ $a = -0.298306 - 1.350070I$ $b = 0.15069 + 1.46462I$	$5.39652 - 5.08088I$	0
$u = 1.69998 - 0.25374I$ $a = -0.298306 + 1.350070I$ $b = 0.15069 - 1.46462I$	$5.39652 + 5.08088I$	0

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.61078 + 0.80572I$ $a = 0.68358 + 1.24119I$ $b = 0.61201 - 1.60557I$	$12.5033 + 17.2116I$	0
$u = 1.61078 - 0.80572I$ $a = 0.68358 - 1.24119I$ $b = 0.61201 + 1.60557I$	$12.5033 - 17.2116I$	0
$u = 0.47534 + 1.81581I$ $a = -0.048480 - 0.458166I$ $b = 0.00271 + 1.46604I$	$7.03462 + 1.24559I$	0
$u = 0.47534 - 1.81581I$ $a = -0.048480 + 0.458166I$ $b = 0.00271 - 1.46604I$	$7.03462 - 1.24559I$	0
$u = -1.87027 + 0.50681I$ $a = 0.337792 - 1.250370I$ $b = 0.45821 + 1.67495I$	$14.6658 - 9.7485I$	0
$u = -1.87027 - 0.50681I$ $a = 0.337792 + 1.250370I$ $b = 0.45821 - 1.67495I$	$14.6658 + 9.7485I$	0
$u = 1.69641 + 0.95826I$ $a = -0.604731 - 1.038330I$ $b = -0.42883 + 1.37214I$	$10.87200 + 8.65298I$	0
$u = 1.69641 - 0.95826I$ $a = -0.604731 + 1.038330I$ $b = -0.42883 - 1.37214I$	$10.87200 - 8.65298I$	0
$u = -1.95267 + 0.75510I$ $a = -0.416349 + 1.035550I$ $b = -0.23344 - 1.52553I$	$13.56600 - 0.76045I$	0
$u = -1.95267 - 0.75510I$ $a = -0.416349 - 1.035550I$ $b = -0.23344 + 1.52553I$	$13.56600 + 0.76045I$	0

$$\text{II. } I_2^u = \langle 85a^5u^3 + 81a^4u^3 + \dots - 914a - 116, 4a^5u^3 + 36a^4u^3 + \dots + 70a - 27, u^4 + 3u^3 + 3u^2 + 2u + 2 \rangle$$

(i) Arc colorings

$$a_3 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ -u^2 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} a \\ -0.702479a^5u^3 - 0.669421a^4u^3 + \dots + 7.55372a + 0.958678 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} 0.305785a^5u^3 - 0.876033a^4u^3 + \dots + 1.58678a + 0.314050 \\ 0.834711a^5u^3 + 2.74380a^4u^3 + \dots - 8.47934a - 1.75207 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 0.239669a^4u^3 - 0.909091a^3u^3 + \dots - 0.727273a + 1.32231 \\ 1.05785a^4u^3 + 1.54545a^3u^3 + \dots + 5.63636a + 0.595041 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -\frac{1}{2}u^3 - \frac{1}{2}u^2 - \frac{1}{2}u - 1 \\ u^3 + 2u^2 + u + 1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 0.0826446a^5u^3 + 0.628099a^4u^3 + \dots - 5.76033a - 1.30579 \\ -0.975207a^5u^3 - 0.611570a^4u^3 + \dots - 4.62810a + 1.55372 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -0.702479a^5u^3 - 0.669421a^4u^3 + \dots + 8.55372a + 0.958678 \\ -0.702479a^5u^3 - 0.669421a^4u^3 + \dots + 7.55372a + 0.958678 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} \frac{1}{2}u^3 + \frac{1}{2}u^2 - \frac{1}{2}u \\ u^3 + u^2 + 1 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -\frac{1}{2}u^3 - \frac{1}{2}u^2 - \frac{1}{2}u - 1 \\ -u^3 - u^2 - 1 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -u \\ -u \end{pmatrix}$$

(ii) Obstruction class = -1

$$\text{(iii) Cusp Shapes} = -\frac{148}{121}a^4u^3 + \frac{28}{11}a^3u^3 + \dots - \frac{136}{11}a + \frac{68}{121}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$(u^4 + u^3 + 4u^2 + u + 1)^6$
c_2, c_4	$(u^4 - u^3 + u + 1)^6$
c_3, c_8	$(u^4 - 3u^3 + 3u^2 - 2u + 2)^6$
c_5, c_7, c_{10} c_{11}	$u^{24} + u^{23} + \dots - 6u + 23$
c_6, c_9	$u^{24} + 3u^{23} + \dots + 82u + 67$
c_{12}	$(u^3 - u^2 + 1)^8$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$(y^4 + 7y^3 + 16y^2 + 7y + 1)^6$
c_2, c_4	$(y^4 - y^3 + 4y^2 - y + 1)^6$
c_3, c_8	$(y^4 - 3y^3 + y^2 + 8y + 4)^6$
c_5, c_7, c_{10} c_{11}	$y^{24} + 21y^{23} + \dots + 24252y + 529$
c_6, c_9	$y^{24} - 7y^{23} + \dots - 56036y + 4489$
c_{12}	$(y^3 - y^2 + 2y - 1)^8$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.066121 + 0.864054I$ $a = 0.847489 - 0.409743I$ $b = 0.435029 + 1.039690I$	$1.24998 - 1.37790I$	$-0.06985 - 1.74429I$
$u = 0.066121 + 0.864054I$ $a = -0.066440 + 1.227610I$ $b = 0.219486 - 1.053480I$	$1.24998 + 4.27835I$	$-0.06985 - 7.70319I$
$u = 0.066121 + 0.864054I$ $a = -0.685301 + 1.029190I$ $b = -0.34890 + 1.62773I$	$5.38756 + 1.45022I$	$6.45941 - 4.72374I$
$u = 0.066121 + 0.864054I$ $a = 1.56525 - 0.14261I$ $b = 0.026299 - 1.366270I$	$5.38756 + 1.45022I$	$6.45941 - 4.72374I$
$u = 0.066121 + 0.864054I$ $a = 1.51833 - 1.27411I$ $b = 0.042954 - 0.201002I$	$1.24998 - 1.37790I$	$-0.06985 - 1.74429I$
$u = 0.066121 + 0.864054I$ $a = -1.63512 + 1.12550I$ $b = -0.940992 + 0.412160I$	$1.24998 + 4.27835I$	$-0.06985 - 7.70319I$
$u = 0.066121 - 0.864054I$ $a = 0.847489 + 0.409743I$ $b = 0.435029 - 1.039690I$	$1.24998 + 1.37790I$	$-0.06985 + 1.74429I$
$u = 0.066121 - 0.864054I$ $a = -0.066440 - 1.227610I$ $b = 0.219486 + 1.053480I$	$1.24998 - 4.27835I$	$-0.06985 + 7.70319I$
$u = 0.066121 - 0.864054I$ $a = -0.685301 - 1.029190I$ $b = -0.34890 - 1.62773I$	$5.38756 - 1.45022I$	$6.45941 + 4.72374I$
$u = 0.066121 - 0.864054I$ $a = 1.56525 + 0.14261I$ $b = 0.026299 + 1.366270I$	$5.38756 - 1.45022I$	$6.45941 + 4.72374I$

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.066121 - 0.864054I$ $a = 1.51833 + 1.27411I$ $b = 0.042954 + 0.201002I$	$1.24998 + 1.37790I$	$-0.06985 + 1.74429I$
$u = 0.066121 - 0.864054I$ $a = -1.63512 - 1.12550I$ $b = -0.940992 - 0.412160I$	$1.24998 - 4.27835I$	$-0.06985 + 7.70319I$
$u = -1.56612 + 0.45882I$ $a = 0.446198 - 1.247430I$ $b = 0.95639 + 1.77228I$	$10.82130 - 6.78371I$	$7.57961 + 4.72374I$
$u = -1.56612 + 0.45882I$ $a = 0.352386 - 1.325540I$ $b = -0.05667 + 1.51321I$	$6.68371 - 3.95559I$	$1.05034 + 1.74429I$
$u = -1.56612 + 0.45882I$ $a = -1.04122 + 1.11468I$ $b = -0.348867 - 1.279900I$	$10.82130 - 6.78371I$	$7.57961 + 4.72374I$
$u = -1.56612 + 0.45882I$ $a = -0.123141 - 0.393851I$ $b = 1.61223 + 0.24522I$	$6.68371 - 9.61184I$	$1.05034 + 7.70319I$
$u = -1.56612 + 0.45882I$ $a = -0.272467 - 0.089477I$ $b = -0.843469 + 0.066206I$	$6.68371 - 3.95559I$	$1.05034 + 1.74429I$
$u = -1.56612 + 0.45882I$ $a = -0.40595 + 1.70866I$ $b = -0.25349 - 1.45295I$	$6.68371 - 9.61184I$	$1.05034 + 7.70319I$
$u = -1.56612 - 0.45882I$ $a = 0.446198 + 1.247430I$ $b = 0.95639 - 1.77228I$	$10.82130 + 6.78371I$	$7.57961 - 4.72374I$
$u = -1.56612 - 0.45882I$ $a = 0.352386 + 1.325540I$ $b = -0.05667 - 1.51321I$	$6.68371 + 3.95559I$	$1.05034 - 1.74429I$

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.56612 - 0.45882I$ $a = -1.04122 - 1.11468I$ $b = -0.348867 + 1.279900I$	$10.82130 + 6.78371I$	$7.57961 - 4.72374I$
$u = -1.56612 - 0.45882I$ $a = -0.123141 + 0.393851I$ $b = 1.61223 - 0.24522I$	$6.68371 + 9.61184I$	$1.05034 - 7.70319I$
$u = -1.56612 - 0.45882I$ $a = -0.272467 + 0.089477I$ $b = -0.843469 - 0.066206I$	$6.68371 + 3.95559I$	$1.05034 - 1.74429I$
$u = -1.56612 - 0.45882I$ $a = -0.40595 - 1.70866I$ $b = -0.25349 + 1.45295I$	$6.68371 + 9.61184I$	$1.05034 - 7.70319I$

III.

$$I_3^u = \langle 4.15 \times 10^6 u^{15} + 7.66 \times 10^6 u^{14} + \dots + 5.35 \times 10^7 b + 7.61 \times 10^6, -1.52 \times 10^8 u^{15} + 4.48 \times 10^7 u^{14} + \dots + 5.35 \times 10^7 a - 3.67 \times 10^8, u^{16} - 3u^{14} + \dots + u + 1 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_3 &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_8 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_9 &= \begin{pmatrix} 1 \\ -u^2 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} 2.83975u^{15} - 0.837240u^{14} + \dots + 10.0880u + 6.86573 \\ -0.0775139u^{15} - 0.143247u^{14} + \dots + 1.48032u - 0.142213 \end{pmatrix} \\ a_7 &= \begin{pmatrix} 1.46611u^{15} + 0.555277u^{14} + \dots - 7.60428u + 7.02320 \\ 0.342955u^{15} - 0.0913471u^{14} + \dots + 0.412915u - 0.146361 \end{pmatrix} \\ a_{10} &= \begin{pmatrix} 2.10671u^{15} + 1.02973u^{14} + \dots + 1.90799u + 11.0945 \\ 0.202635u^{15} + 0.190332u^{14} + \dots + 2.50674u + 0.600158 \end{pmatrix} \\ a_1 &= \begin{pmatrix} -0.478951u^{15} + 0.274850u^{14} + \dots - 3.19806u - 1.01076 \\ 0.134911u^{15} + 0.189098u^{14} + \dots + 0.316322u + 0.253088 \end{pmatrix} \\ a_6 &= \begin{pmatrix} 2.43159u^{15} + 0.321717u^{14} + \dots - 3.47288u + 6.86583 \\ 0.486533u^{15} + 0.0412901u^{14} + \dots + 0.933338u - 0.168123 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 2.76224u^{15} - 0.980487u^{14} + \dots + 11.5683u + 6.72352 \\ -0.0775139u^{15} - 0.143247u^{14} + \dots + 1.48032u - 0.142213 \end{pmatrix} \\ a_5 &= \begin{pmatrix} 0.622529u^{15} - 0.142213u^{14} + \dots + 3.71848u + 0.988996 \\ 0.143578u^{15} + 0.132637u^{14} + \dots + 0.520422u - 0.0217622 \end{pmatrix} \\ a_2 &= \begin{pmatrix} -0.478951u^{15} + 0.274850u^{14} + \dots - 3.19806u - 1.01076 \\ 0.143578u^{15} + 0.132637u^{14} + \dots + 0.520422u - 0.0217622 \end{pmatrix} \\ a_4 &= \begin{pmatrix} u \\ u \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = 1

$$(iii) \text{ Cusp Shapes} = \frac{119862154}{53503381}u^{15} - \frac{50233104}{53503381}u^{14} + \dots + \frac{2012450980}{53503381}u - \frac{93776732}{53503381}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{16} - 10u^{15} + \dots - 7u + 1$
c_2	$u^{16} + 6u^{15} + \dots - u + 1$
c_3	$u^{16} - 3u^{14} + \dots - u + 1$
c_4	$u^{16} - 6u^{15} + \dots + u + 1$
c_5, c_{11}	$u^{16} + 8u^{14} + \dots + 3u + 1$
c_6, c_9	$u^{16} - 3u^{15} + \dots - 6u + 1$
c_7, c_{10}	$u^{16} + 8u^{14} + \dots - 3u + 1$
c_8	$u^{16} - 3u^{14} + \dots + u + 1$
c_{12}	$u^{16} - 5u^{15} + \dots - 5u^3 + 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{16} - 2y^{15} + \dots + 61y + 1$
c_2, c_4	$y^{16} - 10y^{15} + \dots - 7y + 1$
c_3, c_8	$y^{16} - 6y^{15} + \dots + 9y + 1$
c_5, c_7, c_{10} c_{11}	$y^{16} + 16y^{15} + \dots + 11y + 1$
c_6, c_9	$y^{16} - 5y^{15} + \dots - 14y + 1$
c_{12}	$y^{16} + 9y^{15} + \dots + 10y^2 + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.978949 + 0.078957I$ $a = -0.453108 - 0.320975I$ $b = 0.445295 - 0.521652I$	$3.29193 + 2.08322I$	$4.28264 - 4.24407I$
$u = 0.978949 - 0.078957I$ $a = -0.453108 + 0.320975I$ $b = 0.445295 + 0.521652I$	$3.29193 - 2.08322I$	$4.28264 + 4.24407I$
$u = -0.996285 + 0.490791I$ $a = -0.353672 - 0.270556I$ $b = 0.279505 + 0.496867I$	$1.92823 - 6.43556I$	$5.05003 + 5.97847I$
$u = -0.996285 - 0.490791I$ $a = -0.353672 + 0.270556I$ $b = 0.279505 - 0.496867I$	$1.92823 + 6.43556I$	$5.05003 - 5.97847I$
$u = 0.196535 + 0.775710I$ $a = -0.14260 + 1.75182I$ $b = 0.08961 + 1.47734I$	$4.79804 - 0.70603I$	$0.45722 - 4.37222I$
$u = 0.196535 - 0.775710I$ $a = -0.14260 - 1.75182I$ $b = 0.08961 - 1.47734I$	$4.79804 + 0.70603I$	$0.45722 + 4.37222I$
$u = 0.172282 + 0.628783I$ $a = 2.18279 - 0.75635I$ $b = 0.405160 + 0.782797I$	$1.56983 - 2.39298I$	$2.54235 + 6.16314I$
$u = 0.172282 - 0.628783I$ $a = 2.18279 + 0.75635I$ $b = 0.405160 - 0.782797I$	$1.56983 + 2.39298I$	$2.54235 - 6.16314I$
$u = -0.206080 + 0.337926I$ $a = 10.45890 + 6.32490I$ $b = -0.380479 + 0.681918I$	$0.32266 + 2.98693I$	$-4.4966 + 17.7278I$
$u = -0.206080 - 0.337926I$ $a = 10.45890 - 6.32490I$ $b = -0.380479 - 0.681918I$	$0.32266 - 2.98693I$	$-4.4966 - 17.7278I$

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.61349 + 0.06231I$ $a = -0.307309 + 1.257690I$ $b = -0.43055 - 1.58967I$	$10.48480 + 0.24679I$	$0.73600 - 1.75528I$
$u = -1.61349 - 0.06231I$ $a = -0.307309 - 1.257690I$ $b = -0.43055 + 1.58967I$	$10.48480 - 0.24679I$	$0.73600 + 1.75528I$
$u = -0.13825 + 1.62727I$ $a = 0.254873 + 0.700959I$ $b = 0.192282 - 1.004160I$	$-5.54863 + 0.76161I$	$8.2609 + 13.1613I$
$u = -0.13825 - 1.62727I$ $a = 0.254873 - 0.700959I$ $b = 0.192282 + 1.004160I$	$-5.54863 - 0.76161I$	$8.2609 - 13.1613I$
$u = 1.60635 + 0.50423I$ $a = -0.639909 - 1.085690I$ $b = -0.60082 + 1.36455I$	$9.47206 + 6.62853I$	$0.16748 - 3.62136I$
$u = 1.60635 - 0.50423I$ $a = -0.639909 + 1.085690I$ $b = -0.60082 - 1.36455I$	$9.47206 - 6.62853I$	$0.16748 + 3.62136I$

$$\text{IV. } I_4^u = \langle 9.10 \times 10^{19} a^{11} u + 1.90 \times 10^{20} a^{10} u + \dots + 3.64 \times 10^{20} a + 7.10 \times 10^{19}, -a^{11} u - 29a^{10} u + \dots - 3954a + 1387, u^2 - u - 1 \rangle$$

(i) Arc colorings

$$a_3 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ -u - 1 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} a \\ -1.56829a^{11}u - 3.27608a^{10}u + \dots - 6.27166a - 1.22331 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -0.460239a^{11}u - 1.23068a^{10}u + \dots - 5.15203a + 1.64031 \\ 2.83175a^{11}u + 6.50769a^{10}u + \dots + 17.3200a + 1.87860 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -0.837001a^{11}u - 1.69786a^{10}u + \dots - 2.77220a - 2.20499 \\ -1.53854a^{11}u - 3.00157a^{10}u + \dots - 9.06731a - 1.48673 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} 1.83049a^{11}u + 3.69414a^{10}u + \dots + 10.1122a + 0.883061 \\ -1.83049a^{11}u - 3.69414a^{10}u + \dots - 10.1122a - 1.88306 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 2.59830a^{11}u + 5.43442a^{10}u + \dots + 10.6594a + 4.31367 \\ -0.401435a^{11}u - 0.287843a^{10}u + \dots - 1.58392a + 2.69916 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -1.56829a^{11}u - 3.27608a^{10}u + \dots - 5.27166a - 1.22331 \\ -1.56829a^{11}u - 3.27608a^{10}u + \dots - 6.27166a - 1.22331 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -1.13131a^{11}u - 2.28310a^{10}u + \dots - 6.25174a - 0.296436 \\ -2.96180a^{11}u - 5.97724a^{10}u + \dots - 16.3639a - 1.17950 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 1.83049a^{11}u + 3.69414a^{10}u + \dots + 10.1122a + 0.883061 \\ 2.96180a^{11}u + 5.97724a^{10}u + \dots + 16.3639a + 1.17950 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -u \\ -u \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes

$$= \frac{460629544778988}{234092030786251} a^{11} u + \frac{789216386414344}{234092030786251} a^{10} u + \dots - \frac{2682835470744820}{234092030786251} a + \frac{2366946904870098}{234092030786251}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$(u^4 + u^3 + 2u^2 + 4u + 1)^6$
c_2, c_4	$(u^4 - u^3 + 2u - 1)^6$
c_3, c_8	$(u^2 + u - 1)^{12}$
c_5, c_7, c_{10} c_{11}	$u^{24} + u^{23} + \dots - 30u + 59$
c_6, c_9	$u^{24} + 3u^{23} + \dots + 226u + 59$
c_{12}	$(u^3 - u^2 + 1)^8$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$(y^4 + 3y^3 - 2y^2 - 12y + 1)^6$
c_2, c_4	$(y^4 - y^3 + 2y^2 - 4y + 1)^6$
c_3, c_8	$(y^2 - 3y + 1)^{12}$
c_5, c_7, c_{10} c_{11}	$y^{24} + 21y^{23} + \dots - 1844y + 3481$
c_6, c_9	$y^{24} - 7y^{23} + \dots - 22992y + 3481$
c_{12}	$(y^3 - y^2 + 2y - 1)^8$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_4^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.618034$ $a = -0.150728 + 0.935298I$ $b = -0.837750 - 0.491196I$	$-0.39223 + 2.82812I$	$2.49024 - 2.97945I$
$u = -0.618034$ $a = -0.150728 - 0.935298I$ $b = -0.837750 + 0.491196I$	$-0.39223 - 2.82812I$	$2.49024 + 2.97945I$
$u = -0.618034$ $a = 0.423709 + 0.723737I$ $b = 0.589608 - 1.016880I$	$-0.39223 + 2.82812I$	$2.49024 - 2.97945I$
$u = -0.618034$ $a = 0.423709 - 0.723737I$ $b = 0.589608 + 1.016880I$	$-0.39223 - 2.82812I$	$2.49024 + 2.97945I$
$u = -0.618034$ $a = 0.361623 + 0.378726I$ $b = -0.328718 + 0.941057I$	3.74535	$9.01951 + 0.I$
$u = -0.618034$ $a = 0.361623 - 0.378726I$ $b = -0.328718 - 0.941057I$	3.74535	$9.01951 + 0.I$
$u = -0.618034$ $a = -2.38323 + 1.43737I$ $b = 0.492907 + 0.249410I$	$-0.39223 - 2.82812I$	$2.49024 + 2.97945I$
$u = -0.618034$ $a = -2.38323 - 1.43737I$ $b = 0.492907 - 0.249410I$	$-0.39223 + 2.82812I$	$2.49024 - 2.97945I$
$u = -0.618034$ $a = -1.67739 + 5.70634I$ $b = 0.15263 - 1.41932I$	3.74535	$9.01951 + 0.I$
$u = -0.618034$ $a = -1.67739 - 5.70634I$ $b = 0.15263 + 1.41932I$	3.74535	$9.01951 + 0.I$

Solutions to I_4^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.618034$ $a = 1.11700 + 6.25809I$ $b = -0.377692 - 0.949629I$	$-0.39223 - 2.82812I$	0
$u = -0.618034$ $a = 1.11700 - 6.25809I$ $b = -0.377692 + 0.949629I$	$-0.39223 + 2.82812I$	0
$u = 1.61803$ $a = -0.047191 + 1.301750I$ $b = 0.73365 - 1.98613I$	11.6410	$9.01951 + 0.I$
$u = 1.61803$ $a = -0.047191 - 1.301750I$ $b = 0.73365 + 1.98613I$	11.6410	$9.01951 + 0.I$
$u = 1.61803$ $a = -0.07600 + 1.43046I$ $b = -0.37139 - 1.51467I$	$7.50345 - 2.82812I$	$2.49024 + 2.97945I$
$u = 1.61803$ $a = -0.07600 - 1.43046I$ $b = -0.37139 + 1.51467I$	$7.50345 + 2.82812I$	$2.49024 - 2.97945I$
$u = 1.61803$ $a = -0.483052 + 0.076488I$ $b = -0.737687 + 0.246052I$	$7.50345 - 2.82812I$	$2.49024 + 2.97945I$
$u = 1.61803$ $a = -0.483052 - 0.076488I$ $b = -0.737687 - 0.246052I$	$7.50345 + 2.82812I$	$2.49024 - 2.97945I$
$u = 1.61803$ $a = -0.63148 + 1.43380I$ $b = -0.27264 - 1.42678I$	11.6410	$9.01951 + 0.I$
$u = 1.61803$ $a = -0.63148 - 1.43380I$ $b = -0.27264 + 1.42678I$	11.6410	$9.01951 + 0.I$

Solutions to I_4^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.61803$ $a = -0.165330 + 0.160935I$ $b = 1.52448 - 0.54039I$	$7.50345 + 2.82812I$	$2.49024 - 2.97945I$
$u = 1.61803$ $a = -0.165330 - 0.160935I$ $b = 1.52448 + 0.54039I$	$7.50345 - 2.82812I$	$2.49024 + 2.97945I$
$u = 1.61803$ $a = 0.21207 + 1.76756I$ $b = -0.067397 - 1.386770I$	$7.50345 - 2.82812I$	$2.49024 + 2.97945I$
$u = 1.61803$ $a = 0.21207 - 1.76756I$ $b = -0.067397 + 1.386770I$	$7.50345 + 2.82812I$	$2.49024 - 2.97945I$

$$\mathbf{V. } I_1^v = \langle a, 8v^3 + 12v^2 + b + 10v + 3, 8v^4 + 12v^3 + 12v^2 + 5v + 1 \rangle$$

(i) Arc colorings

$$a_3 = \begin{pmatrix} v \\ 0 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 0 \\ -8v^3 - 12v^2 - 10v - 3 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} 1 \\ -8v^3 - 8v^2 - 8v - 1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 8v^3 + 8v^2 + 8v + 2 \\ 16v^3 + 20v^2 + 18v + 5 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} 8v^3 + 12v^2 + 12v + 4 \\ 8v^3 + 12v^2 + 12v + 5 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -4v^2 - 4v - 3 \\ -4v^2 - 4v - 4 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -8v^3 - 12v^2 - 10v - 3 \\ -8v^3 - 12v^2 - 10v - 3 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -8v^3 - 12v^2 - 12v - 4 \\ -8v^3 - 12v^2 - 12v - 5 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 8v^3 + 12v^2 + 13v + 4 \\ 8v^3 + 12v^2 + 12v + 5 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} v \\ 0 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = $-8v^3 - 5v^2 - 3$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_2	$(u - 1)^4$
c_3, c_8	u^4
c_4	$(u + 1)^4$
c_5, c_7	$u^4 + u^2 - u + 1$
c_6	$u^4 + 2u^3 + 3u^2 + u + 1$
c_9	$u^4 - 2u^3 + 3u^2 - u + 1$
c_{10}, c_{11}	$u^4 + u^2 + u + 1$
c_{12}	$u^4 + 3u^3 + 4u^2 + 3u + 2$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_4	$(y - 1)^4$
c_3, c_8	y^4
c_5, c_7, c_{10} c_{11}	$y^4 + 2y^3 + 3y^2 + y + 1$
c_6, c_9	$y^4 + 2y^3 + 7y^2 + 5y + 1$
c_{12}	$y^4 - y^3 + 2y^2 + 7y + 4$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^v	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$v = -0.447562 + 0.776246I$	$-2.62503 - 1.39709I$	$-6.74392 + 3.48426I$
$a = 0$		
$b = 0.547424 + 0.585652I$		
$v = -0.447562 - 0.776246I$	$-2.62503 + 1.39709I$	$-6.74392 - 3.48426I$
$a = 0$		
$b = 0.547424 - 0.585652I$		
$v = -0.302438 + 0.253422I$	$0.98010 - 7.64338I$	$-3.38108 + 0.34032I$
$a = 0$		
$b = -0.547424 - 1.120870I$		
$v = -0.302438 - 0.253422I$	$0.98010 + 7.64338I$	$-3.38108 - 0.34032I$
$a = 0$		
$b = -0.547424 + 1.120870I$		

$$\text{VI. } I_2^v = \langle a, b^6 + b^5 + 2b^4 + 2b^3 + 2b^2 + 2b + 1, v - 1 \rangle$$

(i) Arc colorings

$$a_3 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 0 \\ b \end{pmatrix}$$

$$a_7 = \begin{pmatrix} 1 \\ -b^2 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} b^2 + 1 \\ -b^4 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} b^5 + 2b^3 + b \\ -1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -2b^5 - 3b^3 - b^2 - 2b - 1 \\ -b^5 - b^3 - b^2 - b \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} b \\ b \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -b^5 - 2b^3 - b \\ 1 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} b^5 + 2b^3 + b + 1 \\ -1 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = $4b^3 + 4b - 4$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_2	$(u - 1)^6$
c_3, c_8	u^6
c_4	$(u + 1)^6$
c_5, c_7	$u^6 + u^5 + 2u^4 + 2u^3 + 2u^2 + 2u + 1$
c_6	$u^6 + 3u^5 + 4u^4 + 2u^3 + 1$
c_9	$u^6 - 3u^5 + 4u^4 - 2u^3 + 1$
c_{10}, c_{11}	$u^6 - u^5 + 2u^4 - 2u^3 + 2u^2 - 2u + 1$
c_{12}	$(u^3 - u^2 + 1)^2$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_4	$(y - 1)^6$
c_3, c_8	y^6
c_5, c_7, c_{10} c_{11}	$y^6 + 3y^5 + 4y^4 + 2y^3 + 1$
c_6, c_9	$y^6 - y^5 + 4y^4 - 2y^3 + 8y^2 + 1$
c_{12}	$(y^3 - y^2 + 2y - 1)^2$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^v	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$v = 1.00000$ $a = 0$ $b = 0.498832 + 1.001300I$	$-1.37919 - 2.82812I$	$-7.50976 + 2.97945I$
$v = 1.00000$ $a = 0$ $b = 0.498832 - 1.001300I$	$-1.37919 + 2.82812I$	$-7.50976 - 2.97945I$
$v = 1.00000$ $a = 0$ $b = -0.284920 + 1.115140I$	2.75839	$-6 - 0.980489 + 0.10I$
$v = 1.00000$ $a = 0$ $b = -0.284920 - 1.115140I$	2.75839	$-6 - 0.980489 + 0.10I$
$v = 1.00000$ $a = 0$ $b = -0.713912 + 0.305839I$	$-1.37919 - 2.82812I$	$-7.50976 + 2.97945I$
$v = 1.00000$ $a = 0$ $b = -0.713912 - 0.305839I$	$-1.37919 + 2.82812I$	$-7.50976 - 2.97945I$

VII. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1	$(u-1)^{10}(u^4+u^3+2u^2+4u+1)^6(u^4+u^3+4u^2+u+1)^6$ $\cdot (u^{16}-10u^{15}+\dots-7u+1)(u^{30}+15u^{29}+\dots+5376u+4096)$
c_2	$(u-1)^{10}(u^4-u^3+u+1)^6(u^4-u^3+2u-1)^6$ $\cdot (u^{16}+6u^{15}+\dots-u+1)(u^{30}-5u^{29}+\dots-144u+64)$
c_3	$u^{10}(u^2+u-1)^{12}(u^4-3u^3+\dots-2u+2)^6(u^{16}-3u^{14}+\dots-u+1)$ $\cdot (u^{30}+7u^{29}+\dots+8960u+1024)$
c_4	$(u+1)^{10}(u^4-u^3+u+1)^6(u^4-u^3+2u-1)^6$ $\cdot (u^{16}-6u^{15}+\dots+u+1)(u^{30}-5u^{29}+\dots-144u+64)$
c_5	$(u^4+u^2-u+1)(u^6+u^5+2u^4+2u^3+2u^2+2u+1)$ $\cdot (u^{16}+8u^{14}+\dots+3u+1)(u^{24}+u^{23}+\dots-30u+59)$ $\cdot (u^{24}+u^{23}+\dots-6u+23)(u^{30}+16u^{28}+\dots-4u+1)$
c_6	$(u^4+2u^3+3u^2+u+1)(u^6+3u^5+4u^4+2u^3+1)$ $\cdot (u^{16}-3u^{15}+\dots-6u+1)(u^{24}+3u^{23}+\dots+226u+59)$ $\cdot (u^{24}+3u^{23}+\dots+82u+67)(u^{30}+u^{29}+\dots-3u+1)$
c_7	$(u^4+u^2-u+1)(u^6+u^5+2u^4+2u^3+2u^2+2u+1)$ $\cdot (u^{16}+8u^{14}+\dots-3u+1)(u^{24}+u^{23}+\dots-30u+59)$ $\cdot (u^{24}+u^{23}+\dots-6u+23)(u^{30}+16u^{28}+\dots-4u+1)$
c_8	$u^{10}(u^2+u-1)^{12}(u^4-3u^3+\dots-2u+2)^6(u^{16}-3u^{14}+\dots+u+1)$ $\cdot (u^{30}+7u^{29}+\dots+8960u+1024)$
c_9	$(u^4-2u^3+3u^2-u+1)(u^6-3u^5+4u^4-2u^3+1)$ $\cdot (u^{16}-3u^{15}+\dots-6u+1)(u^{24}+3u^{23}+\dots+226u+59)$ $\cdot (u^{24}+3u^{23}+\dots+82u+67)(u^{30}+u^{29}+\dots-3u+1)$
c_{10}	$(u^4+u^2+u+1)(u^6-u^5+2u^4-2u^3+2u^2-2u+1)$ $\cdot (u^{16}+8u^{14}+\dots-3u+1)(u^{24}+u^{23}+\dots-30u+59)$ $\cdot (u^{24}+u^{23}+\dots-6u+23)(u^{30}+16u^{28}+\dots-4u+1)$
c_{11}	$(u^4+u^2+u+1)(u^6-u^5+2u^4-2u^3+2u^2-2u+1)$ $\cdot (u^{16}+8u^{14}+\dots+3u+1)(u^{24}+u^{23}+\dots-30u+59)$ $\cdot (u^{24}+u^{23}+\dots-6u+23)(u^{30}+16u^{28}+\dots-4u+1)$
c_{12}	$((u^3-u^2+1)^{18})(u^4+3u^3+\dots+3u+2)(u^{16}-5u^{15}+\dots-5u^3+1)$ $\cdot (u^{30}+25u^{29}+\dots+4224u+256)$

VIII. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1	$(y-1)^{10}(y^4+3y^3-2y^2-12y+1)^6(y^4+7y^3+16y^2+7y+1)^6$ $\cdot (y^{16}-2y^{15}+\dots+61y+1)$ $\cdot (y^{30}+5y^{29}+\dots+918224896y+16777216)$
c_2, c_4	$(y-1)^{10}(y^4-y^3+2y^2-4y+1)^6(y^4-y^3+4y^2-y+1)^6$ $\cdot (y^{16}-10y^{15}+\dots-7y+1)(y^{30}-15y^{29}+\dots-5376y+4096)$
c_3, c_8	$y^{10}(y^2-3y+1)^{12}(y^4-3y^3+\dots+8y+4)^6(y^{16}-6y^{15}+\dots+9y+1)$ $\cdot (y^{30}-15y^{29}+\dots-7667712y+1048576)$
c_5, c_7, c_{10} c_{11}	$(y^4+2y^3+3y^2+y+1)(y^6+3y^5+4y^4+2y^3+1)$ $\cdot (y^{16}+16y^{15}+\dots+11y+1)(y^{24}+21y^{23}+\dots+24252y+529)$ $\cdot (y^{24}+21y^{23}+\dots-1844y+3481)(y^{30}+32y^{29}+\dots-14y+1)$
c_6, c_9	$(y^4+2y^3+7y^2+5y+1)(y^6-y^5+4y^4-2y^3+8y^2+1)$ $\cdot (y^{16}-5y^{15}+\dots-14y+1)(y^{24}-7y^{23}+\dots-22992y+3481)$ $\cdot (y^{24}-7y^{23}+\dots-56036y+4489)(y^{30}-13y^{29}+\dots+5y+1)$
c_{12}	$((y^3-y^2+2y-1)^{18})(y^4-y^3+2y^2+7y+4)(y^{16}+9y^{15}+\dots+10y^2+1)$ $\cdot (y^{30}+7y^{29}+\dots+180224y+65536)$