


$12n_{0274}$ ($K12n_{0274}$)



Ideals for irreducible components² of X_{par}

$$\begin{aligned}
 I_1^u &= \langle -u^{12} + 5u^{11} - 18u^{10} + 45u^9 - 87u^8 + 135u^7 - 162u^6 + 154u^5 - 100u^4 + 34u^3 + 12u^2 + 4b - 24u + 8, \\
 &\quad -u^{11} + 5u^{10} - 18u^9 + 41u^8 - 75u^7 + 103u^6 - 110u^5 + 86u^4 - 36u^3 - 2u^2 + 4a + 20u - 12, \\
 &\quad u^{13} - 5u^{12} + 18u^{11} - 45u^{10} + 89u^9 - 141u^8 + 178u^7 - 180u^6 + 134u^5 - 64u^4 + 4u^3 + 24u^2 - 16u + 4 \rangle \\
 I_2^u &= \langle -u^{11}a - 2u^{10}a + \dots - 2a - 3, 3u^{12}a - u^{12} + \dots + 2a^2 - 2, \\
 &\quad u^{13} + 2u^{12} + 7u^{11} + 10u^{10} + 18u^9 + 20u^8 + 21u^7 + 20u^6 + 11u^5 + 10u^4 + u^3 + 4u^2 + 2 \rangle \\
 I_3^u &= \langle -au + 9b + 4a - u + 4, 2a^2 - au + 3u + 5, u^2 + 2 \rangle \\
 I_4^u &= \langle 4b + 2a + u + 2, 2a^2 + 2au + 5, u^2 + 2 \rangle
 \end{aligned}$$

$$I_1^v = \langle a, b^2 - b + 1, v + 1 \rangle$$

$$I_2^v = \langle a, b + v - 1, v^2 - v + 1 \rangle$$

* 6 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 51 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/math/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$\text{I. } I_1^u = \langle -u^{12} + 5u^{11} + \dots + 4b + 8, -u^{11} + 5u^{10} + \dots + 4a - 12, u^{13} - 5u^{12} + \dots - 16u + 4 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_8 &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_4 &= \begin{pmatrix} \frac{1}{4}u^{11} - \frac{5}{4}u^{10} + \dots - 5u + 3 \\ \frac{1}{4}u^{12} - \frac{5}{4}u^{11} + \dots + 6u - 2 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_3 &= \begin{pmatrix} \frac{1}{4}u^{12} - u^{11} + \dots + u + 1 \\ \frac{1}{4}u^{12} - \frac{5}{4}u^{11} + \dots + 6u - 2 \end{pmatrix} \\ a_9 &= \begin{pmatrix} \frac{1}{4}u^{12} - u^{11} + \dots + \frac{1}{2}u^2 - 2u \\ -\frac{1}{4}u^{12} + \frac{3}{4}u^{11} + \dots - u + 1 \end{pmatrix} \\ a_2 &= \begin{pmatrix} \frac{1}{4}u^{12} - \frac{3}{2}u^{11} + \dots + 8u - 2 \\ -\frac{1}{4}u^{12} + \frac{5}{4}u^{11} + \dots - 4u + 1 \end{pmatrix} \\ a_5 &= \begin{pmatrix} -u^{12} + 5u^{11} + \dots - 20u + 7 \\ u^4 + 2u^2 \end{pmatrix} \\ a_7 &= \begin{pmatrix} u \\ u^3 + u \end{pmatrix} \\ a_1 &= \begin{pmatrix} u^2 + 1 \\ u^4 + 2u^2 \end{pmatrix} \\ a_6 &= \begin{pmatrix} \frac{1}{2}u^{12} - \frac{9}{4}u^{11} + \dots + 10u - 4 \\ \frac{1}{4}u^{12} - \frac{5}{4}u^{11} + \dots + 4u - 1 \end{pmatrix} \\ a_{10} &= \begin{pmatrix} \frac{1}{2}u^{12} - \frac{5}{4}u^{11} + \dots - \frac{1}{2}u^2 + u \\ \frac{1}{4}u^{12} - \frac{3}{4}u^{11} + \dots - 2u^2 + u \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes

$$= 2u^{12} - 7u^{11} + 25u^{10} - 55u^9 + 106u^8 - 159u^7 + 190u^6 - 185u^5 + 117u^4 - 42u^3 - 22u^2 + 42u - 14$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{13} + 19u^{12} + \cdots + 15u - 1$
c_2, c_4, c_{10}	$u^{13} - 3u^{12} + \cdots - u + 1$
c_3, c_5, c_9	$u^{13} + u^{12} + 2u^{11} + u^{10} + 6u^9 + 4u^8 + 8u^7 + 3u^6 + 8u^5 + 4u^3 + u^2 + u + 1$
c_6	$u^{13} + u^{12} + \cdots + 1321u + 181$
c_7, c_{11}, c_{12}	$u^{13} - 5u^{12} + \cdots - 16u + 4$
c_8	$u^{13} - u^{12} + \cdots - 39u + 11$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{13} - 61y^{12} + \cdots + 287y - 1$
c_2, c_4, c_{10}	$y^{13} + 19y^{12} + \cdots + 15y - 1$
c_3, c_5, c_9	$y^{13} + 3y^{12} + \cdots - y - 1$
c_6	$y^{13} + 39y^{12} + \cdots + 2051655y - 32761$
c_7, c_{11}, c_{12}	$y^{13} + 11y^{12} + \cdots + 64y - 16$
c_8	$y^{13} - 13y^{12} + \cdots + 1191y - 121$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.148130 + 0.133771I$		
$a = -0.461030 + 0.761015I$	$-12.6764 - 7.1373I$	$-6.57814 + 4.53811I$
$b = -0.953736 - 1.010370I$		
$u = 1.148130 - 0.133771I$		
$a = -0.461030 - 0.761015I$	$-12.6764 + 7.1373I$	$-6.57814 - 4.53811I$
$b = -0.953736 + 1.010370I$		
$u = 0.122498 + 1.377410I$		
$a = -0.69924 + 2.04411I$	$7.00880 - 3.27286I$	$6.62507 + 4.15467I$
$b = -0.201153 - 0.911850I$		
$u = 0.122498 - 1.377410I$		
$a = -0.69924 - 2.04411I$	$7.00880 + 3.27286I$	$6.62507 - 4.15467I$
$b = -0.201153 + 0.911850I$		
$u = -0.12567 + 1.42776I$		
$a = -0.020580 - 1.032540I$	$3.89768 + 2.33726I$	$-4.18156 - 2.46985I$
$b = -0.563900 + 0.510841I$		
$u = -0.12567 - 1.42776I$		
$a = -0.020580 + 1.032540I$	$3.89768 - 2.33726I$	$-4.18156 + 2.46985I$
$b = -0.563900 - 0.510841I$		
$u = -0.534170$		
$a = 0.577891$	-0.888404	-11.3990
$b = 0.497979$		
$u = 0.67458 + 1.31287I$		
$a = -0.656821 + 0.500993I$	$-9.09521 + 0.75227I$	$-5.18637 - 1.36359I$
$b = 0.999822 - 0.879885I$		
$u = 0.67458 - 1.31287I$		
$a = -0.656821 - 0.500993I$	$-9.09521 - 0.75227I$	$-5.18637 + 1.36359I$
$b = 0.999822 + 0.879885I$		
$u = 0.420458 + 0.308747I$		
$a = 1.11560 - 0.96767I$	$1.70562 - 1.30597I$	$1.36439 + 5.54852I$
$b = 0.108686 + 0.774322I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.420458 - 0.308747I$		
$a = 1.11560 + 0.96767I$	$1.70562 + 1.30597I$	$1.36439 - 5.54852I$
$b = 0.108686 - 0.774322I$		
$u = 0.52709 + 1.45902I$		
$a = 0.43312 - 1.92910I$	$-7.6681 - 13.1014I$	$-3.34370 + 7.13870I$
$b = 0.861292 + 1.076230I$		
$u = 0.52709 - 1.45902I$		
$a = 0.43312 + 1.92910I$	$-7.6681 + 13.1014I$	$-3.34370 - 7.13870I$
$b = 0.861292 - 1.076230I$		

$$\text{II. } I_2^u = \langle -u^{11}a - 2u^{10}a + \dots - 2a - 3, \ 3u^{12}a - u^{12} + \dots + 2a^2 - 2, \ u^{13} + 2u^{12} + \dots + 4u^2 + 2 \rangle$$

(i) **Arc colorings**

$$\begin{aligned} a_8 &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_4 &= \begin{pmatrix} a \\ \frac{1}{2}u^{11}a + u^{10}a + \dots + a + \frac{3}{2} \end{pmatrix} \\ a_{12} &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_3 &= \begin{pmatrix} \frac{1}{2}u^{11}a + u^{10}a + \dots + 2a + \frac{3}{2} \\ \frac{1}{2}u^{11}a + u^{10}a + \dots + a + \frac{3}{2} \end{pmatrix} \\ a_9 &= \begin{pmatrix} -\frac{1}{2}u^{12}a - \frac{3}{2}u^{11}a + \dots - 3u^2 - \frac{5}{2}u \\ -\frac{1}{2}u^{12}a + \frac{1}{2}u^{12} + \dots + a + \frac{1}{2} \end{pmatrix} \\ a_2 &= \begin{pmatrix} -\frac{1}{2}u^{12}a + \frac{1}{4}u^{12} + \dots + \frac{3}{2}a + \frac{5}{2} \\ \frac{1}{2}u^{11}a + \frac{1}{2}u^{12} + \dots + 2a - \frac{1}{2} \end{pmatrix} \\ a_5 &= \begin{pmatrix} u^{12} + u^{11} + \dots + 2a - 3 \\ u^4 + 2u^2 \end{pmatrix} \\ a_7 &= \begin{pmatrix} u \\ u^3 + u \end{pmatrix} \\ a_1 &= \begin{pmatrix} u^2 + 1 \\ u^4 + 2u^2 \end{pmatrix} \\ a_6 &= \begin{pmatrix} \frac{1}{2}u^{12}a + \frac{1}{2}u^{11} + \dots - 2a + \frac{1}{2} \\ \frac{1}{2}u^{11}a - \frac{1}{2}u^{12} + \dots + 2a - \frac{5}{2} \end{pmatrix} \\ a_{10} &= \begin{pmatrix} \frac{1}{2}u^{12}a - \frac{1}{4}u^{12} + \dots + \frac{3}{2}a + \frac{5}{2} \\ \frac{1}{2}u^{11}a + \frac{1}{2}u^{12} + \dots + 2a + \frac{1}{2} \end{pmatrix} \end{aligned}$$

(ii) **Obstruction class** = -1

(iii) **Cusp Shapes**
 $= -u^{12} - 3u^{10} + 4u^9 + 2u^8 + 14u^7 + 15u^6 + 12u^5 + 17u^4 - 2u^3 + 7u^2 - 6u$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{26} + 30u^{25} + \cdots - 1609u + 81$
c_2, c_4, c_{10}	$u^{26} - 6u^{25} + \cdots - 95u + 9$
c_3, c_5, c_9	$u^{26} + 2u^{25} + \cdots - 5u + 3$
c_6	$u^{26} + 3u^{25} + \cdots - 8726u + 1181$
c_7, c_{11}, c_{12}	$(u^{13} + 2u^{12} + \cdots + 4u^2 + 2)^2$
c_8	$u^{26} + u^{25} + \cdots - 456u + 241$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{26} - 50y^{25} + \cdots + 5625167y + 6561$
c_2, c_4, c_{10}	$y^{26} + 30y^{25} + \cdots - 1609y + 81$
c_3, c_5, c_9	$y^{26} + 6y^{25} + \cdots + 95y + 9$
c_6	$y^{26} + 37y^{25} + \cdots - 40866606y + 1394761$
c_7, c_{11}, c_{12}	$(y^{13} + 10y^{12} + \cdots - 16y - 4)^2$
c_8	$y^{26} - 15y^{25} + \cdots - 481230y + 58081$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.190251 + 0.933207I$		
$a = 0.298460 - 0.362265I$	$1.31987 + 0.84014I$	$-1.01632 - 1.58660I$
$b = 0.719395 + 0.860977I$		
$u = -0.190251 + 0.933207I$		
$a = -0.31610 - 2.95092I$	$1.31987 + 0.84014I$	$-1.01632 - 1.58660I$
$b = -0.406685 + 1.088750I$		
$u = -0.190251 - 0.933207I$		
$a = 0.298460 + 0.362265I$	$1.31987 - 0.84014I$	$-1.01632 + 1.58660I$
$b = 0.719395 - 0.860977I$		
$u = -0.190251 - 0.933207I$		
$a = -0.31610 + 2.95092I$	$1.31987 - 0.84014I$	$-1.01632 + 1.58660I$
$b = -0.406685 - 1.088750I$		
$u = -0.522806 + 0.734222I$		
$a = 0.408662 + 0.796811I$	$-1.83264 + 2.12437I$	$-6.63093 - 3.54511I$
$b = 0.798739 - 0.694333I$		
$u = -0.522806 + 0.734222I$		
$a = 0.671809 - 0.426036I$	$-1.83264 + 2.12437I$	$-6.63093 - 3.54511I$
$b = -0.780847 - 0.269963I$		
$u = -0.522806 - 0.734222I$		
$a = 0.408662 - 0.796811I$	$-1.83264 - 2.12437I$	$-6.63093 + 3.54511I$
$b = 0.798739 + 0.694333I$		
$u = -0.522806 - 0.734222I$		
$a = 0.671809 + 0.426036I$	$-1.83264 - 2.12437I$	$-6.63093 + 3.54511I$
$b = -0.780847 + 0.269963I$		
$u = 0.354216 + 1.088690I$		
$a = 0.804166 - 1.057380I$	$1.27349 - 6.51495I$	$-1.32958 + 6.90681I$
$b = 0.780806 + 0.869479I$		
$u = 0.354216 + 1.088690I$		
$a = -0.53542 + 2.66449I$	$1.27349 - 6.51495I$	$-1.32958 + 6.90681I$
$b = -0.376849 - 1.157840I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.354216 - 1.088690I$		
$a = 0.804166 + 1.057380I$	$1.27349 + 6.51495I$	$-1.32958 - 6.90681I$
$b = 0.780806 - 0.869479I$		
$u = 0.354216 - 1.088690I$		
$a = -0.53542 - 2.66449I$	$1.27349 + 6.51495I$	$-1.32958 - 6.90681I$
$b = -0.376849 + 1.157840I$		
$u = -1.16445$		
$a = -0.292211 + 0.542220I$	-12.9066	-6.99580
$b = -0.996994 - 0.942106I$		
$u = -1.16445$		
$a = -0.292211 - 0.542220I$	-12.9066	-6.99580
$b = -0.996994 + 0.942106I$		
$u = 0.475729 + 0.397522I$		
$a = 0.255684 + 0.929938I$	$-0.73347 + 3.14087I$	$-5.81698 - 0.90558I$
$b = 0.587541 - 0.997816I$		
$u = 0.475729 + 0.397522I$		
$a = 0.907624 + 1.022940I$	$-0.73347 + 3.14087I$	$-5.81698 - 0.90558I$
$b = -0.603946 + 0.535433I$		
$u = 0.475729 - 0.397522I$		
$a = 0.255684 - 0.929938I$	$-0.73347 - 3.14087I$	$-5.81698 + 0.90558I$
$b = 0.587541 + 0.997816I$		
$u = 0.475729 - 0.397522I$		
$a = 0.907624 - 1.022940I$	$-0.73347 - 3.14087I$	$-5.81698 + 0.90558I$
$b = -0.603946 - 0.535433I$		
$u = 0.065416 + 1.409480I$		
$a = -0.944118 + 0.771575I$	$5.12373 + 1.84437I$	$0.90018 - 3.35466I$
$b = -0.110601 - 0.414458I$		
$u = 0.065416 + 1.409480I$		
$a = 0.14726 - 1.84752I$	$5.12373 + 1.84437I$	$0.90018 - 3.35466I$
$b = -0.534312 + 0.945546I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.065416 - 1.409480I$		
$a = -0.944118 - 0.771575I$	$5.12373 - 1.84437I$	$0.90018 + 3.35466I$
$b = -0.110601 + 0.414458I$		
$u = 0.065416 - 1.409480I$		
$a = 0.14726 + 1.84752I$	$5.12373 - 1.84437I$	$0.90018 + 3.35466I$
$b = -0.534312 - 0.945546I$		
$u = -0.60008 + 1.40135I$		
$a = -0.751907 - 0.379224I$	$-8.56727 + 6.23778I$	$-4.60847 - 2.87458I$
$b = 1.015400 + 0.803176I$		
$u = -0.60008 + 1.40135I$		
$a = 0.34609 + 1.75574I$	$-8.56727 + 6.23778I$	$-4.60847 - 2.87458I$
$b = 0.908351 - 1.039740I$		
$u = -0.60008 - 1.40135I$		
$a = -0.751907 + 0.379224I$	$-8.56727 - 6.23778I$	$-4.60847 + 2.87458I$
$b = 1.015400 - 0.803176I$		
$u = -0.60008 - 1.40135I$		
$a = 0.34609 - 1.75574I$	$-8.56727 - 6.23778I$	$-4.60847 + 2.87458I$
$b = 0.908351 + 1.039740I$		

$$\text{III. } I_3^u = \langle -au + 9b + 4a - u + 4, \ 2a^2 - au + 3u + 5, \ u^2 + 2 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_8 &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_4 &= \begin{pmatrix} a \\ \frac{1}{9}au - \frac{4}{9}a + \frac{1}{9}u - \frac{4}{9} \end{pmatrix} \\ a_{12} &= \begin{pmatrix} 1 \\ -2 \end{pmatrix} \\ a_3 &= \begin{pmatrix} \frac{1}{9}au + \frac{5}{9}a + \frac{1}{9}u - \frac{4}{9} \\ \frac{1}{9}au - \frac{4}{9}a + \frac{1}{9}u - \frac{4}{9} \end{pmatrix} \\ a_9 &= \begin{pmatrix} \frac{2}{3}au + \frac{1}{3}a + \frac{1}{6}u - \frac{5}{3} \\ \frac{1}{9}au - \frac{4}{9}a + \frac{1}{9}u + \frac{5}{9} \end{pmatrix} \\ a_2 &= \begin{pmatrix} \frac{2}{9}au + \frac{1}{9}a - \frac{5}{18}u - \frac{17}{9} \\ \frac{1}{9}au - \frac{4}{9}a + \frac{1}{9}u + \frac{5}{9} \end{pmatrix} \\ a_5 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_7 &= \begin{pmatrix} u \\ -u \end{pmatrix} \\ a_1 &= \begin{pmatrix} -1 \\ 0 \end{pmatrix} \\ a_6 &= \begin{pmatrix} -\frac{5}{9}au - \frac{7}{9}a - \frac{1}{18}u + \frac{20}{9} \\ -\frac{1}{9}au + \frac{4}{9}a - \frac{1}{9}u - \frac{5}{9} \end{pmatrix} \\ a_{10} &= \begin{pmatrix} a + u \\ \frac{1}{9}au - \frac{4}{9}a + \frac{1}{9}u - \frac{4}{9} \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = $-\frac{8}{9}au + \frac{32}{9}a - \frac{8}{9}u - \frac{4}{9}$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_3, c_4 c_9, c_{10}	$(u^2 - u + 1)^2$
c_2, c_5	$(u^2 + u + 1)^2$
c_6	$u^4 + 2u^3 + u^2 - 6u + 3$
c_7, c_{11}, c_{12}	$(u^2 + 2)^2$
c_8	$u^4 - 2u^3 + u^2 + 6u + 3$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_3 c_4, c_5, c_9 c_{10}	$(y^2 + y + 1)^2$
c_6, c_8	$y^4 - 2y^3 + 31y^2 - 30y + 9$
c_7, c_{11}, c_{12}	$(y + 2)^4$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.414210I$		
$a = 0.61237 - 1.37850I$	$4.93480 + 4.05977I$	$0. - 6.92820I$
$b = -0.500000 + 0.866025I$		
$u = 1.414210I$		
$a = -0.61237 + 2.08560I$	$4.93480 - 4.05977I$	$0. + 6.92820I$
$b = -0.500000 - 0.866025I$		
$u = -1.414210I$		
$a = 0.61237 + 1.37850I$	$4.93480 - 4.05977I$	$0. + 6.92820I$
$b = -0.500000 - 0.866025I$		
$u = -1.414210I$		
$a = -0.61237 - 2.08560I$	$4.93480 + 4.05977I$	$0. - 6.92820I$
$b = -0.500000 + 0.866025I$		

$$\text{IV. } I_4^u = \langle 4b + 2a + u + 2, 2a^2 + 2au + 5, u^2 + 2 \rangle$$

(i) Arc colorings

$$a_8 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} a \\ -\frac{1}{2}a - \frac{1}{4}u - \frac{1}{2} \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 1 \\ -2 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} \frac{1}{2}a - \frac{1}{4}u - \frac{1}{2} \\ -\frac{1}{2}a - \frac{1}{4}u - \frac{1}{2} \end{pmatrix}$$

$$a_9 = \begin{pmatrix} \frac{1}{2}au - a + \frac{1}{2}u + \frac{1}{2} \\ \frac{1}{2}a + \frac{1}{4}u + \frac{1}{2} \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -\frac{1}{4}au - \frac{1}{4}u - \frac{7}{4} \\ -\frac{1}{2}a - \frac{1}{4}u + \frac{1}{2} \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} u \\ -u \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -\frac{1}{2}au + \frac{3}{2}a - \frac{1}{4}u \\ -\frac{1}{2}a - \frac{1}{4}u - \frac{1}{2} \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} \frac{3}{4}au - \frac{1}{4}u + \frac{9}{4} \\ \frac{1}{2}a + \frac{1}{4}u - \frac{1}{2} \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = 0

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_3, c_4 c_9, c_{10}	$(u^2 - u + 1)^2$
c_2, c_5	$(u^2 + u + 1)^2$
c_6	$u^4 - 4u^3 + 4u^2 + 3$
c_7, c_{11}, c_{12}	$(u^2 + 2)^2$
c_8	$u^4 + 4u^3 + 4u^2 + 3$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_3 c_4, c_5, c_9 c_{10}	$(y^2 + y + 1)^2$
c_6, c_8	$y^4 - 8y^3 + 22y^2 + 24y + 9$
c_7, c_{11}, c_{12}	$(y + 2)^4$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_4^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.414210I$		
$a = 1.024940I$	4.93480	0
$b = -0.500000 - 0.866025I$		
$u = -1.414210I$		
$a = -1.024940I$	4.93480	0
$b = -0.500000 + 0.866025I$		
$u = -1.414210I$		
$a = 2.43916I$	4.93480	0
$b = -0.500000 - 0.866025I$		

$$\mathbf{V}. \quad I_1^v = \langle a, \ b^2 - b + 1, \ v + 1 \rangle$$

(i) Arc colorings

$$a_8 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} 0 \\ b \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} b \\ b \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -b \\ -b + 1 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} b \\ b - 1 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -1 \\ -b + 1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 0 \\ -b \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = $8b - 4$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_4, c_5 c_{10}	$u^2 - u + 1$
c_2, c_3, c_6 c_8, c_9	$u^2 + u + 1$
c_7, c_{11}, c_{12}	u^2

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_3 c_4, c_5, c_6 c_8, c_9, c_{10}	$y^2 + y + 1$
c_7, c_{11}, c_{12}	y^2

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^v	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$v = -1.00000$		
$a = 0$	$-4.05977I$	$0. + 6.92820I$
$b = 0.500000 + 0.866025I$		
$v = -1.00000$		
$a = 0$	$4.05977I$	$0. - 6.92820I$
$b = 0.500000 - 0.866025I$		

$$\text{VI. } I_2^v = \langle a, b + v - 1, v^2 - v + 1 \rangle$$

(i) Arc colorings

$$a_8 = \begin{pmatrix} v \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} 0 \\ -v + 1 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -v + 1 \\ -v + 1 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ -v + 1 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -v + 1 \\ -v \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} v \\ 0 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} v \\ -v + 1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -v + 2 \\ -v \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = -6

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_4, c_5 c_{10}	$u^2 - u + 1$
c_2, c_3, c_9	$u^2 + u + 1$
c_6, c_8	$(u - 1)^2$
c_7, c_{11}, c_{12}	u^2

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_3 c_4, c_5, c_9 c_{10}	$y^2 + y + 1$
c_6, c_8	$(y - 1)^2$
c_7, c_{11}, c_{12}	y^2

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^v	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$v = 0.500000 + 0.866025I$		
$a = 0$	0	-6.00000
$b = 0.500000 - 0.866025I$		
$v = 0.500000 - 0.866025I$		
$a = 0$	0	-6.00000
$b = 0.500000 + 0.866025I$		

VII. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1	$((u^2 - u + 1)^6)(u^{13} + 19u^{12} + \dots + 15u - 1)$ $\cdot (u^{26} + 30u^{25} + \dots - 1609u + 81)$
c_2	$((u^2 + u + 1)^6)(u^{13} - 3u^{12} + \dots - u + 1)(u^{26} - 6u^{25} + \dots - 95u + 9)$
c_3, c_9	$(u^2 - u + 1)^4(u^2 + u + 1)^2$ $\cdot (u^{13} + u^{12} + 2u^{11} + u^{10} + 6u^9 + 4u^8 + 8u^7 + 3u^6 + 8u^5 + 4u^3 + u^2 + u + 1)$ $\cdot (u^{26} + 2u^{25} + \dots - 5u + 3)$
c_4, c_{10}	$((u^2 - u + 1)^6)(u^{13} - 3u^{12} + \dots - u + 1)(u^{26} - 6u^{25} + \dots - 95u + 9)$
c_5	$(u^2 - u + 1)^2(u^2 + u + 1)^4$ $\cdot (u^{13} + u^{12} + 2u^{11} + u^{10} + 6u^9 + 4u^8 + 8u^7 + 3u^6 + 8u^5 + 4u^3 + u^2 + u + 1)$ $\cdot (u^{26} + 2u^{25} + \dots - 5u + 3)$
c_6	$(u - 1)^2(u^2 + u + 1)(u^4 - 4u^3 + 4u^2 + 3)(u^4 + 2u^3 + u^2 - 6u + 3)$ $\cdot (u^{13} + u^{12} + \dots + 1321u + 181)(u^{26} + 3u^{25} + \dots - 8726u + 1181)$
c_7, c_{11}, c_{12}	$u^4(u^2 + 2)^4(u^{13} - 5u^{12} + \dots - 16u + 4)(u^{13} + 2u^{12} + \dots + 4u^2 + 2)^2$
c_8	$(u - 1)^2(u^2 + u + 1)(u^4 - 2u^3 + u^2 + 6u + 3)(u^4 + 4u^3 + 4u^2 + 3)$ $\cdot (u^{13} - u^{12} + \dots - 39u + 11)(u^{26} + u^{25} + \dots - 456u + 241)$

VIII. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1	$((y^2 + y + 1)^6)(y^{13} - 61y^{12} + \dots + 287y - 1)$ $\cdot (y^{26} - 50y^{25} + \dots + 5625167y + 6561)$
c_2, c_4, c_{10}	$((y^2 + y + 1)^6)(y^{13} + 19y^{12} + \dots + 15y - 1)$ $\cdot (y^{26} + 30y^{25} + \dots - 1609y + 81)$
c_3, c_5, c_9	$((y^2 + y + 1)^6)(y^{13} + 3y^{12} + \dots - y - 1)(y^{26} + 6y^{25} + \dots + 95y + 9)$
c_6	$(y - 1)^2(y^2 + y + 1)(y^4 - 8y^3 + 22y^2 + 24y + 9)$ $\cdot (y^4 - 2y^3 + 31y^2 - 30y + 9)(y^{13} + 39y^{12} + \dots + 2051655y - 32761)$ $\cdot (y^{26} + 37y^{25} + \dots - 40866606y + 1394761)$
c_7, c_{11}, c_{12}	$y^4(y + 2)^8(y^{13} + 10y^{12} + \dots - 16y - 4)^2$ $\cdot (y^{13} + 11y^{12} + \dots + 64y - 16)$
c_8	$(y - 1)^2(y^2 + y + 1)(y^4 - 8y^3 + 22y^2 + 24y + 9)$ $\cdot (y^4 - 2y^3 + 31y^2 - 30y + 9)(y^{13} - 13y^{12} + \dots + 1191y - 121)$ $\cdot (y^{26} - 15y^{25} + \dots - 481230y + 58081)$