



12n₀₂₈₇ (K12n₀₂₈₇)



A knot diagram¹

Linearized knot diagram



Solving Sequence

$$2,6 \xrightarrow{c_2} 3 \xrightarrow{c_6} 7 \xrightarrow{c_3} 4,10 \xrightarrow{c_7} 8 \xrightarrow{c_1} 1 \xrightarrow{c_{10}} 11 \xrightarrow{c_5} 5 \xrightarrow{c_9} 9 \xrightarrow{c_{12}} 12 \rightsquigarrow c_4, c_8, c_{11}$$

Ideals for irreducible components² of X_{par}

$$I_1^u = \langle -14u^{35} - 85u^{34} + \dots + b - 25, -27u^{35} - 160u^{34} + \dots + 2a - 59, u^{36} + 6u^{35} + \dots + 11u + 2 \rangle$$

$$I_2^u = \langle 2u^{15} + 7u^{13} - u^{12} + 12u^{11} - 4u^{10} + 7u^9 - 5u^8 - 2u^7 - u^6 - 5u^5 + 5u^4 + 3u^2 + b - u, \\ u^{16} + 5u^{14} - u^{13} + 11u^{12} - 5u^{11} + 12u^{10} - 10u^9 + 4u^8 - 8u^7 - 3u^6 + 2u^5 - 3u^4 + 8u^3 - u^2 + a + 3u - 2, \\ u^{17} - u^{16} + 5u^{15} - 4u^{14} + 12u^{13} - 9u^{12} + 16u^{11} - 10u^{10} + 10u^9 - 5u^8 + 3u^6 - 6u^5 + 6u^4 - 4u^3 + 4u^2 - 2u \rangle$$

$$I_3^u = \langle -u^{17} + u^{16} + \dots + b + 1, u^{17}a + 4u^{17} + \dots + a^2 - 5, u^{18} - u^{17} + \dots - u + 1 \rangle$$

* 3 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 89 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/maths/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$\mathbf{I. } I_1^u = \langle -14u^{35} - 85u^{34} + \cdots + b - 25, -27u^{35} - 160u^{34} + \cdots + 2a - 59, u^{36} + 6u^{35} + \cdots + 11u + 2 \rangle$$

(i) Arc colorings

$$a_2 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} 1 \\ -u^2 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} u \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u^4 + u^2 + 1 \\ u^4 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} \frac{27}{2}u^{35} + 80u^{34} + \cdots + \frac{287}{2}u + \frac{59}{2} \\ 14u^{35} + 85u^{34} + \cdots + 135u + 25 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} \frac{5}{2}u^{35} + 15u^{34} + \cdots + \frac{55}{2}u + \frac{13}{2} \\ 3u^{35} + 17u^{34} + \cdots + 27u + 5 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} u^2 + 1 \\ -u^4 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} \frac{17}{2}u^{35} + 51u^{34} + \cdots + \frac{205}{2}u + \frac{41}{2} \\ 5u^{35} + 36u^{34} + \cdots + 85u + 17 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -\frac{1}{2}u^{35} - 5u^{34} + \cdots + \frac{1}{2}u + \frac{3}{2} \\ -2u^{35} - 14u^{34} + \cdots - 19u - 3 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} \frac{7}{2}u^{35} + 21u^{34} + \cdots + \frac{85}{2}u + \frac{17}{2} \\ u^{35} + 6u^{34} + \cdots + 30u + 7 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} \frac{13}{2}u^{35} + 38u^{34} + \cdots + \frac{121}{2}u + \frac{27}{2} \\ 11u^{35} + 63u^{34} + \cdots + 69u + 11 \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes

$$= -10u^{35} - 47u^{34} - 193u^{33} - 528u^{32} - 1308u^{31} - 2680u^{30} - 5100u^{29} - 8677u^{28} - 13841u^{27} - 20439u^{26} - 28425u^{25} - 37282u^{24} - 46259u^{23} - 54570u^{22} - 61115u^{21} - 65180u^{20} - 66272u^{19} - 64122u^{18} - 59268u^{17} - 52003u^{16} - 43524u^{15} - 34581u^{14} - 26106u^{13} - 18689u^{12} - 12638u^{11} - 8103u^{10} - 4980u^9 - 2882u^8 - 1585u^7 - 744u^6 - 322u^5 - 123u^4 - 56u^3 - 11u^2 + 11u + 16$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{36} + 18u^{35} + \dots + 11u + 4$
c_2, c_6	$u^{36} - 6u^{35} + \dots - 11u + 2$
c_3	$u^{36} + 6u^{35} + \dots + 721u + 74$
c_4, c_{12}	$u^{36} - 28u^{34} + \dots + u + 1$
c_5, c_9	$u^{36} - 8u^{34} + \dots - 19u + 23$
c_7, c_{10}	$u^{36} - 3u^{35} + \dots - 4u + 1$
c_8, c_{11}	$u^{36} + 17u^{35} + \dots - 1495u + 160$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{36} + 2y^{35} + \cdots + 415y + 16$
c_2, c_6	$y^{36} + 18y^{35} + \cdots + 11y + 4$
c_3	$y^{36} - 20y^{35} + \cdots - 111805y + 5476$
c_4, c_{12}	$y^{36} - 56y^{35} + \cdots - y + 1$
c_5, c_9	$y^{36} - 16y^{35} + \cdots - 5421y + 529$
c_7, c_{10}	$y^{36} + 9y^{35} + \cdots + 16y + 1$
c_8, c_{11}	$y^{36} - 29y^{35} + \cdots - 1047185y + 25600$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.149973 + 0.998728I$ $a = 0.826426 + 0.069933I$ $b = 0.978572 - 0.499137I$	$-2.48443 - 1.62652I$	$-2.33035 + 4.22312I$
$u = -0.149973 - 0.998728I$ $a = 0.826426 - 0.069933I$ $b = 0.978572 + 0.499137I$	$-2.48443 + 1.62652I$	$-2.33035 - 4.22312I$
$u = -0.991952 + 0.201272I$ $a = -0.095835 - 0.534611I$ $b = 0.1012230 - 0.0449035I$	$5.44078 - 0.44653I$	$17.6525 + 14.6221I$
$u = -0.991952 - 0.201272I$ $a = -0.095835 + 0.534611I$ $b = 0.1012230 + 0.0449035I$	$5.44078 + 0.44653I$	$17.6525 - 14.6221I$
$u = 0.722213 + 0.749533I$ $a = -0.458236 - 0.270213I$ $b = -0.182561 + 1.023860I$	$10.02280 + 8.45449I$	$8.41836 - 6.51159I$
$u = 0.722213 - 0.749533I$ $a = -0.458236 + 0.270213I$ $b = -0.182561 - 1.023860I$	$10.02280 - 8.45449I$	$8.41836 + 6.51159I$
$u = 0.704411 + 0.851124I$ $a = 0.304244 - 0.316418I$ $b = -0.793602 - 0.473087I$	$9.73048 - 3.08341I$	$8.49473 + 1.45967I$
$u = 0.704411 - 0.851124I$ $a = 0.304244 + 0.316418I$ $b = -0.793602 + 0.473087I$	$9.73048 + 3.08341I$	$8.49473 - 1.45967I$
$u = -0.839708 + 0.298444I$ $a = -2.16620 + 0.59359I$ $b = -0.604870 - 0.306470I$	$7.48792 + 11.15390I$	$7.02947 - 5.33510I$
$u = -0.839708 - 0.298444I$ $a = -2.16620 - 0.59359I$ $b = -0.604870 + 0.306470I$	$7.48792 - 11.15390I$	$7.02947 + 5.33510I$

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.459586 + 1.035300I$ $a = -0.189766 + 0.533405I$ $b = 0.983727 + 0.590961I$	$-0.955654 + 0.821545I$	$0. - 3.95614I$
$u = 0.459586 - 1.035300I$ $a = -0.189766 - 0.533405I$ $b = 0.983727 - 0.590961I$	$-0.955654 - 0.821545I$	$0. + 3.95614I$
$u = -0.349884 + 1.107310I$ $a = -0.40589 - 1.81289I$ $b = -1.19270 - 2.50026I$	$-4.05377 + 0.63572I$	$0.484295 - 0.443572I$
$u = -0.349884 - 1.107310I$ $a = -0.40589 + 1.81289I$ $b = -1.19270 + 2.50026I$	$-4.05377 - 0.63572I$	$0.484295 + 0.443572I$
$u = 0.489524 + 1.057530I$ $a = 0.323550 - 0.327963I$ $b = -0.560466 - 0.804802I$	$-0.66063 + 5.65062I$	$0.65729 - 7.68260I$
$u = 0.489524 - 1.057530I$ $a = 0.323550 + 0.327963I$ $b = -0.560466 + 0.804802I$	$-0.66063 - 5.65062I$	$0.65729 + 7.68260I$
$u = -0.477205 + 1.078410I$ $a = -0.219824 + 1.300630I$ $b = -0.24654 + 1.84787I$	$-0.93062 - 3.45682I$	$3.69827 + 2.78118I$
$u = -0.477205 - 1.078410I$ $a = -0.219824 - 1.300630I$ $b = -0.24654 - 1.84787I$	$-0.93062 + 3.45682I$	$3.69827 - 2.78118I$
$u = -0.516662 + 1.118490I$ $a = 0.37409 - 2.28317I$ $b = 0.64651 - 3.41250I$	$-2.89301 - 8.21064I$	$2.73606 + 6.67381I$
$u = -0.516662 - 1.118490I$ $a = 0.37409 + 2.28317I$ $b = 0.64651 + 3.41250I$	$-2.89301 + 8.21064I$	$2.73606 - 6.67381I$

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.235187 + 1.215000I$ $a = 0.269343 + 1.318210I$ $b = 1.29431 + 2.00975I$	$2.56736 + 7.86975I$	$1.57367 - 3.75977I$
$u = -0.235187 - 1.215000I$ $a = 0.269343 - 1.318210I$ $b = 1.29431 - 2.00975I$	$2.56736 - 7.86975I$	$1.57367 + 3.75977I$
$u = 0.437735 + 0.549718I$ $a = 0.890033 + 0.585768I$ $b = 0.312734 - 1.109250I$	$0.56302 + 2.97658I$	$7.22028 + 1.12421I$
$u = 0.437735 - 0.549718I$ $a = 0.890033 - 0.585768I$ $b = 0.312734 + 1.109250I$	$0.56302 - 2.97658I$	$7.22028 - 1.12421I$
$u = -0.579068 + 1.162280I$ $a = -0.18131 + 2.07088I$ $b = -0.66974 + 3.26898I$	$4.9093 - 16.4009I$	$4.03368 + 8.81881I$
$u = -0.579068 - 1.162280I$ $a = -0.18131 - 2.07088I$ $b = -0.66974 - 3.26898I$	$4.9093 + 16.4009I$	$4.03368 - 8.81881I$
$u = -0.645401 + 0.250421I$ $a = 2.56211 - 0.68246I$ $b = 0.796733 - 0.012403I$	$-0.44102 + 3.68764I$	$6.93148 - 2.97663I$
$u = -0.645401 - 0.250421I$ $a = 2.56211 + 0.68246I$ $b = 0.796733 + 0.012403I$	$-0.44102 - 3.68764I$	$6.93148 + 2.97663I$
$u = -0.379298 + 1.270510I$ $a = -0.367307 + 0.599503I$ $b = -0.495507 + 1.221510I$	$0.75632 - 4.99885I$	$0. + 9.16274I$
$u = -0.379298 - 1.270510I$ $a = -0.367307 - 0.599503I$ $b = -0.495507 - 1.221510I$	$0.75632 + 4.99885I$	$0. - 9.16274I$

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.507489 + 0.424879I$		
$a = -0.697425 - 0.253280I$	$1.18510 - 1.52172I$	$5.49872 + 4.62592I$
$b = 0.190814 + 0.745125I$		
$u = 0.507489 - 0.424879I$		
$a = -0.697425 + 0.253280I$	$1.18510 + 1.52172I$	$5.49872 - 4.62592I$
$b = 0.190814 - 0.745125I$		
$u = -0.505579 + 0.388688I$		
$a = -1.192940 + 0.572586I$	$1.086520 - 0.606535I$	$7.97928 + 3.45064I$
$b = -0.480909 + 0.224460I$		
$u = -0.505579 - 0.388688I$		
$a = -1.192940 - 0.572586I$	$1.086520 + 0.606535I$	$7.97928 - 3.45064I$
$b = -0.480909 - 0.224460I$		
$u = -0.651042 + 1.222950I$		
$a = 0.174939 - 0.327220I$	$2.39061 - 5.47394I$	$20.9532 + 20.3381I$
$b = 0.422269 - 0.499103I$		
$u = -0.651042 - 1.222950I$		
$a = 0.174939 + 0.327220I$	$2.39061 + 5.47394I$	$20.9532 - 20.3381I$
$b = 0.422269 + 0.499103I$		

II.

$$I_2^u = \langle 2u^{15} + 7u^{13} + \dots + b - u, u^{16} + 5u^{14} + \dots + a - 2, u^{17} - u^{16} + \dots - 2u + 1 \rangle$$

(i) Arc colorings

$$a_2 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} 1 \\ -u^2 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} u \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u^4 + u^2 + 1 \\ u^4 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -u^{16} - 5u^{14} + \dots - 3u + 2 \\ -2u^{15} - 7u^{13} + \dots - 3u^2 + u \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 2u^{16} - u^{15} + \dots + 4u - 3 \\ u^{16} + 5u^{14} + \dots + 2u - 1 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} u^2 + 1 \\ -u^4 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -2u^{16} + u^{15} + \dots - 4u + 2 \\ -u^{16} - 4u^{14} + \dots - u + 1 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 3u^{16} - 5u^{15} + \dots + 11u - 7 \\ u^{16} - 4u^{15} + \dots + 6u - 5 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -2u^{16} + u^{15} + \dots - 5u + 2 \\ -u^{16} + u^{15} + \dots - 2u + 2 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -5u^{16} + 2u^{15} + \dots - 12u + 8 \\ -2u^{16} - 2u^{15} + \dots - 2u + 2 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = $6u^{15} - 4u^{14} + 24u^{13} - 12u^{12} + 48u^{11} - 20u^{10} + 44u^9 - 6u^8 + 6u^7 + 12u^6 - 21u^5 + 22u^4 - 14u^3 + 10u^2 - u + 6$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{17} - 9u^{16} + \dots - 4u + 1$
c_2	$u^{17} - u^{16} + \dots - 2u + 1$
c_3	$u^{17} + u^{16} + \dots + 2u + 5$
c_4, c_{12}	$u^{17} - 2u^{16} + \dots - 3u + 1$
c_5, c_9	$u^{17} - 8u^{15} + \dots - u + 1$
c_6	$u^{17} + u^{16} + \dots - 2u - 1$
c_7, c_{10}	$u^{17} - 3u^{16} + \dots + 2u + 1$
c_8	$u^{17} + 14u^{16} + \dots + 23u + 5$
c_{11}	$u^{17} - 14u^{16} + \dots + 23u - 5$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{17} + y^{16} + \dots - 8y - 1$
c_2, c_6	$y^{17} + 9y^{16} + \dots - 4y - 1$
c_3	$y^{17} - 13y^{16} + \dots - 86y - 25$
c_4, c_{12}	$y^{17} - 16y^{16} + \dots - 3y - 1$
c_5, c_9	$y^{17} - 16y^{16} + \dots + 9y - 1$
c_7, c_{10}	$y^{17} - 7y^{16} + \dots + 4y - 1$
c_8, c_{11}	$y^{17} - 18y^{16} + \dots - 371y - 25$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.431008 + 0.942233I$ $a = 0.209460 + 0.502451I$ $b = -0.824758 + 0.503665I$	$-0.436145 - 0.105477I$	$5.17968 - 2.38886I$
$u = -0.431008 - 0.942233I$ $a = 0.209460 - 0.502451I$ $b = -0.824758 - 0.503665I$	$-0.436145 + 0.105477I$	$5.17968 + 2.38886I$
$u = -0.874680$ $a = -0.192244$ $b = 0.315231$	5.41472	11.0230
$u = -0.468457 + 0.715892I$ $a = -0.678237 + 0.210938I$ $b = -0.173523 - 0.977457I$	$0.24900 - 3.62004I$	$1.98563 + 8.95063I$
$u = -0.468457 - 0.715892I$ $a = -0.678237 - 0.210938I$ $b = -0.173523 + 0.977457I$	$0.24900 + 3.62004I$	$1.98563 - 8.95063I$
$u = 0.464004 + 1.050320I$ $a = -0.31217 + 3.03749I$ $b = -0.65170 + 4.08273I$	$5.02616 + 3.30361I$	$0.39237 - 4.97703I$
$u = 0.464004 - 1.050320I$ $a = -0.31217 - 3.03749I$ $b = -0.65170 - 4.08273I$	$5.02616 - 3.30361I$	$0.39237 + 4.97703I$
$u = 0.773218 + 0.228170I$ $a = -1.92418 - 0.33198I$ $b = -0.478975 + 0.164410I$	$-2.06386 - 4.46334I$	$0.46360 + 4.84315I$
$u = 0.773218 - 0.228170I$ $a = -1.92418 + 0.33198I$ $b = -0.478975 - 0.164410I$	$-2.06386 + 4.46334I$	$0.46360 - 4.84315I$
$u = 0.305634 + 1.188350I$ $a = 0.051305 - 1.236500I$ $b = 0.65454 - 1.98487I$	$-6.36852 - 1.01335I$	$-5.48366 + 1.62767I$

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.305634 - 1.188350I$		
$a = 0.051305 + 1.236500I$	$-6.36852 + 1.01335I$	$-5.48366 - 1.62767I$
$b = 0.65454 + 1.98487I$		
$u = 0.539603 + 1.156090I$		
$a = -0.00613 - 1.80864I$	$-4.76857 + 9.35771I$	$-2.38330 - 7.93311I$
$b = -0.17533 - 2.86609I$		
$u = 0.539603 - 1.156090I$		
$a = -0.00613 + 1.80864I$	$-4.76857 - 9.35771I$	$-2.38330 + 7.93311I$
$b = -0.17533 + 2.86609I$		
$u = -0.596101 + 1.176370I$		
$a = -0.0194796 - 0.1364700I$	$2.18794 - 5.27328I$	$-0.326316 - 0.621275I$
$b = 0.343510 - 0.109245I$		
$u = -0.596101 - 1.176370I$		
$a = -0.0194796 + 0.1364700I$	$2.18794 + 5.27328I$	$-0.326316 + 0.621275I$
$b = 0.343510 + 0.109245I$		
$u = 0.350446 + 0.524920I$		
$a = 2.77555 - 1.60523I$	$6.75650 + 0.40196I$	$5.16061 + 1.96558I$
$b = 1.64862 - 0.37193I$		
$u = 0.350446 - 0.524920I$		
$a = 2.77555 + 1.60523I$	$6.75650 - 0.40196I$	$5.16061 - 1.96558I$
$b = 1.64862 + 0.37193I$		

III.

$$I_3^u = \langle -u^{17} + u^{16} + \dots + b + 1, u^{17}a + 4u^{17} + \dots + a^2 - 5, u^{18} - u^{17} + \dots - u + 1 \rangle$$

(i) Arc colorings

$$a_2 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_3 = \begin{pmatrix} 1 \\ -a^2 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} u \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u^4 + u^2 + 1 \\ u^4 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} a \\ u^{17} - u^{16} + \dots + u - 1 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} -u^{17} + u^{16} + \dots - a - 1 \\ -2u^{17} + 2u^{16} + \dots - 2u + 1 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} u^2 + 1 \\ -u^4 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} u^{17} - u^{16} + \dots + a + 1 \\ 2u^{17} - 2u^{16} + \dots + 2u - 1 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 3u^{17} - 3u^{16} + \dots - a - 4 \\ 4u^{17} - 4u^{16} + \dots + 4u - 3 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} u^{17} - u^{16} + \dots + a + 1 \\ 2u^{17} - 2u^{16} + \dots + u - 1 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -3u^{17} + 2u^{16} + \dots - 3a - 2 \\ -6u^{17} + 5u^{16} + \dots - 6u + 3 \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes = $-4u^{17} + 4u^{16} - 16u^{15} + 12u^{14} - 32u^{13} + 24u^{12} - 36u^{11} + 28u^{10} - 24u^9 + 28u^8 - 12u^7 + 16u^6 - 8u^5 + 8u^4 - 8u^3 - 4u + 14$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$(u^{18} + 9u^{17} + \dots + u + 1)^2$
c_2, c_6	$(u^{18} + u^{17} + \dots + u + 1)^2$
c_3	$(u^{18} - u^{17} + \dots - u + 5)^2$
c_4, c_{12}	$u^{36} + u^{35} + \dots - 4000u + 889$
c_5, c_9	$u^{36} + u^{35} + \dots - 84188u + 25097$
c_7, c_{10}	$u^{36} - 15u^{35} + \dots - 340u + 25$
c_8, c_{11}	$(u^{18} - 15u^{17} + \dots + 29u + 3)^2$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$(y^{18} + y^{17} + \cdots + 9y + 1)^2$
c_2, c_6	$(y^{18} + 9y^{17} + \cdots + y + 1)^2$
c_3	$(y^{18} - 7y^{17} + \cdots - 91y + 25)^2$
c_4, c_{12}	$y^{36} - 45y^{35} + \cdots - 11217180y + 790321$
c_5, c_9	$y^{36} - 25y^{35} + \cdots - 7934994452y + 629859409$
c_7, c_{10}	$y^{36} - 9y^{35} + \cdots + 14100y + 625$
c_8, c_{11}	$(y^{18} - 35y^{17} + \cdots - 211y + 9)^2$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.606951 + 0.762732I$ $a = -0.672810 + 0.417119I$ $b = -0.184656 - 0.365220I$	$1.63218 - 2.36433I$	$8.96106 + 3.34702I$
$u = -0.606951 + 0.762732I$ $a = 0.085009 + 0.384811I$ $b = -0.428365 + 0.927160I$	$1.63218 - 2.36433I$	$8.96106 + 3.34702I$
$u = -0.606951 - 0.762732I$ $a = -0.672810 - 0.417119I$ $b = -0.184656 + 0.365220I$	$1.63218 + 2.36433I$	$8.96106 - 3.34702I$
$u = -0.606951 - 0.762732I$ $a = 0.085009 - 0.384811I$ $b = -0.428365 - 0.927160I$	$1.63218 + 2.36433I$	$8.96106 - 3.34702I$
$u = -0.320154 + 1.065080I$ $a = -1.344920 + 0.399822I$ $b = -0.333397 + 0.527308I$	$2.99038 - 0.58479I$	$3.81506 - 0.42463I$
$u = -0.320154 + 1.065080I$ $a = -0.54509 + 1.40985I$ $b = -1.52871 + 2.64353I$	$2.99038 - 0.58479I$	$3.81506 - 0.42463I$
$u = -0.320154 - 1.065080I$ $a = -1.344920 - 0.399822I$ $b = -0.333397 - 0.527308I$	$2.99038 + 0.58479I$	$3.81506 + 0.42463I$
$u = -0.320154 - 1.065080I$ $a = -0.54509 - 1.40985I$ $b = -1.52871 - 2.64353I$	$2.99038 + 0.58479I$	$3.81506 + 0.42463I$
$u = 0.483861 + 1.030980I$ $a = -0.41599 + 2.66215I$ $b = -0.88355 + 4.52483I$	$5.97153 + 3.09151I$	$11.11493 - 2.77317I$
$u = 0.483861 + 1.030980I$ $a = -0.32117 - 3.24954I$ $b = -0.56235 - 3.23198I$	$5.97153 + 3.09151I$	$11.11493 - 2.77317I$

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.483861 - 1.030980I$ $a = -0.41599 - 2.66215I$ $b = -0.88355 - 4.52483I$	$5.97153 - 3.09151I$	$11.11493 + 2.77317I$
$u = 0.483861 - 1.030980I$ $a = -0.32117 + 3.24954I$ $b = -0.56235 + 3.23198I$	$5.97153 - 3.09151I$	$11.11493 + 2.77317I$
$u = 0.781793 + 0.257942I$ $a = 1.50733 + 0.37688I$ $b = 0.649577 - 0.040046I$	$-0.79783 - 3.98828I$	$8.01934 + 2.30410I$
$u = 0.781793 + 0.257942I$ $a = -1.81527 - 0.39003I$ $b = -0.249789 + 0.261114I$	$-0.79783 - 3.98828I$	$8.01934 + 2.30410I$
$u = 0.781793 - 0.257942I$ $a = 1.50733 - 0.37688I$ $b = 0.649577 + 0.040046I$	$-0.79783 + 3.98828I$	$8.01934 - 2.30410I$
$u = 0.781793 - 0.257942I$ $a = -1.81527 + 0.39003I$ $b = -0.249789 - 0.261114I$	$-0.79783 + 3.98828I$	$8.01934 - 2.30410I$
$u = 0.286599 + 1.176040I$ $a = 0.129696 - 1.018410I$ $b = 0.84736 - 1.88349I$	$-5.21072 - 0.69909I$	$2.61745 - 0.31146I$
$u = 0.286599 + 1.176040I$ $a = 0.111176 + 1.187900I$ $b = -0.28946 + 1.60976I$	$-5.21072 - 0.69909I$	$2.61745 - 0.31146I$
$u = 0.286599 - 1.176040I$ $a = 0.129696 + 1.018410I$ $b = 0.84736 + 1.88349I$	$-5.21072 + 0.69909I$	$2.61745 + 0.31146I$
$u = 0.286599 - 1.176040I$ $a = 0.111176 - 1.187900I$ $b = -0.28946 - 1.60976I$	$-5.21072 + 0.69909I$	$2.61745 + 0.31146I$

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.527745 + 1.103190I$ $a = 1.191070 + 0.466746I$ $b = 2.91002 + 0.99091I$	$4.40791 - 6.64525I$	$7.35959 + 7.71274I$
$u = -0.527745 + 1.103190I$ $a = 1.43943 + 1.42860I$ $b = 0.83094 + 1.94919I$	$4.40791 - 6.64525I$	$7.35959 + 7.71274I$
$u = -0.527745 - 1.103190I$ $a = 1.191070 - 0.466746I$ $b = 2.91002 - 0.99091I$	$4.40791 + 6.64525I$	$7.35959 - 7.71274I$
$u = -0.527745 - 1.103190I$ $a = 1.43943 - 1.42860I$ $b = 0.83094 - 1.94919I$	$4.40791 + 6.64525I$	$7.35959 - 7.71274I$
$u = 0.500651 + 0.525564I$ $a = -2.56302 - 0.46993I$ $b = -2.37571 + 0.29310I$	$7.49746 + 0.97328I$	$14.1139 - 4.5518I$
$u = 0.500651 + 0.525564I$ $a = 3.00132 - 1.06603I$ $b = 0.551934 - 0.024389I$	$7.49746 + 0.97328I$	$14.1139 - 4.5518I$
$u = 0.500651 - 0.525564I$ $a = -2.56302 + 0.46993I$ $b = -2.37571 - 0.29310I$	$7.49746 - 0.97328I$	$14.1139 + 4.5518I$
$u = 0.500651 - 0.525564I$ $a = 3.00132 + 1.06603I$ $b = 0.551934 + 0.024389I$	$7.49746 - 0.97328I$	$14.1139 + 4.5518I$
$u = 0.548853 + 1.153160I$ $a = 0.00604 - 1.65136I$ $b = -0.30591 - 2.86421I$	$-3.42686 + 8.95499I$	$4.97585 - 5.84784I$
$u = 0.548853 + 1.153160I$ $a = 0.22040 + 1.64693I$ $b = 0.40384 + 2.31431I$	$-3.42686 + 8.95499I$	$4.97585 - 5.84784I$

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.548853 - 1.153160I$ $a = 0.00604 + 1.65136I$ $b = -0.30591 + 2.86421I$	$-3.42686 - 8.95499I$	$4.97585 + 5.84784I$
$u = 0.548853 - 1.153160I$ $a = 0.22040 - 1.64693I$ $b = 0.40384 - 2.31431I$	$-3.42686 - 8.95499I$	$4.97585 + 5.84784I$
$u = -0.646907 + 0.309141I$ $a = -1.50508 - 0.65497I$ $b = -0.59498 - 1.52441I$	$6.67515 + 2.06052I$	$11.02279 - 4.27827I$
$u = -0.646907 + 0.309141I$ $a = -1.00812 - 2.23459I$ $b = 0.043192 + 0.359954I$	$6.67515 + 2.06052I$	$11.02279 - 4.27827I$
$u = -0.646907 - 0.309141I$ $a = -1.50508 + 0.65497I$ $b = -0.59498 + 1.52441I$	$6.67515 - 2.06052I$	$11.02279 + 4.27827I$
$u = -0.646907 - 0.309141I$ $a = -1.00812 + 2.23459I$ $b = 0.043192 - 0.359954I$	$6.67515 - 2.06052I$	$11.02279 + 4.27827I$

IV. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1	$(u^{17} - 9u^{16} + \dots - 4u + 1)(u^{18} + 9u^{17} + \dots + u + 1)^2$ $\cdot (u^{36} + 18u^{35} + \dots + 11u + 4)$
c_2	$(u^{17} - u^{16} + \dots - 2u + 1)(u^{18} + u^{17} + \dots + u + 1)^2$ $\cdot (u^{36} - 6u^{35} + \dots - 11u + 2)$
c_3	$(u^{17} + u^{16} + \dots + 2u + 5)(u^{18} - u^{17} + \dots - u + 5)^2$ $\cdot (u^{36} + 6u^{35} + \dots + 721u + 74)$
c_4, c_{12}	$(u^{17} - 2u^{16} + \dots - 3u + 1)(u^{36} - 28u^{34} + \dots + u + 1)$ $\cdot (u^{36} + u^{35} + \dots - 4000u + 889)$
c_5, c_9	$(u^{17} - 8u^{15} + \dots - u + 1)(u^{36} - 8u^{34} + \dots - 19u + 23)$ $\cdot (u^{36} + u^{35} + \dots - 84188u + 25097)$
c_6	$(u^{17} + u^{16} + \dots - 2u - 1)(u^{18} + u^{17} + \dots + u + 1)^2$ $\cdot (u^{36} - 6u^{35} + \dots - 11u + 2)$
c_7, c_{10}	$(u^{17} - 3u^{16} + \dots + 2u + 1)(u^{36} - 15u^{35} + \dots - 340u + 25)$ $\cdot (u^{36} - 3u^{35} + \dots - 4u + 1)$
c_8	$(u^{17} + 14u^{16} + \dots + 23u + 5)(u^{18} - 15u^{17} + \dots + 29u + 3)^2$ $\cdot (u^{36} + 17u^{35} + \dots - 1495u + 160)$
c_{11}	$(u^{17} - 14u^{16} + \dots + 23u - 5)(u^{18} - 15u^{17} + \dots + 29u + 3)^2$ $\cdot (u^{36} + 17u^{35} + \dots - 1495u + 160)$

V. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1	$(y^{17} + y^{16} + \dots - 8y - 1)(y^{18} + y^{17} + \dots + 9y + 1)^2$ $\cdot (y^{36} + 2y^{35} + \dots + 415y + 16)$
c_2, c_6	$(y^{17} + 9y^{16} + \dots - 4y - 1)(y^{18} + 9y^{17} + \dots + y + 1)^2$ $\cdot (y^{36} + 18y^{35} + \dots + 11y + 4)$
c_3	$(y^{17} - 13y^{16} + \dots - 86y - 25)(y^{18} - 7y^{17} + \dots - 91y + 25)^2$ $\cdot (y^{36} - 20y^{35} + \dots - 111805y + 5476)$
c_4, c_{12}	$(y^{17} - 16y^{16} + \dots - 3y - 1)(y^{36} - 56y^{35} + \dots - y + 1)$ $\cdot (y^{36} - 45y^{35} + \dots - 11217180y + 790321)$
c_5, c_9	$(y^{17} - 16y^{16} + \dots + 9y - 1)$ $\cdot (y^{36} - 25y^{35} + \dots - 7934994452y + 629859409)$ $\cdot (y^{36} - 16y^{35} + \dots - 5421y + 529)$
c_7, c_{10}	$(y^{17} - 7y^{16} + \dots + 4y - 1)(y^{36} - 9y^{35} + \dots + 14100y + 625)$ $\cdot (y^{36} + 9y^{35} + \dots + 16y + 1)$
c_8, c_{11}	$(y^{17} - 18y^{16} + \dots - 371y - 25)(y^{18} - 35y^{17} + \dots - 211y + 9)^2$ $\cdot (y^{36} - 29y^{35} + \dots - 1047185y + 25600)$