


12n₀₇₂₉ (K12n₀₇₂₉)



Ideals for irreducible components² of X_{par}

$$I_1^u = \langle 8.98709 \times 10^{197} u^{57} + 3.28099 \times 10^{198} u^{56} + \dots + 1.56771 \times 10^{200} b + 3.12866 \times 10^{201}, \\ 4.24469 \times 10^{201} u^{57} + 9.55967 \times 10^{201} u^{56} + \dots + 1.61004 \times 10^{203} a + 8.17316 \times 10^{204}, \\ u^{58} + u^{57} + \dots + 7117u - 1027 \rangle$$

$$I_2^u = \langle 2.79079 \times 10^{21} u^{21} + 1.76550 \times 10^{20} u^{20} + \dots + 2.38585 \times 10^{23} b - 1.48787 \times 10^{23}, \\ 4.29144 \times 10^{23} u^{21} + 2.28509 \times 10^{23} u^{20} + \dots + 2.14726 \times 10^{24} a - 1.05273 \times 10^{24}, u^{22} - 10u^{20} + \dots + 73u - 10 \rangle$$

* 2 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 80 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/math/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILS/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$\text{I. } I_1^u = \langle 8.99 \times 10^{197} u^{57} + 3.28 \times 10^{198} u^{56} + \dots + 1.57 \times 10^{200} b + 3.13 \times 10^{201}, 4.24 \times 10^{201} u^{57} + 9.56 \times 10^{201} u^{56} + \dots + 1.61 \times 10^{203} a + 8.17 \times 10^{204}, u^{58} + u^{57} + \dots + 7117u - 1027 \rangle$$

(i) **Arc colorings**

$$\begin{aligned} a_4 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_{10} &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_5 &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_7 &= \begin{pmatrix} -0.0263639u^{57} - 0.0593753u^{56} + \dots + 306.471u - 50.7637 \\ -0.00573262u^{57} - 0.0209285u^{56} + \dots + 140.907u - 19.9569 \end{pmatrix} \\ a_8 &= \begin{pmatrix} -0.0320965u^{57} - 0.0803038u^{56} + \dots + 447.378u - 70.7206 \\ -0.00573262u^{57} - 0.0209285u^{56} + \dots + 140.907u - 19.9569 \end{pmatrix} \\ a_9 &= \begin{pmatrix} -0.0276476u^{57} - 0.0285016u^{56} + \dots - 4.44141u - 21.8460 \\ -0.0133010u^{57} - 0.0190926u^{56} + \dots + 17.6945u - 4.17250 \end{pmatrix} \\ a_3 &= \begin{pmatrix} 0.00962893u^{57} + 0.0347661u^{56} + \dots - 153.709u + 17.1775 \\ 0.00137276u^{57} + 0.0136626u^{56} + \dots - 118.572u + 17.6094 \end{pmatrix} \\ a_2 &= \begin{pmatrix} 0.00962893u^{57} + 0.0347661u^{56} + \dots - 153.709u + 17.1775 \\ -0.0130047u^{57} - 0.0227497u^{56} + \dots + 50.4397u - 8.20641 \end{pmatrix} \\ a_1 &= \begin{pmatrix} -0.00337579u^{57} + 0.0120164u^{56} + \dots - 103.269u + 8.97114 \\ -0.0130047u^{57} - 0.0227497u^{56} + \dots + 50.4397u - 8.20641 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 0.0273854u^{57} + 0.0503706u^{56} + \dots - 200.581u + 48.9873 \\ 0.0113327u^{57} + 0.0369171u^{56} + \dots - 227.216u + 32.7951 \end{pmatrix} \\ a_6 &= \begin{pmatrix} -0.00894778u^{57} - 0.00254877u^{56} + \dots - 150.638u + 30.0737 \\ 0.0204778u^{57} + 0.0335185u^{56} + \dots - 83.8092u + 15.3951 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} -0.0314997u^{57} - 0.0527340u^{56} + \dots + 69.4621u - 13.0531 \\ -0.0300231u^{57} - 0.0787118u^{56} + \dots + 398.575u - 61.1428 \end{pmatrix} \end{aligned}$$

(ii) **Obstruction class = -1**

(iii) **Cusp Shapes** = $-0.0373141u^{57} - 0.119218u^{56} + \dots + 753.945u - 118.284$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{58} + 7u^{57} + \cdots - 22459u + 1219$
c_2	$u^{58} + 4u^{57} + \cdots + 1084u + 167$
c_3	$u^{58} + u^{57} + \cdots - 283u - 43$
c_4	$u^{58} - u^{57} + \cdots - 7117u - 1027$
c_5, c_{12}	$u^{58} + 2u^{57} + \cdots + 1960u + 664$
c_6	$u^{58} + 6u^{57} + \cdots + 10u + 1$
c_7	$u^{58} + 9u^{56} + \cdots - 10008u - 584$
c_8, c_{11}	$u^{58} + u^{57} + \cdots + 1066u + 97$
c_9	$u^{58} + u^{57} + \cdots + u - 1$
c_{10}	$u^{58} - 4u^{57} + \cdots + 129032u - 9829$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{58} - 61y^{57} + \cdots - 240710163y + 1485961$
c_2	$y^{58} - 8y^{57} + \cdots - 98574y + 27889$
c_3	$y^{58} - 9y^{57} + \cdots - 167293y + 1849$
c_4	$y^{58} - 67y^{57} + \cdots - 53586855y + 1054729$
c_5, c_{12}	$y^{58} + 36y^{57} + \cdots - 6938496y + 440896$
c_6	$y^{58} + 8y^{57} + \cdots - 184y + 1$
c_7	$y^{58} + 18y^{57} + \cdots - 5014784y + 341056$
c_8, c_{11}	$y^{58} + 23y^{57} + \cdots + 106020y + 9409$
c_9	$y^{58} - 19y^{57} + \cdots - 27y + 1$
c_{10}	$y^{58} - 52y^{57} + \cdots - 2754353568y + 96609241$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.984047 + 0.106547I$		
$a = 0.732984 + 0.485124I$	$-5.10938 - 4.31500I$	0
$b = -0.71312 + 2.03208I$		
$u = -0.984047 - 0.106547I$		
$a = 0.732984 - 0.485124I$	$-5.10938 + 4.31500I$	0
$b = -0.71312 - 2.03208I$		
$u = 0.951459 + 0.036280I$		
$a = 0.756616 + 0.985473I$	$2.83534 - 4.32094I$	0
$b = -0.14380 + 1.51989I$		
$u = 0.951459 - 0.036280I$		
$a = 0.756616 - 0.985473I$	$2.83534 + 4.32094I$	0
$b = -0.14380 - 1.51989I$		
$u = 0.889685 + 0.338008I$		
$a = 1.06446 - 1.48155I$	$-0.25157 + 3.22294I$	0
$b = -0.428769 - 0.679768I$		
$u = 0.889685 - 0.338008I$		
$a = 1.06446 + 1.48155I$	$-0.25157 - 3.22294I$	0
$b = -0.428769 + 0.679768I$		
$u = 0.938379 + 0.011522I$		
$a = -0.161328 - 0.401628I$	$-1.102690 - 0.062220I$	0
$b = 0.308280 + 0.060433I$		
$u = 0.938379 - 0.011522I$		
$a = -0.161328 + 0.401628I$	$-1.102690 + 0.062220I$	0
$b = 0.308280 - 0.060433I$		
$u = -0.735401 + 0.007028I$		
$a = 1.37029 - 1.13833I$	$4.02435 + 6.74078I$	$-6.40897 - 6.18806I$
$b = -0.71739 - 1.26490I$		
$u = -0.735401 - 0.007028I$		
$a = 1.37029 + 1.13833I$	$4.02435 - 6.74078I$	$-6.40897 + 6.18806I$
$b = -0.71739 + 1.26490I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.611627 + 0.386837I$		
$a = 1.49854 - 0.36748I$	$0.571664 - 1.277470I$	$-8.81085 + 0.40906I$
$b = -0.269457 + 0.990833I$		
$u = 0.611627 - 0.386837I$		
$a = 1.49854 + 0.36748I$	$0.571664 + 1.277470I$	$-8.81085 - 0.40906I$
$b = -0.269457 - 0.990833I$		
$u = -0.543172 + 0.422725I$		
$a = 0.719118 + 0.935767I$	$4.86373 + 2.09347I$	$-6.47153 - 2.48732I$
$b = -1.320790 + 0.324187I$		
$u = -0.543172 - 0.422725I$		
$a = 0.719118 - 0.935767I$	$4.86373 - 2.09347I$	$-6.47153 + 2.48732I$
$b = -1.320790 - 0.324187I$		
$u = 1.208100 + 0.579060I$		
$a = -0.037426 - 0.757251I$	$-0.975445 - 0.218387I$	0
$b = 0.521975 - 0.291948I$		
$u = 1.208100 - 0.579060I$		
$a = -0.037426 + 0.757251I$	$-0.975445 + 0.218387I$	0
$b = 0.521975 + 0.291948I$		
$u = 0.660076$		
$a = 0.366125$	-1.09302	-7.08600
$b = 0.323077$		
$u = -0.554976 + 0.140362I$		
$a = 1.75463 - 1.19601I$	$4.60137 - 0.26241I$	$-5.57820 + 1.37492I$
$b = 0.273584 + 0.107138I$		
$u = -0.554976 - 0.140362I$		
$a = 1.75463 + 1.19601I$	$4.60137 + 0.26241I$	$-5.57820 - 1.37492I$
$b = 0.273584 - 0.107138I$		
$u = 1.34287 + 0.51625I$		
$a = 0.360673 - 0.444881I$	$-1.29465 - 2.40549I$	0
$b = -0.86318 - 2.71719I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.34287 - 0.51625I$		
$a = 0.360673 + 0.444881I$	$-1.29465 + 2.40549I$	0
$b = -0.86318 + 2.71719I$		
$u = -1.09636 + 0.94746I$		
$a = 0.401125 + 0.121349I$	$-4.43749 - 1.69406I$	0
$b = 0.133549 - 1.228370I$		
$u = -1.09636 - 0.94746I$		
$a = 0.401125 - 0.121349I$	$-4.43749 + 1.69406I$	0
$b = 0.133549 + 1.228370I$		
$u = -0.479931 + 0.256201I$		
$a = 0.80829 + 1.69787I$	$4.89868 - 5.92269I$	$-8.28735 + 6.67085I$
$b = 0.357974 - 0.899794I$		
$u = -0.479931 - 0.256201I$		
$a = 0.80829 - 1.69787I$	$4.89868 + 5.92269I$	$-8.28735 - 6.67085I$
$b = 0.357974 + 0.899794I$		
$u = -1.45716$		
$a = -0.652587$	-6.65741	0
$b = 1.66458$		
$u = 1.51660 + 0.17170I$		
$a = -0.397509 - 0.123089I$	$-2.05878 - 4.62210I$	0
$b = 2.78994 - 1.29874I$		
$u = 1.51660 - 0.17170I$		
$a = -0.397509 + 0.123089I$	$-2.05878 + 4.62210I$	0
$b = 2.78994 + 1.29874I$		
$u = 0.159997 + 0.444223I$		
$a = -0.891005 - 0.942470I$	$5.40486 + 3.03261I$	$-1.43535 + 3.06435I$
$b = -0.215135 + 1.292810I$		
$u = 0.159997 - 0.444223I$		
$a = -0.891005 + 0.942470I$	$5.40486 - 3.03261I$	$-1.43535 - 3.06435I$
$b = -0.215135 - 1.292810I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.55519 + 0.00032I$		
$a = -0.587461 - 0.437982I$	$-6.26258 - 2.81974I$	0
$b = 0.67801 - 1.40728I$		
$u = -1.55519 - 0.00032I$		
$a = -0.587461 + 0.437982I$	$-6.26258 + 2.81974I$	0
$b = 0.67801 + 1.40728I$		
$u = 0.127053 + 0.418851I$		
$a = 1.23833 - 1.01400I$	$-0.617637 - 1.214780I$	$-6.10791 + 5.67942I$
$b = -0.126610 + 0.836327I$		
$u = 0.127053 - 0.418851I$		
$a = 1.23833 + 1.01400I$	$-0.617637 + 1.214780I$	$-6.10791 - 5.67942I$
$b = -0.126610 - 0.836327I$		
$u = -1.63145$		
$a = 0.436826$	-9.90939	0
$b = -1.65196$		
$u = 1.76333 + 0.03132I$		
$a = -0.750182 + 0.469345I$	$-4.94459 + 6.18934I$	0
$b = 1.20433 + 1.24812I$		
$u = 1.76333 - 0.03132I$		
$a = -0.750182 - 0.469345I$	$-4.94459 - 6.18934I$	0
$b = 1.20433 - 1.24812I$		
$u = 0.222347 + 0.057423I$		
$a = 4.52994 + 3.59770I$	$1.83627 - 4.60135I$	$-3.06433 + 1.23782I$
$b = -0.062776 - 0.383104I$		
$u = 0.222347 - 0.057423I$		
$a = 4.52994 - 3.59770I$	$1.83627 + 4.60135I$	$-3.06433 - 1.23782I$
$b = -0.062776 + 0.383104I$		
$u = -0.48982 + 1.70817I$		
$a = 0.647515 + 0.422020I$	$6.18677 + 0.35665I$	0
$b = -1.71821 + 0.15561I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.48982 - 1.70817I$		
$a = 0.647515 - 0.422020I$	$6.18677 - 0.35665I$	0
$b = -1.71821 - 0.15561I$		
$u = -1.79356 + 0.11460I$		
$a = -0.715949 + 0.568710I$	$-7.85182 + 4.36860I$	0
$b = 0.49033 + 1.44237I$		
$u = -1.79356 - 0.11460I$		
$a = -0.715949 - 0.568710I$	$-7.85182 - 4.36860I$	0
$b = 0.49033 - 1.44237I$		
$u = 1.62131 + 0.77787I$		
$a = 0.785352 + 0.143559I$	$-2.94636 - 6.95100I$	0
$b = -1.54041 + 1.26825I$		
$u = 1.62131 - 0.77787I$		
$a = 0.785352 - 0.143559I$	$-2.94636 + 6.95100I$	0
$b = -1.54041 - 1.26825I$		
$u = -1.75534 + 0.48896I$		
$a = 0.227519 + 0.805258I$	$0.70429 + 7.95243I$	0
$b = 0.410897 + 0.818197I$		
$u = -1.75534 - 0.48896I$		
$a = 0.227519 - 0.805258I$	$0.70429 - 7.95243I$	0
$b = 0.410897 - 0.818197I$		
$u = -1.80778 + 0.38348I$		
$a = -0.171765 - 0.343297I$	$-3.74606 - 2.72375I$	0
$b = 0.191963 - 0.552646I$		
$u = -1.80778 - 0.38348I$		
$a = -0.171765 + 0.343297I$	$-3.74606 + 2.72375I$	0
$b = 0.191963 + 0.552646I$		
$u = 1.90742 + 0.23418I$		
$a = -0.392220 - 0.010291I$	$-2.82054 + 3.11371I$	0
$b = -0.096402 - 0.583283I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.90742 - 0.23418I$		
$a = -0.392220 + 0.010291I$	$-2.82054 - 3.11371I$	0
$b = -0.096402 + 0.583283I$		
$u = -1.87242 + 0.53585I$		
$a = 0.710787 - 0.294332I$	$-0.6525 + 15.7491I$	0
$b = -1.36569 - 1.62804I$		
$u = -1.87242 - 0.53585I$		
$a = 0.710787 + 0.294332I$	$-0.6525 - 15.7491I$	0
$b = -1.36569 + 1.62804I$		
$u = -1.96832$		
$a = -1.31909$	-11.0216	0
$b = 0.902042$		
$u = 0.12762 + 2.16706I$		
$a = -0.451547 + 0.382090I$	$6.61103 - 6.39087I$	0
$b = 1.61400 - 0.44696I$		
$u = 0.12762 - 2.16706I$		
$a = -0.451547 - 0.382090I$	$6.61103 + 6.39087I$	0
$b = 1.61400 + 0.44696I$		
$u = 1.97864 + 0.91027I$		
$a = 0.389495 + 0.063606I$	$-1.21978 - 5.87932I$	0
$b = -0.011945 + 1.259660I$		
$u = 1.97864 - 0.91027I$		
$a = 0.389495 - 0.063606I$	$-1.21978 + 5.87932I$	0
$b = -0.011945 - 1.259660I$		

$$\text{II. } I_2^u = \langle 2.79 \times 10^{21} u^{21} + 1.77 \times 10^{20} u^{20} + \dots + 2.39 \times 10^{23} b - 1.49 \times 10^{23}, \ 4.29 \times 10^{23} u^{21} + 2.29 \times 10^{23} u^{20} + \dots + 2.15 \times 10^{24} a - 1.05 \times 10^{24}, \ u^{22} - 10u^{20} + \dots + 73u - 9 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_4 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_{10} &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_5 &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_7 &= \begin{pmatrix} -0.199856u^{21} - 0.106419u^{20} + \dots + 19.6045u + 0.490264 \\ -0.0116973u^{21} - 0.000739990u^{20} + \dots - 7.08742u + 0.623622 \end{pmatrix} \\ a_8 &= \begin{pmatrix} -0.211553u^{21} - 0.107159u^{20} + \dots + 12.5171u + 1.11389 \\ -0.0116973u^{21} - 0.000739990u^{20} + \dots - 7.08742u + 0.623622 \end{pmatrix} \\ a_9 &= \begin{pmatrix} -0.336127u^{21} - 0.0767089u^{20} + \dots + 51.7425u - 6.09472 \\ -0.00505495u^{21} + 0.0250633u^{20} + \dots - 1.05021u + 0.800312 \end{pmatrix} \\ a_3 &= \begin{pmatrix} 0.324297u^{21} + 0.300057u^{20} + \dots - 20.7701u + 6.75595 \\ 0.179573u^{21} + 0.111102u^{20} + \dots - 23.5518u + 3.45052 \end{pmatrix} \\ a_2 &= \begin{pmatrix} 0.324297u^{21} + 0.300057u^{20} + \dots - 20.7701u + 6.75595 \\ -0.0962234u^{21} - 0.138807u^{20} + \dots - 4.56630u + 0.750012 \end{pmatrix} \\ a_1 &= \begin{pmatrix} 0.228074u^{21} + 0.161250u^{20} + \dots - 25.3364u + 7.50597 \\ -0.0962234u^{21} - 0.138807u^{20} + \dots - 4.56630u + 0.750012 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 0.318413u^{21} + 0.152269u^{20} + \dots - 39.8613u + 2.56934 \\ -0.0745587u^{21} - 0.0585677u^{20} + \dots + 12.8859u - 2.38557 \end{pmatrix} \\ a_6 &= \begin{pmatrix} 0.417332u^{21} + 0.134515u^{20} + \dots - 84.2017u + 11.3295 \\ 0.0872425u^{21} + 0.0721007u^{20} + \dots - 7.96511u - 0.936957 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} -0.196498u^{21} - 0.237839u^{20} + \dots + 11.6692u - 7.14808 \\ -0.238971u^{21} - 0.175726u^{20} + \dots + 32.6164u - 4.66605 \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = 1

$$(iii) \text{ Cusp Shapes} = -\frac{457412961431615476330833}{238584851591663803618957}u^{21} - \frac{254429452136571387242067}{238584851591663803618957}u^{20} + \dots + \frac{58966595376988644279933534}{238584851591663803618957}u - \frac{10233901156098748389216510}{238584851591663803618957}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{22} - 12u^{21} + \cdots - 9u + 1$
c_2	$u^{22} - u^{21} + \cdots + 72u - 63$
c_3	$u^{22} - u^{20} + \cdots - 7u + 1$
c_4	$u^{22} - 10u^{20} + \cdots + 73u - 9$
c_5	$u^{22} - u^{21} + \cdots + 24u + 8$
c_6	$u^{22} - u^{21} + \cdots - 16u - 3$
c_7	$u^{22} + 3u^{21} + \cdots + 8u - 56$
c_8	$u^{22} - 2u^{21} + \cdots + 2u + 1$
c_9	$u^{22} + 6u^{21} + \cdots + 5u + 1$
c_{10}	$u^{22} - 7u^{21} + \cdots + 20u - 7$
c_{11}	$u^{22} + 2u^{21} + \cdots - 2u + 1$
c_{12}	$u^{22} + u^{21} + \cdots - 24u + 8$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{22} - 22y^{21} + \cdots + 83y + 1$
c_2	$y^{22} - 13y^{21} + \cdots - 12240y + 3969$
c_3	$y^{22} - 2y^{21} + \cdots - 31y + 1$
c_4	$y^{22} - 20y^{21} + \cdots + 395y + 81$
c_5, c_{12}	$y^{22} + 7y^{21} + \cdots + 64y + 64$
c_6	$y^{22} - 5y^{21} + \cdots - 34y + 9$
c_7	$y^{22} + 5y^{21} + \cdots + 39360y + 3136$
c_8, c_{11}	$y^{22} + 6y^{21} + \cdots - 2y + 1$
c_9	$y^{22} - 12y^{21} + \cdots - 13y + 1$
c_{10}	$y^{22} - 21y^{21} + \cdots + 6y + 49$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.074930 + 0.105891I$	$-0.49759 + 2.56520I$	$-11.45169 - 0.03293I$
$a = 1.065490 - 0.834834I$		
$b = -0.116473 - 0.664580I$		
$u = 1.074930 - 0.105891I$	$-0.49759 - 2.56520I$	$-11.45169 + 0.03293I$
$a = 1.065490 + 0.834834I$		
$b = -0.116473 + 0.664580I$		
$u = -0.534288 + 0.431482I$	$-1.81004 - 0.52019I$	$-15.0020 + 3.2051I$
$a = -0.434617 - 1.025970I$		
$b = -0.157192 + 0.475540I$		
$u = -0.534288 - 0.431482I$	$-1.81004 + 0.52019I$	$-15.0020 - 3.2051I$
$a = -0.434617 + 1.025970I$		
$b = -0.157192 - 0.475540I$		
$u = 0.661139 + 0.110313I$	$1.36966 - 5.12657I$	$-11.4526 + 9.5918I$
$a = 1.56420 + 1.41244I$		
$b = -0.229898 + 0.895312I$		
$u = 0.661139 - 0.110313I$	$1.36966 + 5.12657I$	$-11.4526 - 9.5918I$
$a = 1.56420 - 1.41244I$		
$b = -0.229898 - 0.895312I$		
$u = 1.395400 + 0.151449I$	$-6.51584 + 4.02828I$	$-15.4736 - 6.2870I$
$a = -0.635171 + 0.368343I$		
$b = 1.07115 + 1.63083I$		
$u = 1.395400 - 0.151449I$	$-6.51584 - 4.02828I$	$-15.4736 + 6.2870I$
$a = -0.635171 - 0.368343I$		
$b = 1.07115 - 1.63083I$		
$u = -0.51433 + 1.35727I$	$6.84628 + 0.34484I$	$0.584936 + 0.037274I$
$a = 0.844430 + 0.510048I$		
$b = -1.50481 + 0.10273I$		
$u = -0.51433 - 1.35727I$	$6.84628 - 0.34484I$	$0.584936 - 0.037274I$
$a = 0.844430 - 0.510048I$		
$b = -1.50481 - 0.10273I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.07066 + 1.58496I$		
$a = -0.581666 + 0.511900I$	$7.58470 - 6.19359I$	$-1.94969 + 4.30160I$
$b = 1.29498 - 0.63093I$		
$u = 0.07066 - 1.58496I$		
$a = -0.581666 - 0.511900I$	$7.58470 + 6.19359I$	$-1.94969 - 4.30160I$
$b = 1.29498 + 0.63093I$		
$u = 1.62037$		
$a = 0.458428$	-9.98265	-65.6220
$b = -1.67473$		
$u = 1.38984 + 0.85119I$		
$a = -0.299446 + 0.142790I$	$-4.73458 + 2.11091I$	$-18.7205 - 5.4471I$
$b = 0.092686 - 1.119880I$		
$u = 1.38984 - 0.85119I$		
$a = -0.299446 - 0.142790I$	$-4.73458 - 2.11091I$	$-18.7205 + 5.4471I$
$b = 0.092686 + 1.119880I$		
$u = -1.67782 + 0.22721I$		
$a = -0.181061 + 0.126576I$	$-1.70506 + 4.08011I$	$-6.63396 - 1.41336I$
$b = 1.29986 + 1.32033I$		
$u = -1.67782 - 0.22721I$		
$a = -0.181061 - 0.126576I$	$-1.70506 - 4.08011I$	$-6.63396 + 1.41336I$
$b = 1.29986 - 1.32033I$		
$u = 0.124026 + 0.167304I$		
$a = 3.57897 - 0.31377I$	$5.23332 - 3.56725I$	$-5.47085 + 9.86595I$
$b = -0.39813 - 1.50149I$		
$u = 0.124026 - 0.167304I$		
$a = 3.57897 + 0.31377I$	$5.23332 + 3.56725I$	$-5.47085 - 9.86595I$
$b = -0.39813 + 1.50149I$		
$u = -1.82014 + 0.10403I$		
$a = -0.726378 + 0.567727I$	$-8.37280 + 4.49491I$	$-17.7083 - 5.1691I$
$b = 0.55876 + 1.35106I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.82014 - 0.10403I$		
$a = -0.726378 - 0.567727I$	$-8.37280 - 4.49491I$	$-17.7083 + 5.1691I$
$b = 0.55876 - 1.35106I$		
$u = -1.95919$		
$a = -1.29238$	-11.1324	-44.8210
$b = 0.852875$		

III. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1	$(u^{22} - 12u^{21} + \dots - 9u + 1)(u^{58} + 7u^{57} + \dots - 22459u + 1219)$
c_2	$(u^{22} - u^{21} + \dots + 72u - 63)(u^{58} + 4u^{57} + \dots + 1084u + 167)$
c_3	$(u^{22} - u^{20} + \dots - 7u + 1)(u^{58} + u^{57} + \dots - 283u - 43)$
c_4	$(u^{22} - 10u^{20} + \dots + 73u - 9)(u^{58} - u^{57} + \dots - 7117u - 1027)$
c_5	$(u^{22} - u^{21} + \dots + 24u + 8)(u^{58} + 2u^{57} + \dots + 1960u + 664)$
c_6	$(u^{22} - u^{21} + \dots - 16u - 3)(u^{58} + 6u^{57} + \dots + 10u + 1)$
c_7	$(u^{22} + 3u^{21} + \dots + 8u - 56)(u^{58} + 9u^{56} + \dots - 10008u - 584)$
c_8	$(u^{22} - 2u^{21} + \dots + 2u + 1)(u^{58} + u^{57} + \dots + 1066u + 97)$
c_9	$(u^{22} + 6u^{21} + \dots + 5u + 1)(u^{58} + u^{57} + \dots + u - 1)$
c_{10}	$(u^{22} - 7u^{21} + \dots + 20u - 7)(u^{58} - 4u^{57} + \dots + 129032u - 9829)$
c_{11}	$(u^{22} + 2u^{21} + \dots - 2u + 1)(u^{58} + u^{57} + \dots + 1066u + 97)$
c_{12}	$(u^{22} + u^{21} + \dots - 24u + 8)(u^{58} + 2u^{57} + \dots + 1960u + 664)$

IV. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1	$(y^{22} - 22y^{21} + \dots + 83y + 1)$ $\cdot (y^{58} - 61y^{57} + \dots - 240710163y + 1485961)$
c_2	$(y^{22} - 13y^{21} + \dots - 12240y + 3969)$ $\cdot (y^{58} - 8y^{57} + \dots - 98574y + 27889)$
c_3	$(y^{22} - 2y^{21} + \dots - 31y + 1)(y^{58} - 9y^{57} + \dots - 167293y + 1849)$
c_4	$(y^{22} - 20y^{21} + \dots + 395y + 81)$ $\cdot (y^{58} - 67y^{57} + \dots - 53586855y + 1054729)$
c_5, c_{12}	$(y^{22} + 7y^{21} + \dots + 64y + 64)$ $\cdot (y^{58} + 36y^{57} + \dots - 6938496y + 440896)$
c_6	$(y^{22} - 5y^{21} + \dots - 34y + 9)(y^{58} + 8y^{57} + \dots - 184y + 1)$
c_7	$(y^{22} + 5y^{21} + \dots + 39360y + 3136)$ $\cdot (y^{58} + 18y^{57} + \dots - 5014784y + 341056)$
c_8, c_{11}	$(y^{22} + 6y^{21} + \dots - 2y + 1)(y^{58} + 23y^{57} + \dots + 106020y + 9409)$
c_9	$(y^{22} - 12y^{21} + \dots - 13y + 1)(y^{58} - 19y^{57} + \dots - 27y + 1)$
c_{10}	$(y^{22} - 21y^{21} + \dots + 6y + 49)$ $\cdot (y^{58} - 52y^{57} + \dots - 2754353568y + 96609241)$