



12n₀₇₆₄ (K12n₀₇₆₄)



A knot diagram¹

Linearized knot diagram



Solving Sequence

$$7,11 \xrightarrow{c_7} 8 \xrightarrow{c_{11}} 5,12 \xrightarrow{c_4} 4 \xrightarrow{c_{10}} 10 \xrightarrow{c_{12}} 1 \xrightarrow{c_1} 2 \xrightarrow{c_3} 3 \xrightarrow{c_6} 6 \xrightarrow{c_9} 9 \rightsquigarrow c_2, c_5, c_8$$

Ideals for irreducible components² of X_{par}

$$I_1^u = \langle -521239u^{17} - 378608u^{16} + \dots + 1930047b + 369959, \\ -4207072u^{17} - 369959u^{16} + \dots + 1930047a + 27427427, u^{18} + 9u^{16} + \dots - 7u + 1 \rangle$$

$$I_2^u = \langle u^2 + b, u^6 - u^5 + 2u^4 - 4u^3 + u^2 + a - 2u + 2, u^7 - u^6 + 2u^5 - 4u^4 + u^3 - 2u^2 + u + 1 \rangle$$

$$I_3^u = \langle -47u^{11} - 232u^{10} + \dots + 125b - 849, 3669u^{11} + 13414u^{10} + \dots + 14875a + 13098, \\ u^{12} + 2u^{11} + 8u^{10} + 12u^9 + 29u^8 + 33u^7 + 54u^6 + 51u^5 + 54u^4 + 43u^3 + 28u^2 + 15u + 7 \rangle$$

$$I_4^u = \langle -2931u^{11} + 6882u^{10} + \dots + 36253b - 67921, 20673u^{11} - 60690u^{10} + \dots + 471289a - 219170, \\ u^{12} - 2u^{11} - 2u^{10} + 4u^9 + 11u^8 - u^7 - 6u^6 - 7u^5 - 14u^4 - 19u^3 + 10u^2 + 25u + 13 \rangle$$

* 4 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 49 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/maths/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

I.

$$I_1^u = \langle -5.21 \times 10^5 u^{17} - 3.79 \times 10^5 u^{16} + \dots + 1.93 \times 10^6 b + 3.70 \times 10^5, -4.21 \times 10^6 u^{17} - 3.70 \times 10^5 u^{16} + \dots + 1.93 \times 10^6 a + 2.74 \times 10^7, u^{18} + 9u^{16} + \dots - 7u + 1 \rangle$$

(i) Arc colorings

$$a_7 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} 2.17978u^{17} + 0.191684u^{16} + \dots + 26.2071u - 14.2108 \\ 0.270065u^{17} + 0.196165u^{16} + \dots - 1.83799u - 0.191684 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -u \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} 2.17978u^{17} + 0.191684u^{16} + \dots + 25.2071u - 14.2108 \\ 0.270065u^{17} + 0.196165u^{16} + \dots - 1.83799u - 0.191684 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -2.64153u^{17} - 0.657915u^{16} + \dots - 25.4168u + 18.5822 \\ -0.576170u^{17} - 0.104194u^{16} + \dots - 0.125886u + 0.849598 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} 7.90979u^{17} + 1.38868u^{16} + \dots + 54.3071u - 42.6181 \\ 0.275909u^{17} + 0.141446u^{16} + \dots + 7.93863u - 3.32710 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 2.85131u^{17} + 0.717915u^{16} + \dots + 18.8121u - 16.5368 \\ 0.0775567u^{17} - 0.0705465u^{16} + \dots + 6.12764u - 1.93842 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -2.77375u^{17} - 0.788461u^{16} + \dots - 12.6845u + 14.5983 \\ -0.0775567u^{17} + 0.0705465u^{16} + \dots - 6.12764u + 1.93842 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 2.06984u^{17} + 0.123529u^{16} + \dots + 25.9420u - 13.8229 \\ 0.380005u^{17} + 0.264320u^{16} + \dots - 1.57289u - 0.579533 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -5.96916u^{17} - 1.48818u^{16} + \dots - 52.2245u + 38.4019 \\ -0.707998u^{17} - 0.0464724u^{16} + \dots - 2.87520u + 2.06771 \end{pmatrix}$$

(ii) Obstruction class = -1

$$(iii) \text{ Cusp Shapes} = -\frac{464573}{91907}u^{17} - \frac{454121}{643349}u^{16} + \dots - \frac{30228500}{643349}u + \frac{10577045}{643349}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{18} + u^{17} + \dots + 10u + 7$
c_2, c_6, c_8	$u^{18} + u^{17} + \dots - 3u - 1$
c_3, c_5, c_9	$u^{18} - 16u^{16} + \dots - 20u - 11$
c_4, c_7, c_{11}	$u^{18} + 9u^{16} + \dots + 7u + 1$
c_{10}	$u^{18} + 7u^{17} + \dots - 57u - 7$
c_{12}	$u^{18} - 4u^{17} + \dots + 24u + 117$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{18} - 29y^{17} + \dots - 1724y + 49$
c_2, c_6, c_8	$y^{18} - 15y^{17} + \dots - 3y + 1$
c_3, c_5, c_9	$y^{18} - 32y^{17} + \dots - 1060y + 121$
c_4, c_7, c_{11}	$y^{18} + 18y^{17} + \dots - 31y + 1$
c_{10}	$y^{18} + y^{17} + \dots - 337y + 49$
c_{12}	$y^{18} - 38y^{17} + \dots - 200412y + 13689$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.017806 + 1.047910I$		
$a = 0.678997 - 0.386228I$	$-0.15119 + 2.09745I$	$-12.63201 - 4.12977I$
$b = -0.748794 - 0.598570I$		
$u = 0.017806 - 1.047910I$		
$a = 0.678997 + 0.386228I$	$-0.15119 - 2.09745I$	$-12.63201 + 4.12977I$
$b = -0.748794 + 0.598570I$		
$u = -0.346086 + 0.994636I$		
$a = -0.819378 + 0.554280I$	$5.46595 - 1.70319I$	$-1.84461 + 1.40793I$
$b = 1.44016 - 0.91249I$		
$u = -0.346086 - 0.994636I$		
$a = -0.819378 - 0.554280I$	$5.46595 + 1.70319I$	$-1.84461 - 1.40793I$
$b = 1.44016 + 0.91249I$		
$u = -0.442971 + 1.297020I$		
$a = -0.306761 - 0.349541I$	$-0.12898 + 2.98542I$	$-11.42448 - 1.55661I$
$b = 0.497178 - 0.425094I$		
$u = -0.442971 - 1.297020I$		
$a = -0.306761 + 0.349541I$	$-0.12898 - 2.98542I$	$-11.42448 + 1.55661I$
$b = 0.497178 + 0.425094I$		
$u = 0.680846 + 1.209390I$		
$a = 0.815864 - 0.142059I$	$-0.03662 - 6.82707I$	$-12.5292 + 8.6563I$
$b = -1.262000 + 0.276114I$		
$u = 0.680846 - 1.209390I$		
$a = 0.815864 + 0.142059I$	$-0.03662 + 6.82707I$	$-12.5292 - 8.6563I$
$b = -1.262000 - 0.276114I$		
$u = -0.76192 + 1.23703I$		
$a = 0.045171 - 1.091190I$	$-13.04820 + 4.46526I$	$-11.23742 - 2.32071I$
$b = -1.337910 - 0.285858I$		
$u = -0.76192 - 1.23703I$		
$a = 0.045171 + 1.091190I$	$-13.04820 - 4.46526I$	$-11.23742 + 2.32071I$
$b = -1.337910 + 0.285858I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.24569 + 1.44514I$ $a = -1.051510 - 0.623676I$ $b = 1.93533 + 0.56642I$	$8.63765 + 4.81272I$	$-14.8469 + 6.2266I$
$u = -0.24569 - 1.44514I$ $a = -1.051510 + 0.623676I$ $b = 1.93533 - 0.56642I$	$8.63765 - 4.81272I$	$-14.8469 - 6.2266I$
$u = -0.172111 + 0.488659I$ $a = -1.21271 + 2.12963I$ $b = 0.783985 - 0.730119I$	$2.63033 - 2.95805I$	$-12.64066 + 3.45085I$
$u = -0.172111 - 0.488659I$ $a = -1.21271 - 2.12963I$ $b = 0.783985 + 0.730119I$	$2.63033 + 2.95805I$	$-12.64066 - 3.45085I$
$u = 0.292901$ $a = 1.34245$ $b = -0.177730$	-0.535468	-18.6140
$u = 0.178761$ $a = -8.26850$ $b = -0.442984$	-7.55325	4.27620
$u = 1.03430 + 1.57774I$ $a = 0.813344 - 0.058598I$ $b = -1.99758 + 1.15997I$	$-11.6616 - 12.5169I$	$-11.17612 + 5.12731I$
$u = 1.03430 - 1.57774I$ $a = 0.813344 + 0.058598I$ $b = -1.99758 - 1.15997I$	$-11.6616 + 12.5169I$	$-11.17612 - 5.12731I$

$$\text{II. } I_2^u = \langle u^2 + b, u^6 - u^5 + 2u^4 - 4u^3 + u^2 + a - 2u + 2, u^7 - u^6 + 2u^5 - 4u^4 + u^3 - 2u^2 + u + 1 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_7 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_8 &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_5 &= \begin{pmatrix} -u^6 + u^5 - 2u^4 + 4u^3 - u^2 + 2u - 2 \\ -u^2 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} -u \\ u \end{pmatrix} \\ a_4 &= \begin{pmatrix} -u^6 + u^5 - 2u^4 + 4u^3 - u^2 + u - 2 \\ -u^3 - u^2 \end{pmatrix} \\ a_{10} &= \begin{pmatrix} -u^6 + u^5 - 2u^4 + 4u^3 - u^2 + 3u - 3 \\ u^3 - u^2 + u \end{pmatrix} \\ a_1 &= \begin{pmatrix} u^6 - 2u^5 + 5u^4 - 9u^3 + 8u^2 - 10u + 5 \\ u^6 - 2u^5 + u^4 - 4u^3 + u^2 + u + 1 \end{pmatrix} \\ a_2 &= \begin{pmatrix} u^6 - u^5 + 3u^4 - 5u^3 + 3u^2 - 5u + 2 \\ u^6 - u^5 + 2u^4 - 3u^3 + u^2 - u \end{pmatrix} \\ a_3 &= \begin{pmatrix} u^4 - 2u^3 + 2u^2 - 4u + 2 \\ u^6 - u^5 + 2u^4 - 3u^3 + u^2 - u \end{pmatrix} \\ a_6 &= \begin{pmatrix} -u^6 + u^5 - u^4 + 4u^3 + u - 2 \\ -u^4 - 2u^2 + u \end{pmatrix} \\ a_9 &= \begin{pmatrix} -2u^6 + 2u^5 - 3u^4 + 9u^3 - 2u^2 + 5u - 6 \\ u^6 + u^5 + u^4 + u^3 - 2u^2 + u \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = 1

$$\text{(iii) Cusp Shapes} = -5u^6 + 5u^5 - 11u^4 + 28u^3 - 15u^2 + 14u - 24$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^7 - 6u^5 - u^4 + 9u^3 + u^2 - 4u + 1$
c_2, c_8	$u^7 + 2u^6 - u^5 + 2u^4 + u^3 - 2u^2 - u - 1$
c_3, c_5	$u^7 + u^6 - 5u^5 + 5u^4 - 9u^3 + 7u^2 - 1$
c_4, c_7	$u^7 - u^6 + 2u^5 - 4u^4 + u^3 - 2u^2 + u + 1$
c_6	$u^7 - 2u^6 - u^5 - 2u^4 + u^3 + 2u^2 - u + 1$
c_9	$u^7 - u^6 - 5u^5 - 5u^4 - 9u^3 - 7u^2 + 1$
c_{10}	$u^7 + 6u^6 + 17u^5 + 26u^4 + 20u^3 + 3u^2 - 5u - 1$
c_{11}	$u^7 + u^6 + 2u^5 + 4u^4 + u^3 + 2u^2 + u - 1$
c_{12}	$u^7 + 5u^6 + 6u^5 + 7u^4 + 18u^3 + 5u^2 + 10u + 7$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^7 - 12y^6 + 54y^5 - 117y^4 + 131y^3 - 71y^2 + 14y - 1$
c_2, c_6, c_8	$y^7 - 6y^6 - 5y^5 + 15y^3 - 2y^2 - 3y - 1$
c_3, c_5, c_9	$y^7 - 11y^6 - 3y^5 + 51y^4 + 13y^3 - 39y^2 + 14y - 1$
c_4, c_7, c_{11}	$y^7 + 3y^6 - 2y^5 - 14y^4 - 9y^3 + 6y^2 + 5y - 1$
c_{10}	$y^7 - 2y^6 + 17y^5 - 42y^4 + 86y^3 - 157y^2 + 31y - 1$
c_{12}	$y^7 - 13y^6 + 2y^5 + 137y^4 + 304y^3 + 237y^2 + 30y - 49$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.180603 + 0.994309I$ $a = -1.17684 - 0.97360I$ $b = 0.956033 + 0.359151I$	$3.86399 + 3.81570I$	$-7.59315 - 5.07181I$
$u = -0.180603 - 0.994309I$ $a = -1.17684 + 0.97360I$ $b = 0.956033 - 0.359151I$	$3.86399 - 3.81570I$	$-7.59315 + 5.07181I$
$u = 0.799230$ $a = 0.251205$ $b = -0.638768$	-3.11260	-12.2590
$u = 1.42119$ $a = -0.296362$ $b = -2.01977$	-14.5025	-11.1100
$u = -0.22015 + 1.41755I$ $a = -1.106980 - 0.688829I$ $b = 1.96097 + 0.62416I$	$8.82336 + 5.00709I$	$6.2703 - 15.1192I$
$u = -0.22015 - 1.41755I$ $a = -1.106980 + 0.688829I$ $b = 1.96097 - 0.62416I$	$8.82336 - 5.00709I$	$6.2703 + 15.1192I$
$u = -0.418901$ $a = -3.38720$ $b = -0.175478$	-7.75956	-34.9850

$$\text{III. } I_3^u = \langle -47u^{11} - 232u^{10} + \dots + 125b - 849, 3669u^{11} + 13414u^{10} + \dots + 14875a + 13098, u^{12} + 2u^{11} + \dots + 15u + 7 \rangle$$

(i) Arc colorings

$$a_7 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -0.246655u^{11} - 0.901782u^{10} + \dots - 4.03341u - 0.880538 \\ 0.376000u^{11} + 1.85600u^{10} + \dots + 17.0880u + 6.79200 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -u \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -0.154420u^{11} - 0.00719328u^{10} + \dots + 5.20094u + 3.05217 \\ 0.287059u^{11} + 0.957647u^{10} + \dots + 5.79059u + 1.82118 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 0.389513u^{11} + 1.18750u^{10} + \dots + 8.03341u + 3.02339 \\ -0.301647u^{11} - 0.798118u^{10} + \dots - 5.36847u - 3.08094 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} 0.154420u^{11} + 0.00719328u^{10} + \dots - 5.20094u - 3.05217 \\ -0.187294u^{11} - 0.500235u^{10} + \dots - 2.12894u + 0.310118 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 0.0135126u^{11} + 0.331496u^{10} + \dots + 5.94541u + 2.23139 \\ 0.362824u^{11} + 0.871059u^{10} + \dots + 4.34024u + 0.944471 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -0.349311u^{11} - 0.539563u^{10} + \dots + 1.60518u + 1.28692 \\ 0.362824u^{11} + 0.871059u^{10} + \dots + 4.34024u + 0.944471 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -0.346420u^{11} - 0.359193u^{10} + \dots + 7.30494u + 3.98817 \\ 0.475765u^{11} + 1.31341u^{10} + \dots + 5.74965u + 1.92329 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 0.778958u^{11} + 1.90568u^{10} + \dots + 5.02729u + 1.17002 \\ -0.515765u^{11} - 1.55341u^{10} + \dots - 5.26965u - 1.60329 \end{pmatrix}$$

(ii) Obstruction class = 1

$$\text{(iii) Cusp Shapes} = \frac{3466}{2125}u^{11} + \frac{5146}{2125}u^{10} + \frac{20662}{2125}u^9 + \frac{4013}{425}u^8 + \frac{54149}{2125}u^7 + \frac{27374}{2125}u^6 + \frac{10937}{425}u^5 + \frac{7631}{2125}u^4 + \frac{2288}{2125}u^3 - \frac{3202}{425}u^2 - \frac{16867}{2125}u - \frac{37678}{2125}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$(u^3 - 3u^2 + 2u + 1)^4$
c_2, c_8	$u^{12} - u^{11} - u^{10} + u^9 - u^8 + 2u^7 + u^6 - 9u^5 + 15u^4 - 13u^3 + 9u^2 - 4u + 1$
c_3, c_5	$u^{12} + 4u^{11} + \dots + 2u + 1$
c_4, c_7	$u^{12} + 2u^{11} + \dots + 15u + 7$
c_6	$u^{12} + u^{11} - u^{10} - u^9 - u^8 - 2u^7 + u^6 + 9u^5 + 15u^4 + 13u^3 + 9u^2 + 4u + 1$
c_9	$u^{12} - 4u^{11} + \dots - 2u + 1$
c_{10}	$(u^2 - u + 1)^6$
c_{11}	$u^{12} - 2u^{11} + \dots - 15u + 7$
c_{12}	$u^{12} - 4u^{11} + \dots + 21u + 7$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$(y^3 - 5y^2 + 10y - 1)^4$
c_2, c_6, c_8	$y^{12} - 3y^{11} + \dots + 2y + 1$
c_3, c_5, c_9	$y^{12} - 2y^{11} + \dots + 26y + 1$
c_4, c_7, c_{11}	$y^{12} + 12y^{11} + \dots + 167y + 49$
c_{10}	$(y^2 + y + 1)^6$
c_{12}	$y^{12} + 4y^{11} + \dots + 105y + 49$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.180135 + 0.927755I$ $a = 0.798712 + 0.694015I$ $b = -1.29718 + 1.09845I$	$0.265740 - 0.798239I$	$-10.21508 - 2.15696I$
$u = 0.180135 - 0.927755I$ $a = 0.798712 - 0.694015I$ $b = -1.29718 - 1.09845I$	$0.265740 + 0.798239I$	$-10.21508 + 2.15696I$
$u = -0.206453 + 1.188860I$ $a = -0.636230 + 0.531056I$ $b = 1.80380 - 0.74655I$	$4.40332 - 2.02988I$	$-10.56984 + 3.46410I$
$u = -0.206453 - 1.188860I$ $a = -0.636230 - 0.531056I$ $b = 1.80380 + 0.74655I$	$4.40332 + 2.02988I$	$-10.56984 - 3.46410I$
$u = 0.300960 + 1.170050I$ $a = -0.797324 + 0.222244I$ $b = 1.87013 + 0.59068I$	$0.26574 - 4.85801I$	$-10.21508 + 4.77124I$
$u = 0.300960 - 1.170050I$ $a = -0.797324 - 0.222244I$ $b = 1.87013 - 0.59068I$	$0.26574 + 4.85801I$	$-10.21508 - 4.77124I$
$u = -0.670986 + 0.330909I$ $a = 0.087396 + 1.333780I$ $b = 0.828521 - 0.773223I$	$4.40332 - 2.02988I$	$-10.56984 + 3.46410I$
$u = -0.670986 - 0.330909I$ $a = 0.087396 - 1.333780I$ $b = 0.828521 + 0.773223I$	$4.40332 + 2.02988I$	$-10.56984 - 3.46410I$
$u = 0.40365 + 1.40633I$ $a = -0.663216 + 0.165175I$ $b = 0.835953 - 0.124983I$	$0.265740 + 0.798239I$	$-10.21508 + 2.15696I$
$u = 0.40365 - 1.40633I$ $a = -0.663216 - 0.165175I$ $b = 0.835953 + 0.124983I$	$0.265740 - 0.798239I$	$-10.21508 - 2.15696I$

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.00731 + 1.43634I$	$0.26574 + 4.85801I$	$-10.21508 - 4.77124I$
$a = 0.567805 - 0.050096I$		
$b = -1.041220 - 0.420474I$		
$u = -1.00731 - 1.43634I$	$0.26574 - 4.85801I$	$-10.21508 + 4.77124I$
$a = 0.567805 + 0.050096I$		
$b = -1.041220 + 0.420474I$		

$$\text{IV. } I_4^u = \langle -2931u^{11} + 6882u^{10} + \dots + 36253b - 67921, 20673u^{11} - 60690u^{10} + \dots + 471289a - 219170, u^{12} - 2u^{11} + \dots + 25u + 13 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_7 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_8 &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_5 &= \begin{pmatrix} -0.0438648u^{11} + 0.128774u^{10} + \dots - 0.996698u + 0.465044 \\ 0.0808485u^{11} - 0.189833u^{10} + \dots + 2.60001u + 1.87353 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} -u \\ u \end{pmatrix} \\ a_4 &= \begin{pmatrix} 0.0941100u^{11} - 0.160829u^{10} + \dots + 2.05919u + 2.87215 \\ 0.0281356u^{11} + 0.0393623u^{10} + \dots + 1.14768u + 2.05103 \end{pmatrix} \\ a_{10} &= \begin{pmatrix} 0.120788u^{11} - 0.282621u^{10} + \dots + 1.76593u + 1.45803 \\ 0.0273908u^{11} - 0.164621u^{10} + \dots + 0.519405u - 1.22343 \end{pmatrix} \\ a_1 &= \begin{pmatrix} -0.0941100u^{11} + 0.160829u^{10} + \dots - 2.05919u - 2.87215 \\ -0.0324939u^{11} + 0.0685461u^{10} + \dots + 0.472016u - 0.826442 \end{pmatrix} \\ a_2 &= \begin{pmatrix} -0.133483u^{11} + 0.361165u^{10} + \dots - 2.47857u - 2.33702 \\ -0.190081u^{11} + 0.415166u^{10} + \dots - 2.22379u - 2.79433 \end{pmatrix} \\ a_3 &= \begin{pmatrix} -0.0565980u^{11} + 0.0540008u^{10} + \dots + 0.254780u - 0.457318 \\ 0.190081u^{11} - 0.415166u^{10} + \dots + 2.22379u + 2.79433 \end{pmatrix} \\ a_6 &= \begin{pmatrix} 0.108895u^{11} - 0.270117u^{10} + \dots - 0.193179u + 0.632864 \\ -0.0719113u^{11} + 0.209059u^{10} + \dots + 1.79649u + 1.70571 \end{pmatrix} \\ a_9 &= \begin{pmatrix} 0.177965u^{11} - 0.558920u^{10} + \dots + 2.12679u + 1.81136 \\ 0.514054u^{11} - 1.53965u^{10} + \dots + 9.23946u + 4.75588 \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = -1

$$\text{(iii) Cusp Shapes} = -\frac{3530}{36253}u^{11} + \frac{19484}{36253}u^{10} - \frac{23246}{36253}u^9 - \frac{11803}{36253}u^8 - \frac{17415}{36253}u^7 + \frac{113796}{36253}u^6 + \frac{5597}{36253}u^5 + \frac{68101}{36253}u^4 - \frac{58140}{36253}u^3 - \frac{1074}{36253}u^2 - \frac{209035}{36253}u - \frac{58638}{5179}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$(u^3 - u^2 - 2u + 1)^4$
c_2, c_6, c_8	$u^{12} + u^{11} + \dots - 308u + 91$
c_3, c_5, c_9	$u^{12} + 4u^{11} + \dots + 315u^2 + 189$
c_4, c_7, c_{11}	$u^{12} + 2u^{11} + \dots - 25u + 13$
c_{10}	$(u^2 - u + 1)^6$
c_{12}	$u^{12} - 18u^{10} + \dots - 525u + 127$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$(y^3 - 5y^2 + 6y - 1)^4$
c_2, c_6, c_8	$y^{12} - 31y^{11} + \dots - 27342y + 8281$
c_3, c_5, c_9	$y^{12} - 38y^{11} + \dots + 119070y + 35721$
c_4, c_7, c_{11}	$y^{12} - 8y^{11} + \dots - 365y + 169$
c_{10}	$(y^2 + y + 1)^6$
c_{12}	$y^{12} - 36y^{11} + \dots - 25943y + 16129$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_4^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.935871 + 0.612228I$ $a = 0.798080 - 0.403281I$ $b = -1.27562 - 0.98498I$	$-3.05488 + 2.02988I$	$-11.80194 - 3.46410I$
$u = -0.935871 - 0.612228I$ $a = 0.798080 + 0.403281I$ $b = -1.27562 + 0.98498I$	$-3.05488 - 2.02988I$	$-11.80194 + 3.46410I$
$u = 0.267965 + 1.122990I$ $a = -0.830154 + 0.247151I$ $b = 1.50749 - 0.36507I$	$2.58490 - 2.02988I$	$-8.75302 + 3.46410I$
$u = 0.267965 - 1.122990I$ $a = -0.830154 - 0.247151I$ $b = 1.50749 + 0.36507I$	$2.58490 + 2.02988I$	$-8.75302 - 3.46410I$
$u = -0.668934 + 0.428490I$ $a = 1.118010 - 0.578487I$ $b = -0.304581 + 0.329429I$	$2.58490 + 2.02988I$	$-8.75302 - 3.46410I$
$u = -0.668934 - 0.428490I$ $a = 1.118010 + 0.578487I$ $b = -0.304581 - 0.329429I$	$2.58490 - 2.02988I$	$-8.75302 + 3.46410I$
$u = 1.213350 + 0.131620I$ $a = -0.483814 - 0.661265I$ $b = 0.443182 - 0.504374I$	$-3.05488 - 2.02988I$	$-11.80194 + 3.46410I$
$u = 1.213350 - 0.131620I$ $a = -0.483814 + 0.661265I$ $b = 0.443182 + 0.504374I$	$-3.05488 + 2.02988I$	$-11.80194 - 3.46410I$
$u = -0.894944 + 0.895021I$ $a = 0.763167 - 0.204455I$ $b = -2.34607 - 2.27347I$	$-14.3344 + 2.0299I$	$-10.44504 - 3.46410I$
$u = -0.894944 - 0.895021I$ $a = 0.763167 + 0.204455I$ $b = -2.34607 + 2.27347I$	$-14.3344 - 2.0299I$	$-10.44504 + 3.46410I$

Solutions to I_4^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 2.01843 + 1.05092I$		
$a = -0.019135 + 0.439021I$	$-14.3344 + 2.0299I$	$-10.44504 - 3.46410I$
$b = -1.024400 + 0.327533I$		
$u = 2.01843 - 1.05092I$		
$a = -0.019135 - 0.439021I$	$-14.3344 - 2.0299I$	$-10.44504 + 3.46410I$
$b = -1.024400 - 0.327533I$		

V. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1	$(u^3 - 3u^2 + 2u + 1)^4(u^3 - u^2 - 2u + 1)^4$ $\cdot (u^7 - 6u^5 - u^4 + 9u^3 + u^2 - 4u + 1)(u^{18} + u^{17} + \dots + 10u + 7)$
c_2, c_8	$(u^7 + 2u^6 - u^5 + 2u^4 + u^3 - 2u^2 - u - 1)$ $\cdot (u^{12} - u^{11} - u^{10} + u^9 - u^8 + 2u^7 + u^6 - 9u^5 + 15u^4 - 13u^3 + 9u^2 - 4u + 1)$ $\cdot (u^{12} + u^{11} + \dots - 308u + 91)(u^{18} + u^{17} + \dots - 3u - 1)$
c_3, c_5	$(u^7 + u^6 + \dots + 7u^2 - 1)(u^{12} + 4u^{11} + \dots + 315u^2 + 189)$ $\cdot (u^{12} + 4u^{11} + \dots + 2u + 1)(u^{18} - 16u^{16} + \dots - 20u - 11)$
c_4, c_7	$(u^7 - u^6 + \dots + u + 1)(u^{12} + 2u^{11} + \dots - 25u + 13)$ $\cdot (u^{12} + 2u^{11} + \dots + 15u + 7)(u^{18} + 9u^{16} + \dots + 7u + 1)$
c_6	$(u^7 - 2u^6 + \dots - u + 1)(u^{12} + u^{11} + \dots - 308u + 91)$ $\cdot (u^{12} + u^{11} - u^{10} - u^9 - u^8 - 2u^7 + u^6 + 9u^5 + 15u^4 + 13u^3 + 9u^2 + 4u + 1)$ $\cdot (u^{18} + u^{17} + \dots - 3u - 1)$
c_9	$(u^7 - u^6 + \dots - 7u^2 + 1)(u^{12} - 4u^{11} + \dots - 2u + 1)$ $\cdot (u^{12} + 4u^{11} + \dots + 315u^2 + 189)(u^{18} - 16u^{16} + \dots - 20u - 11)$
c_{10}	$(u^2 - u + 1)^{12}(u^7 + 6u^6 + 17u^5 + 26u^4 + 20u^3 + 3u^2 - 5u - 1)$ $\cdot (u^{18} + 7u^{17} + \dots - 57u - 7)$
c_{11}	$(u^7 + u^6 + \dots + u - 1)(u^{12} - 2u^{11} + \dots - 15u + 7)$ $\cdot (u^{12} + 2u^{11} + \dots - 25u + 13)(u^{18} + 9u^{16} + \dots + 7u + 1)$
c_{12}	$(u^7 + 5u^6 + 6u^5 + 7u^4 + 18u^3 + 5u^2 + 10u + 7)$ $\cdot (u^{12} - 18u^{10} + \dots - 525u + 127)(u^{12} - 4u^{11} + \dots + 21u + 7)$ $\cdot (u^{18} - 4u^{17} + \dots + 24u + 117)$

VI. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1	$(y^3 - 5y^2 + 6y - 1)^4(y^3 - 5y^2 + 10y - 1)^4$ $\cdot (y^7 - 12y^6 + 54y^5 - 117y^4 + 131y^3 - 71y^2 + 14y - 1)$ $\cdot (y^{18} - 29y^{17} + \dots - 1724y + 49)$
c_2, c_6, c_8	$(y^7 - 6y^6 - 5y^5 + 15y^3 - 2y^2 - 3y - 1)$ $\cdot (y^{12} - 31y^{11} + \dots - 27342y + 8281)(y^{12} - 3y^{11} + \dots + 2y + 1)$ $\cdot (y^{18} - 15y^{17} + \dots - 3y + 1)$
c_3, c_5, c_9	$(y^7 - 11y^6 - 3y^5 + 51y^4 + 13y^3 - 39y^2 + 14y - 1)$ $\cdot (y^{12} - 38y^{11} + \dots + 119070y + 35721)(y^{12} - 2y^{11} + \dots + 26y + 1)$ $\cdot (y^{18} - 32y^{17} + \dots - 1060y + 121)$
c_4, c_7, c_{11}	$(y^7 + 3y^6 - 2y^5 - 14y^4 - 9y^3 + 6y^2 + 5y - 1)$ $\cdot (y^{12} - 8y^{11} + \dots - 365y + 169)(y^{12} + 12y^{11} + \dots + 167y + 49)$ $\cdot (y^{18} + 18y^{17} + \dots - 31y + 1)$
c_{10}	$(y^2 + y + 1)^{12}(y^7 - 2y^6 + 17y^5 - 42y^4 + 86y^3 - 157y^2 + 31y - 1)$ $\cdot (y^{18} + y^{17} + \dots - 337y + 49)$
c_{12}	$(y^7 - 13y^6 + 2y^5 + 137y^4 + 304y^3 + 237y^2 + 30y - 49)$ $\cdot (y^{12} - 36y^{11} + \dots - 25943y + 16129)(y^{12} + 4y^{11} + \dots + 105y + 49)$ $\cdot (y^{18} - 38y^{17} + \dots - 200412y + 13689)$