


$12n_{0855}$ ($K12n_{0855}$)



Ideals for irreducible components² of X_{par}

$$I_1^u = \langle -5u^{24} - 19u^{23} + \dots + 2b + 18, -u^{24} - 3u^{23} + \dots + 4a + 4, u^{25} + 5u^{24} + \dots + 10u - 4 \rangle$$

$$I_2^u = \langle 20u^7a^3 - 10u^7a^2 + \dots - 62a + 47, -3u^7a^2 + 6u^7a + \dots - 20a - 11, u^8 - u^7 - 3u^6 + 2u^5 + 3u^4 - 2u - 1 \rangle$$

$$I_3^u = \langle u^{14} - u^{13} - 5u^{12} + 5u^{11} + 8u^{10} - 8u^9 - u^8 + u^7 - 8u^6 + 8u^5 + 3u^4 - 4u^3 + 4u^2 + b - 3u,$$

$$u^{14} - 6u^{12} + 13u^{10} - 9u^8 - 7u^6 + 11u^4 - u^3 + a + u - 3,$$

$$u^{15} - 7u^{13} + 19u^{11} - 22u^9 + 3u^7 + 14u^5 - u^4 - 6u^3 + 2u^2 - 3u - 1 \rangle$$

* 3 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 72 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/math/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$\text{I. } I_1^u = \langle -5u^{24} - 19u^{23} + \dots + 2b + 18, -u^{24} - 3u^{23} + \dots + 4a + 4, u^{25} + 5u^{24} + \dots + 10u - 4 \rangle$$

(i) Arc colorings

$$\begin{aligned} a_5 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ a_{12} &= \begin{pmatrix} 0 \\ u \end{pmatrix} \\ a_6 &= \begin{pmatrix} 1 \\ u^2 \end{pmatrix} \\ a_9 &= \begin{pmatrix} \frac{1}{4}u^{24} + \frac{3}{4}u^{23} + \dots + \frac{25}{4}u - 1 \\ \frac{5}{2}u^{24} + \frac{19}{2}u^{23} + \dots + \frac{61}{2}u - 9 \end{pmatrix} \\ a_1 &= \begin{pmatrix} -7u^{24} - \frac{51}{2}u^{23} + \dots - \frac{131}{2}u + \frac{41}{2} \\ -\frac{45}{2}u^{24} - \frac{161}{2}u^{23} + \dots - \frac{391}{2}u + 62 \end{pmatrix} \\ a_8 &= \begin{pmatrix} \frac{1}{4}u^{24} + \frac{3}{4}u^{23} + \dots + \frac{25}{4}u - 1 \\ \frac{7}{2}u^{24} + \frac{25}{2}u^{23} + \dots + \frac{73}{2}u - 11 \end{pmatrix} \\ a_{11} &= \begin{pmatrix} u \\ u \end{pmatrix} \\ a_4 &= \begin{pmatrix} -u^2 + 1 \\ -u^2 \end{pmatrix} \\ a_2 &= \begin{pmatrix} \frac{1}{2}u^{24} + 3u^{23} + \dots + 11u - \frac{7}{2} \\ -\frac{11}{2}u^{24} - \frac{35}{2}u^{23} + \dots - \frac{55}{2}u + 10 \end{pmatrix} \\ a_3 &= \begin{pmatrix} -7u^{24} - \frac{51}{2}u^{23} + \dots - \frac{131}{2}u + \frac{43}{2} \\ -\frac{45}{2}u^{24} - \frac{161}{2}u^{23} + \dots - \frac{393}{2}u + 62 \end{pmatrix} \\ a_7 &= \begin{pmatrix} u^6 - 3u^4 + 2u^2 + 1 \\ u^8 - 2u^6 + 2u^2 \end{pmatrix} \\ a_{10} &= \begin{pmatrix} -u^3 + 2u \\ -u^5 + u^3 + u \end{pmatrix} \end{aligned}$$

(ii) Obstruction class = -1

$$(iii) \text{ Cusp Shapes} = 11u^{24} + 41u^{23} - 32u^{22} - 241u^{21} + 37u^{20} + 663u^{19} - 146u^{18} - 999u^{17} + 733u^{16} + 604u^{15} - 1604u^{14} + 758u^{13} + 1393u^{12} - 2013u^{11} + 411u^{10} + 1454u^9 - 1775u^8 + 369u^7 + 901u^6 - 1088u^5 + 350u^4 + 208u^3 - 373u^2 + 164u - 46$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{25} - 21u^{24} + \cdots + 1792u - 256$
c_2, c_7, c_9	$u^{25} + u^{24} + \cdots + u + 1$
c_3, c_8, c_{12}	$u^{25} + 18u^{23} + \cdots + 20u^2 + 1$
c_4, c_5, c_{11}	$u^{25} - 5u^{24} + \cdots + 10u + 4$
c_6, c_{10}	$u^{25} + 15u^{24} + \cdots - 1986u - 196$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{25} - 13y^{24} + \cdots - 131072y - 65536$
c_2, c_7, c_9	$y^{25} - 15y^{24} + \cdots + 9y - 1$
c_3, c_8, c_{12}	$y^{25} + 36y^{24} + \cdots - 40y - 1$
c_4, c_5, c_{11}	$y^{25} - 21y^{24} + \cdots - 100y - 16$
c_6, c_{10}	$y^{25} + 15y^{24} + \cdots - 132996y - 38416$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.042442 + 0.868462I$ $a = -0.716248 + 0.009860I$ $b = -0.315314 - 0.333618I$	$6.65854 - 2.22451I$	$1.85427 + 3.24248I$
$u = 0.042442 - 0.868462I$ $a = -0.716248 - 0.009860I$ $b = -0.315314 + 0.333618I$	$6.65854 + 2.22451I$	$1.85427 - 3.24248I$
$u = 0.170420 + 0.851806I$ $a = -1.68479 - 1.05733I$ $b = -0.313941 - 1.151630I$	$1.36106 - 10.21960I$	$3.12273 + 6.18876I$
$u = 0.170420 - 0.851806I$ $a = -1.68479 + 1.05733I$ $b = -0.313941 + 1.151630I$	$1.36106 + 10.21960I$	$3.12273 - 6.18876I$
$u = 0.689863 + 0.516195I$ $a = 0.90028 + 1.54729I$ $b = 0.157173 + 0.858656I$	$-2.89781 - 4.73922I$	$0.64335 + 6.01647I$
$u = 0.689863 - 0.516195I$ $a = 0.90028 - 1.54729I$ $b = 0.157173 - 0.858656I$	$-2.89781 + 4.73922I$	$0.64335 - 6.01647I$
$u = 1.062620 + 0.439244I$ $a = -0.93017 - 1.06985I$ $b = -0.393102 - 0.665533I$	$-1.37637 + 5.59077I$	$0.49701 - 2.52369I$
$u = 1.062620 - 0.439244I$ $a = -0.93017 + 1.06985I$ $b = -0.393102 + 0.665533I$	$-1.37637 - 5.59077I$	$0.49701 + 2.52369I$
$u = 0.370519 + 0.720063I$ $a = 1.56474 + 0.89351I$ $b = 0.401573 + 0.852679I$	$-1.86125 + 0.38147I$	$3.19263 - 0.66430I$
$u = 0.370519 - 0.720063I$ $a = 1.56474 - 0.89351I$ $b = 0.401573 - 0.852679I$	$-1.86125 - 0.38147I$	$3.19263 + 0.66430I$

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.27648$		
$a = 0.500312$	-2.96177	-0.698220
$b = 0.280537$		
$u = -1.296580 + 0.136250I$		
$a = 0.004316 - 0.323358I$	$-4.47329 + 2.69307I$	$-5.03326 - 5.98784I$
$b = 0.429924 - 1.218120I$		
$u = -1.296580 - 0.136250I$		
$a = 0.004316 + 0.323358I$	$-4.47329 - 2.69307I$	$-5.03326 + 5.98784I$
$b = 0.429924 + 1.218120I$		
$u = 1.235760 + 0.416433I$		
$a = -0.139402 - 0.504528I$	$2.97397 - 2.37583I$	$-1.057681 + 0.915229I$
$b = -0.038519 - 0.297655I$		
$u = 1.235760 - 0.416433I$		
$a = -0.139402 + 0.504528I$	$2.97397 + 2.37583I$	$-1.057681 - 0.915229I$
$b = -0.038519 + 0.297655I$		
$u = -1.304980 + 0.396966I$		
$a = -0.149198 + 0.398993I$	$2.45389 + 6.76060I$	$-2.28268 - 6.77573I$
$b = -0.962816 + 0.648234I$		
$u = -1.304980 - 0.396966I$		
$a = -0.149198 - 0.398993I$	$2.45389 - 6.76060I$	$-2.28268 + 6.77573I$
$b = -0.962816 - 0.648234I$		
$u = -1.37760 + 0.36657I$		
$a = 0.142768 + 1.259210I$	$-3.5266 + 14.6054I$	$-1.02784 - 7.76061I$
$b = -1.32923 + 3.16497I$		
$u = -1.37760 - 0.36657I$		
$a = 0.142768 - 1.259210I$	$-3.5266 - 14.6054I$	$-1.02784 + 7.76061I$
$b = -1.32923 - 3.16497I$		
$u = -1.44314 + 0.25974I$		
$a = 0.006895 - 1.095370I$	$-7.68655 + 3.14182I$	$0.46512 + 1.43904I$
$b = 1.09681 - 2.92836I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.44314 - 0.25974I$		
$a = 0.006895 + 1.095370I$	$-7.68655 - 3.14182I$	$0.46512 - 1.43904I$
$b = 1.09681 + 2.92836I$		
$u = -1.46408 + 0.08648I$		
$a = -0.108140 - 1.228800I$	$-9.92989 + 6.47027I$	$-4.20740 - 5.48144I$
$b = 0.14349 - 3.82971I$		
$u = -1.46408 - 0.08648I$		
$a = -0.108140 + 1.228800I$	$-9.92989 - 6.47027I$	$-4.20740 + 5.48144I$
$b = 0.14349 + 3.82971I$		
$u = 0.176540 + 0.371053I$		
$a = 0.608793 + 0.633615I$	$0.045964 - 0.832254I$	$1.18286 + 8.33998I$
$b = -0.016319 + 0.292219I$		
$u = 0.176540 - 0.371053I$		
$a = 0.608793 - 0.633615I$	$0.045964 + 0.832254I$	$1.18286 - 8.33998I$
$b = -0.016319 - 0.292219I$		

$$\text{II. } I_2^u = \langle 20u^7a^3 - 10u^7a^2 + \cdots - 62a + 47, -3u^7a^2 + 6u^7a + \cdots - 20a - 11, u^8 - u^7 - 3u^6 + 2u^5 + 3u^4 - 2u - 1 \rangle$$

(i) **Arc colorings**

$$a_5 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} a \\ -0.465116a^3u^7 + 0.232558a^2u^7 + \cdots + 1.44186a - 1.09302 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} a^2u \\ -0.790698a^3u^7 + 1.39535a^2u^7 + \cdots + 0.651163a - 0.558140 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} a \\ -0.465116a^3u^7 + 0.232558a^2u^7 + \cdots + 1.44186a - 1.09302 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} u \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -u^2 + 1 \\ -u^2 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 0.116279a^3u^7 + 0.441860a^2u^7 + \cdots + 0.139535a + 1.02326 \\ -0.302326a^3u^7 + 1.65116a^2u^7 + \cdots + 0.837209a + 0.139535 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} 0.186047a^3u^7 - 0.0930233a^2u^7 + \cdots + 0.0232558a + 0.837209 \\ 1.90698a^3u^7 - 1.95349a^2u^7 + \cdots - 0.511628a - 0.418605 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} u^6 - 3u^4 + 2u^2 + 1 \\ u^7 + u^6 - 2u^5 - 3u^4 + 2u^2 + 2u + 1 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -u^3 + 2u \\ -u^5 + u^3 + u \end{pmatrix}$$

(ii) **Obstruction class** = -1

(iii) **Cusp Shapes** = $-4u^6 + 12u^4 + 4u^3 - 8u^2 - 8u - 6$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$(u^2 + u - 1)^{16}$
c_2, c_7, c_9	$u^{32} + u^{31} + \dots + 318u + 199$
c_3, c_8, c_{12}	$u^{32} - u^{31} + \dots - 1792u - 271$
c_4, c_5, c_{11}	$(u^8 + u^7 - 3u^6 - 2u^5 + 3u^4 + 2u - 1)^4$
c_6, c_{10}	$(u^8 - 3u^7 + 7u^6 - 10u^5 + 11u^4 - 10u^3 + 6u^2 - 4u + 1)^4$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$(y^2 - 3y + 1)^{16}$
c_2, c_7, c_9	$y^{32} - 13y^{31} + \cdots - 651956y + 39601$
c_3, c_8, c_{12}	$y^{32} + 19y^{31} + \cdots - 1289332y + 73441$
c_4, c_5, c_{11}	$(y^8 - 7y^7 + 19y^6 - 22y^5 + 3y^4 + 14y^3 - 6y^2 - 4y + 1)^4$
c_6, c_{10}	$(y^8 + 5y^7 + 11y^6 + 6y^5 - 17y^4 - 34y^3 - 22y^2 - 4y + 1)^4$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.180120 + 0.268597I$		
$a = -0.573305 + 0.724047I$	$-4.98850 + 1.13123I$	$0.584775 - 0.510791I$
$b = 0.069236 - 0.301245I$		
$u = -1.180120 + 0.268597I$		
$a = 0.977884 + 0.579436I$	$2.90719 + 1.13123I$	$0.584775 - 0.510791I$
$b = -0.33020 + 2.17451I$		
$u = -1.180120 + 0.268597I$		
$a = 0.371867 - 1.152550I$	$-4.98850 + 1.13123I$	$0.584775 - 0.510791I$
$b = 0.48288 - 1.38255I$		
$u = -1.180120 + 0.268597I$		
$a = -0.450513 + 0.542402I$	$2.90719 + 1.13123I$	$0.584775 - 0.510791I$
$b = -1.11527 + 2.23372I$		
$u = -1.180120 - 0.268597I$		
$a = -0.573305 - 0.724047I$	$-4.98850 - 1.13123I$	$0.584775 + 0.510791I$
$b = 0.069236 + 0.301245I$		
$u = -1.180120 - 0.268597I$		
$a = 0.977884 - 0.579436I$	$2.90719 - 1.13123I$	$0.584775 + 0.510791I$
$b = -0.33020 - 2.17451I$		
$u = -1.180120 - 0.268597I$		
$a = 0.371867 + 1.152550I$	$-4.98850 - 1.13123I$	$0.584775 + 0.510791I$
$b = 0.48288 + 1.38255I$		
$u = -1.180120 - 0.268597I$		
$a = -0.450513 - 0.542402I$	$2.90719 - 1.13123I$	$0.584775 + 0.510791I$
$b = -1.11527 - 2.23372I$		
$u = -0.108090 + 0.747508I$		
$a = -1.029300 + 0.246196I$	$6.10726 + 2.57849I$	$3.72292 - 3.56796I$
$b = 0.524935 - 0.257181I$		
$u = -0.108090 + 0.747508I$		
$a = -1.46947 + 1.05203I$	$-1.78843 + 2.57849I$	$3.72292 - 3.56796I$
$b = -0.428409 + 1.312810I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.108090 + 0.747508I$		
$a = -0.64475 - 1.85535I$	$6.10726 + 2.57849I$	$3.72292 - 3.56796I$
$b = -0.208281 - 1.216220I$		
$u = -0.108090 + 0.747508I$		
$a = 2.10891 - 0.43739I$	$-1.78843 + 2.57849I$	$3.72292 - 3.56796I$
$b = 0.307458 - 0.750018I$		
$u = -0.108090 - 0.747508I$		
$a = -1.029300 - 0.246196I$	$6.10726 - 2.57849I$	$3.72292 + 3.56796I$
$b = 0.524935 + 0.257181I$		
$u = -0.108090 - 0.747508I$		
$a = -1.46947 - 1.05203I$	$-1.78843 - 2.57849I$	$3.72292 + 3.56796I$
$b = -0.428409 - 1.312810I$		
$u = -0.108090 - 0.747508I$		
$a = -0.64475 + 1.85535I$	$6.10726 - 2.57849I$	$3.72292 + 3.56796I$
$b = -0.208281 + 1.216220I$		
$u = -0.108090 - 0.747508I$		
$a = 2.10891 + 0.43739I$	$-1.78843 - 2.57849I$	$3.72292 + 3.56796I$
$b = 0.307458 + 0.750018I$		
$u = 1.37100$		
$a = 0.975338$	-2.55489	-5.86400
$b = 1.58426$		
$u = 1.37100$		
$a = -0.247524 + 1.200480I$	-10.4506	-5.86400
$b = -0.14313 + 4.43627I$		
$u = 1.37100$		
$a = -0.247524 - 1.200480I$	-10.4506	-5.86400
$b = -0.14313 - 4.43627I$		
$u = 1.37100$		
$a = 0.320715$	-2.55489	-5.86400
$b = -0.834836$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.334530 + 0.318930I$	$-6.32752 - 6.44354I$	$-1.42845 + 5.29417I$
$a = 0.283327 - 1.065010I$		
$b = -1.51898 - 2.93006I$		
$u = 1.334530 + 0.318930I$	$1.56816 - 6.44354I$	$-1.42845 + 5.29417I$
$a = -1.130370 + 0.118813I$		
$b = -1.00572 + 1.40159I$		
$u = 1.334530 + 0.318930I$	$-6.32752 - 6.44354I$	$-1.42845 + 5.29417I$
$a = 0.237323 + 1.244130I$		
$b = 1.81040 + 3.06377I$		
$u = 1.334530 + 0.318930I$	$1.56816 - 6.44354I$	$-1.42845 + 5.29417I$
$a = -0.232706 - 0.587777I$		
$b = 0.24276 - 1.75166I$		
$u = 1.334530 - 0.318930I$	$-6.32752 + 6.44354I$	$-1.42845 - 5.29417I$
$a = 0.283327 + 1.065010I$		
$b = -1.51898 + 2.93006I$		
$u = 1.334530 - 0.318930I$	$1.56816 + 6.44354I$	$-1.42845 - 5.29417I$
$a = -1.130370 - 0.118813I$		
$b = -1.00572 - 1.40159I$		
$u = 1.334530 - 0.318930I$	$-6.32752 + 6.44354I$	$-1.42845 - 5.29417I$
$a = 0.237323 - 1.244130I$		
$b = 1.81040 - 3.06377I$		
$u = 1.334530 - 0.318930I$	$1.56816 + 6.44354I$	$-1.42845 - 5.29417I$
$a = -0.232706 + 0.587777I$		
$b = 0.24276 + 1.75166I$		
$u = -0.463640$		
$a = -0.460586$	3.10281	-3.89450
$b = -1.38936$		
$u = -0.463640$		
$a = 2.56602$	3.10281	-3.89450
$b = -0.430619$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.463640$		
$a = -0.40210 + 3.00975I$	-4.79288	-3.89450
$b = 0.347586 + 0.953404I$		
$u = -0.463640$		
$a = -0.40210 - 3.00975I$	-4.79288	-3.89450
$b = 0.347586 - 0.953404I$		

III.

$$I_3^u = \langle u^{14} - u^{13} + \dots + b - 3u, \ u^{14} - 6u^{12} + \dots + a - 3, \ u^{15} - 7u^{13} + \dots - 3u - 1 \rangle$$

(i) **Arc colorings**

$$a_5 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -u^{14} + 6u^{12} - 13u^{10} + 9u^8 + 7u^6 - 11u^4 + u^3 - u + 3 \\ -u^{14} + u^{13} + \dots - 4u^2 + 3u \end{pmatrix}$$

$$a_1 = \begin{pmatrix} u^{11} - 5u^9 + 9u^7 - 5u^5 - 3u^3 + u^2 + 3u - 2 \\ -u^{14} + u^{13} + \dots + u + 1 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} -u^{14} + 6u^{12} - 13u^{10} + 9u^8 + 7u^6 - 11u^4 + u^3 - u + 3 \\ -2u^{14} + u^{13} + \dots - 4u^2 + 2u \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} u \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} -u^2 + 1 \\ -u^2 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} u^{14} - 6u^{12} + \dots + 4u - 2 \\ u^{13} + u^{12} + \dots + u + 1 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} u^{11} - 5u^9 + 9u^7 - 5u^5 - 3u^3 + 3u - 1 \\ -u^{14} + u^{13} + \dots + 6u^3 + 1 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} u^6 - 3u^4 + 2u^2 + 1 \\ u^8 - 2u^6 + 2u^2 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} -u^3 + 2u \\ -u^5 + u^3 + u \end{pmatrix}$$

(ii) **Obstruction class = 1**

(iii) **Cusp Shapes** = $-4u^{14} + 2u^{13} + 28u^{12} - 10u^{11} - 72u^{10} + 16u^9 + 72u^8 - 2u^7 + 5u^6 - 16u^5 - 46u^4 + 13u^3 + 3u^2 - 5u + 10$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{15} - 6u^{14} + \cdots + 7u - 1$
c_2, c_9	$u^{15} + u^{14} + \cdots + 3u^2 + 1$
c_3, c_8	$u^{15} + 3u^{13} + \cdots + u + 1$
c_4, c_5	$u^{15} - 7u^{13} + 19u^{11} - 22u^9 + 3u^7 + 14u^5 - u^4 - 6u^3 + 2u^2 - 3u - 1$
c_6	$u^{15} + 5u^{13} + \cdots - 3u - 1$
c_7	$u^{15} - u^{14} + \cdots - 3u^2 - 1$
c_{10}	$u^{15} + 5u^{13} + \cdots - 3u + 1$
c_{11}	$u^{15} - 7u^{13} + 19u^{11} - 22u^9 + 3u^7 + 14u^5 + u^4 - 6u^3 - 2u^2 - 3u + 1$
c_{12}	$u^{15} + 3u^{13} + \cdots + u - 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{15} - 10y^{14} + \cdots + 65y - 1$
c_2, c_7, c_9	$y^{15} - 9y^{14} + \cdots - 6y - 1$
c_3, c_8, c_{12}	$y^{15} + 6y^{14} + \cdots + 9y - 1$
c_4, c_5, c_{11}	$y^{15} - 14y^{14} + \cdots + 13y - 1$
c_6, c_{10}	$y^{15} + 10y^{14} + \cdots + 15y - 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.067646 + 0.825365I$		
$a = -0.380175 - 0.868214I$	$7.99475 + 1.92226I$	$9.12294 - 1.22401I$
$b = 0.310598 - 0.548513I$		
$u = -0.067646 - 0.825365I$		
$a = -0.380175 + 0.868214I$	$7.99475 - 1.92226I$	$9.12294 + 1.22401I$
$b = 0.310598 + 0.548513I$		
$u = -1.20404$		
$a = -0.582456$	0.894455	-3.21180
$b = -3.43713$		
$u = 1.202120 + 0.181369I$		
$a = 0.277876 + 0.914400I$	$-6.17070 - 1.65898I$	$-7.52412 + 4.06635I$
$b = 0.118572 + 0.512284I$		
$u = 1.202120 - 0.181369I$		
$a = 0.277876 - 0.914400I$	$-6.17070 + 1.65898I$	$-7.52412 - 4.06635I$
$b = 0.118572 - 0.512284I$		
$u = -1.216500 + 0.366926I$		
$a = 0.493698 + 0.310572I$	4.46605 + 2.37005I	5.16588 - 2.66961I
$b = 0.45473 + 1.67906I$		
$u = -1.216500 - 0.366926I$		
$a = 0.493698 - 0.310572I$	4.46605 - 2.37005I	5.16588 + 2.66961I
$b = 0.45473 - 1.67906I$		
$u = 1.322630 + 0.369202I$		
$a = -0.597920 - 0.072527I$	3.63687 - 6.22447I	4.70740 + 3.89231I
$b = -0.341991 - 0.081983I$		
$u = 1.322630 - 0.369202I$		
$a = -0.597920 + 0.072527I$	3.63687 + 6.22447I	4.70740 - 3.89231I
$b = -0.341991 + 0.081983I$		
$u = 1.38580$		
$a = 0.772897$	-1.59422	4.26710
$b = 0.448939$		

Solutions to I_3^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.191847 + 0.580989I$	$-3.17279 - 1.01583I$	$0.86576 + 1.51873I$
$a = 2.04075 + 1.02550I$		
$b = 0.318346 + 1.004690I$		
$u = 0.191847 - 0.580989I$	$-3.17279 + 1.01583I$	$0.86576 - 1.51873I$
$a = 2.04075 - 1.02550I$		
$b = 0.318346 - 1.004690I$		
$u = -1.393270 + 0.230580I$	$-8.28883 + 4.00739I$	$-4.40865 - 4.46883I$
$a = -0.026700 - 1.147080I$		
$b = 1.19545 - 3.34732I$		
$u = -1.393270 - 0.230580I$	$-8.28883 - 4.00739I$	$-4.40865 + 4.46883I$
$a = -0.026700 + 1.147080I$		
$b = 1.19545 + 3.34732I$		
$u = -0.260139$	3.76907	11.0860
$a = 3.19450$		
$b = -1.12320$		

IV. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1	$((u^2 + u - 1)^{16})(u^{15} - 6u^{14} + \dots + 7u - 1)$ $\cdot (u^{25} - 21u^{24} + \dots + 1792u - 256)$
c_2, c_9	$(u^{15} + u^{14} + \dots + 3u^2 + 1)(u^{25} + u^{24} + \dots + u + 1)$ $\cdot (u^{32} + u^{31} + \dots + 318u + 199)$
c_3, c_8	$(u^{15} + 3u^{13} + \dots + u + 1)(u^{25} + 18u^{23} + \dots + 20u^2 + 1)$ $\cdot (u^{32} - u^{31} + \dots - 1792u - 271)$
c_4, c_5	$(u^8 + u^7 - 3u^6 - 2u^5 + 3u^4 + 2u - 1)^4$ $\cdot (u^{15} - 7u^{13} + 19u^{11} - 22u^9 + 3u^7 + 14u^5 - u^4 - 6u^3 + 2u^2 - 3u - 1)$ $\cdot (u^{25} - 5u^{24} + \dots + 10u + 4)$
c_6	$(u^8 - 3u^7 + 7u^6 - 10u^5 + 11u^4 - 10u^3 + 6u^2 - 4u + 1)^4$ $\cdot (u^{15} + 5u^{13} + \dots - 3u - 1)(u^{25} + 15u^{24} + \dots - 1986u - 196)$
c_7	$(u^{15} - u^{14} + \dots - 3u^2 - 1)(u^{25} + u^{24} + \dots + u + 1)$ $\cdot (u^{32} + u^{31} + \dots + 318u + 199)$
c_{10}	$(u^8 - 3u^7 + 7u^6 - 10u^5 + 11u^4 - 10u^3 + 6u^2 - 4u + 1)^4$ $\cdot (u^{15} + 5u^{13} + \dots - 3u + 1)(u^{25} + 15u^{24} + \dots - 1986u - 196)$
c_{11}	$(u^8 + u^7 - 3u^6 - 2u^5 + 3u^4 + 2u - 1)^4$ $\cdot (u^{15} - 7u^{13} + 19u^{11} - 22u^9 + 3u^7 + 14u^5 + u^4 - 6u^3 - 2u^2 - 3u + 1)$ $\cdot (u^{25} - 5u^{24} + \dots + 10u + 4)$
c_{12}	$(u^{15} + 3u^{13} + \dots + u - 1)(u^{25} + 18u^{23} + \dots + 20u^2 + 1)$ $\cdot (u^{32} - u^{31} + \dots - 1792u - 271)$

V. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1	$((y^2 - 3y + 1)^{16})(y^{15} - 10y^{14} + \dots + 65y - 1)$ $\cdot (y^{25} - 13y^{24} + \dots - 131072y - 65536)$
c_2, c_7, c_9	$(y^{15} - 9y^{14} + \dots - 6y - 1)(y^{25} - 15y^{24} + \dots + 9y - 1)$ $\cdot (y^{32} - 13y^{31} + \dots - 651956y + 39601)$
c_3, c_8, c_{12}	$(y^{15} + 6y^{14} + \dots + 9y - 1)(y^{25} + 36y^{24} + \dots - 40y - 1)$ $\cdot (y^{32} + 19y^{31} + \dots - 1289332y + 73441)$
c_4, c_5, c_{11}	$(y^8 - 7y^7 + 19y^6 - 22y^5 + 3y^4 + 14y^3 - 6y^2 - 4y + 1)^4$ $\cdot (y^{15} - 14y^{14} + \dots + 13y - 1)(y^{25} - 21y^{24} + \dots - 100y - 16)$
c_6, c_{10}	$(y^8 + 5y^7 + 11y^6 + 6y^5 - 17y^4 - 34y^3 - 22y^2 - 4y + 1)^4$ $\cdot (y^{15} + 10y^{14} + \dots + 15y - 1)(y^{25} + 15y^{24} + \dots - 132996y - 38416)$