


## $11a_{45}$ ( $K11a_{45}$ )



### Ideals for irreducible components<sup>2</sup> of $X_{\text{par}}$

$$I_1^u = \langle u^{43} + u^{42} + \dots + b + 1, -u^{46} - 2u^{45} + \dots + a - 2, u^{47} + 2u^{46} + \dots + 4u + 1 \rangle$$

$$I_2^u = \langle b + 1, -u^3 - u^2 + a - 3u, u^4 + u^3 + 3u^2 + 2u + 1 \rangle$$

\* 2 irreducible components of  $\dim_{\mathbb{C}} = 0$ , with total 51 representations.

---

<sup>1</sup>The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/math/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

<sup>2</sup>All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$I_1^u = \langle u^{43} + u^{42} + \cdots + b + 1, -u^{46} - 2u^{45} + \cdots + a - 2, u^{47} + 2u^{46} + \cdots + 4u + 1 \rangle$$

(i) Arc colorings

$$a_5 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 1 \\ -u^2 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} u^{46} + 2u^{45} + \cdots + u + 2 \\ -u^{43} - u^{42} + \cdots - 9u^3 - 1 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} u \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u^{46} + 2u^{45} + \cdots - u + 2 \\ -u^{43} - u^{42} + \cdots - u - 1 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} u^9 + 4u^7 + 3u^5 - 2u^3 + u \\ u^9 + 5u^7 + 7u^5 + 2u^3 + u \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -u \\ u^3 + u \end{pmatrix}$$

$$a_7 = \begin{pmatrix} u^2 + 1 \\ -u^4 - 2u^2 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} u^{46} + 2u^{45} + \cdots + 3u + 3 \\ u^{44} - u^{43} + \cdots + u^2 - 1 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} u^5 + 2u^3 - u \\ u^5 + 3u^3 + u \end{pmatrix}$$

$$a_8 = \begin{pmatrix} u^5 + 2u^3 - u \\ u^5 + 3u^3 + u \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes =  $u^{46} + 2u^{45} + \cdots - 7u - 9$

**(iv) u-Polynomials at the component**

Crossings	u-Polynomials at each crossing
$c_1, c_4$	$u^{47} - 5u^{46} + \cdots - 8u + 1$
$c_2$	$u^{47} + 21u^{46} + \cdots - 6u + 1$
$c_3, c_7$	$u^{47} + u^{46} + \cdots + 40u + 16$
$c_5, c_6, c_9$ $c_{10}$	$u^{47} + 2u^{46} + \cdots + 4u + 1$
$c_8, c_{11}$	$u^{47} - 8u^{46} + \cdots + 616u - 49$

**(v) Riley Polynomials at the component**

Crossings	Riley Polynomials at each crossing
$c_1, c_4$	$y^{47} - 21y^{46} + \cdots - 6y - 1$
$c_2$	$y^{47} + 15y^{46} + \cdots - 262y - 1$
$c_3, c_7$	$y^{47} + 27y^{46} + \cdots - 3264y - 256$
$c_5, c_6, c_9$ $c_{10}$	$y^{47} + 52y^{46} + \cdots + 12y - 1$
$c_8, c_{11}$	$y^{47} + 32y^{46} + \cdots + 287140y - 2401$

(vi) Complex Volumes and Cusp Shapes

Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.604283 + 0.592983I$		
$a = 1.47187 - 1.86934I$	$4.37990 - 10.31740I$	$-1.65034 + 8.68510I$
$b = 1.126700 + 0.685229I$		
$u = -0.604283 - 0.592983I$		
$a = 1.47187 + 1.86934I$	$4.37990 + 10.31740I$	$-1.65034 - 8.68510I$
$b = 1.126700 - 0.685229I$		
$u = 0.232115 + 0.798027I$		
$a = 2.24071 + 0.38222I$	$-1.04927 + 4.94975I$	$-5.93011 - 7.21371I$
$b = 0.974632 - 0.559034I$		
$u = 0.232115 - 0.798027I$		
$a = 2.24071 - 0.38222I$	$-1.04927 - 4.94975I$	$-5.93011 + 7.21371I$
$b = 0.974632 + 0.559034I$		
$u = -0.610300 + 0.553965I$		
$a = -0.902531 - 0.046325I$	$6.31726 - 4.42879I$	$1.23708 + 4.26556I$
$b = 0.490595 - 0.918525I$		
$u = -0.610300 - 0.553965I$		
$a = -0.902531 + 0.046325I$	$6.31726 + 4.42879I$	$1.23708 - 4.26556I$
$b = 0.490595 + 0.918525I$		
$u = 0.564268 + 0.532900I$		
$a = -0.60777 - 2.13982I$	$1.15387 + 4.21075I$	$-2.42283 - 6.46620I$
$b = -0.886230 + 0.571745I$		
$u = 0.564268 - 0.532900I$		
$a = -0.60777 + 2.13982I$	$1.15387 - 4.21075I$	$-2.42283 + 6.46620I$
$b = -0.886230 - 0.571745I$		
$u = 0.368403 + 0.677743I$		
$a = 0.723270 + 1.043890I$	$-0.121886 + 0.686906I$	$-2.62923 - 1.30814I$
$b = 0.725690 + 0.454437I$		
$u = 0.368403 - 0.677743I$		
$a = 0.723270 - 1.043890I$	$-0.121886 - 0.686906I$	$-2.62923 + 1.30814I$
$b = 0.725690 - 0.454437I$		

Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.631315 + 0.431371I$		
$a = 0.384246 - 0.958542I$	$6.67942 + 0.21804I$	$2.26501 + 2.20975I$
$b = 0.543018 + 0.899712I$		
$u = -0.631315 - 0.431371I$		
$a = 0.384246 + 0.958542I$	$6.67942 - 0.21804I$	$2.26501 - 2.20975I$
$b = 0.543018 - 0.899712I$		
$u = -0.641272 + 0.383119I$		
$a = 0.022026 + 0.485329I$	$4.99878 + 6.09831I$	$0.10454 - 2.74650I$
$b = 1.094550 - 0.696965I$		
$u = -0.641272 - 0.383119I$		
$a = 0.022026 - 0.485329I$	$4.99878 - 6.09831I$	$0.10454 + 2.74650I$
$b = 1.094550 + 0.696965I$		
$u = -0.536843 + 0.494469I$		
$a = -0.97848 + 1.17374I$	$-0.20145 - 1.85701I$	$-0.50975 + 4.37782I$
$b = -1.286310 + 0.037305I$		
$u = -0.536843 - 0.494469I$		
$a = -0.97848 - 1.17374I$	$-0.20145 + 1.85701I$	$-0.50975 - 4.37782I$
$b = -1.286310 - 0.037305I$		
$u = 0.565661 + 0.444526I$		
$a = 0.741523 + 0.717629I$	$1.41432 - 0.32518I$	$-1.284718 - 0.498489I$
$b = -0.806231 - 0.561068I$		
$u = 0.565661 - 0.444526I$		
$a = 0.741523 - 0.717629I$	$1.41432 + 0.32518I$	$-1.284718 + 0.498489I$
$b = -0.806231 + 0.561068I$		
$u = -0.073145 + 0.598544I$		
$a = -2.60222 + 1.37444I$	$-2.92920 - 0.94552I$	$-11.86909 + 0.58583I$
$b = -1.034680 - 0.259393I$		
$u = -0.073145 - 0.598544I$		
$a = -2.60222 - 1.37444I$	$-2.92920 + 0.94552I$	$-11.86909 - 0.58583I$
$b = -1.034680 + 0.259393I$		

Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.284667 + 0.487292I$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	
$a = 0.588067 + 0.203086I$	$-0.038903 + 1.106530I$	$-0.72417 - 6.07516I$
$b = 0.141384 + 0.220781I$		
$u = 0.284667 - 0.487292I$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	
$a = 0.588067 - 0.203086I$	$-0.038903 - 1.106530I$	$-0.72417 + 6.07516I$
$b = 0.141384 - 0.220781I$		
$u = -0.15656 + 1.43102I$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	
$a = 0.964258 - 0.136228I$	$-0.77593 + 3.28450I$	0
$b = 1.042580 - 0.719852I$		
$u = -0.15656 - 1.43102I$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	
$a = 0.964258 + 0.136228I$	$-0.77593 - 3.28450I$	0
$b = 1.042580 + 0.719852I$		
$u = 0.534809 + 0.072952I$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	
$a = 0.267834 + 0.634733I$	$1.73762 + 2.33285I$	$1.63431 - 3.88919I$
$b = 0.841064 - 0.586379I$		
$u = 0.534809 - 0.072952I$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	
$a = 0.267834 - 0.634733I$	$1.73762 - 2.33285I$	$1.63431 + 3.88919I$
$b = 0.841064 + 0.586379I$		
$u = -0.17342 + 1.46958I$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	
$a = 0.892690 - 0.159044I$	$0.53093 - 2.62820I$	0
$b = 0.614156 + 0.885412I$		
$u = -0.17342 - 1.46958I$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	
$a = 0.892690 + 0.159044I$	$0.53093 + 2.62820I$	0
$b = 0.614156 - 0.885412I$		
$u = 0.14450 + 1.50474I$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	
$a = 0.086605 + 0.128202I$	$-4.99092 + 2.12497I$	0
$b = -0.704621 - 0.592796I$		
$u = 0.14450 - 1.50474I$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	
$a = 0.086605 - 0.128202I$	$-4.99092 - 2.12497I$	0
$b = -0.704621 + 0.592796I$		

Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.14983 + 1.52897I$	$-6.93429 - 4.28271I$	0
$a = -2.18209 + 0.88257I$		
$b = -1.313780 + 0.094691I$		
$u = -0.14983 - 1.52897I$	$-6.93429 + 4.28271I$	0
$a = -2.18209 - 0.88257I$		
$b = -1.313780 - 0.094691I$		
$u = 0.04949 + 1.54079I$	$-6.95405 + 2.13733I$	0
$a = 0.359833 + 0.420302I$		
$b = 0.022840 + 0.536346I$		
$u = 0.04949 - 1.54079I$	$-6.95405 - 2.13733I$	0
$a = 0.359833 - 0.420302I$		
$b = 0.022840 - 0.536346I$		
$u = 0.16493 + 1.53761I$	$-5.72542 + 6.83119I$	0
$a = -1.47288 - 1.40203I$		
$b = -0.945052 + 0.589157I$		
$u = 0.16493 - 1.53761I$	$-5.72542 - 6.83119I$	0
$a = -1.47288 + 1.40203I$		
$b = -0.945052 - 0.589157I$		
$u = -0.18579 + 1.54140I$	$-0.62096 - 7.31850I$	0
$a = -0.364549 - 0.727740I$		
$b = 0.443402 - 0.937849I$		
$u = -0.18579 - 1.54140I$	$-0.62096 + 7.31850I$	0
$a = -0.364549 + 0.727740I$		
$b = 0.443402 + 0.937849I$		
$u = -0.01255 + 1.55908I$	$-10.25720 - 1.20723I$	0
$a = -2.70333 + 0.61165I$		
$b = -1.133330 - 0.334796I$		
$u = -0.01255 - 1.55908I$	$-10.25720 + 1.20723I$	0
$a = -2.70333 - 0.61165I$		
$b = -1.133330 + 0.334796I$		

Solutions to $I_1^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.18591 + 1.55945I$		
$a = 2.26208 - 1.08076I$	$-2.78197 - 13.21230I$	0
$b = 1.152470 + 0.673628I$		
$u = -0.18591 - 1.55945I$		
$a = 2.26208 + 1.08076I$	$-2.78197 + 13.21230I$	0
$b = 1.152470 - 0.673628I$		
$u = 0.10845 + 1.58073I$		
$a = 1.54084 + 0.82842I$	$-7.73076 + 2.45052I$	0
$b = 0.782367 + 0.305537I$		
$u = 0.10845 - 1.58073I$		
$a = 1.54084 - 0.82842I$	$-7.73076 - 2.45052I$	0
$b = 0.782367 - 0.305537I$		
$u = 0.04923 + 1.60296I$		
$a = 2.55612 - 0.07351I$	$-9.19770 + 5.90087I$	0
$b = 1.041330 - 0.500879I$		
$u = 0.04923 - 1.60296I$		
$a = 2.55612 + 0.07351I$	$-9.19770 - 5.90087I$	0
$b = 1.041330 + 0.500879I$		
$u = -0.210582$		
$a = 2.42377$	$-1.24674$	$-7.85810$
$b = -0.853085$		

$$\text{II. } I_2^u = \langle b + 1, -u^3 - u^2 + a - 3u, u^4 + u^3 + 3u^2 + 2u + 1 \rangle$$

(i) Arc colorings

$$a_5 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 1 \\ -u^2 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} u^3 + u^2 + 3u \\ -1 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} u \\ u \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u^3 + u^2 + 3u + 1 \\ -1 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} -u \\ u^3 + u \end{pmatrix}$$

$$a_7 = \begin{pmatrix} u^2 + 1 \\ u^3 + u^2 + 2u + 1 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} u^3 + u^2 + 3u + 1 \\ -1 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} u^2 + 1 \\ u^3 + u^2 + 2u + 1 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} u^2 + 1 \\ u^3 + u^2 + 2u + 1 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes =  $3u^3 + 3u^2 + 10u - 4$

**(iv) u-Polynomials at the component**

Crossings	u-Polynomials at each crossing
$c_1$	$(u - 1)^4$
$c_2, c_4$	$(u + 1)^4$
$c_3, c_7$	$u^4$
$c_5, c_6$	$u^4 + u^3 + 3u^2 + 2u + 1$
$c_8$	$u^4 + u^3 + u^2 + 1$
$c_9, c_{10}$	$u^4 - u^3 + 3u^2 - 2u + 1$
$c_{11}$	$u^4 - u^3 + u^2 + 1$

**(v) Riley Polynomials at the component**

Crossings	Riley Polynomials at each crossing
$c_1, c_2, c_4$	$(y - 1)^4$
$c_3, c_7$	$y^4$
$c_5, c_6, c_9$ $c_{10}$	$y^4 + 5y^3 + 7y^2 + 2y + 1$
$c_8, c_{11}$	$y^4 + y^3 + 3y^2 + 2y + 1$

**(vi) Complex Volumes and Cusp Shapes**

Solutions to $I_2^u$	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.395123 + 0.506844I$		
$a = -1.04332 + 1.22719I$	$-1.43393 - 1.41510I$	$-7.52507 + 4.18840I$
$b = -1.00000$		
$u = -0.395123 - 0.506844I$		
$a = -1.04332 - 1.22719I$	$-1.43393 + 1.41510I$	$-7.52507 - 4.18840I$
$b = -1.00000$		
$u = -0.10488 + 1.55249I$		
$a = -1.95668 + 0.64120I$	$-8.43568 - 3.16396I$	$-9.97493 + 3.47609I$
$b = -1.00000$		
$u = -0.10488 - 1.55249I$		
$a = -1.95668 - 0.64120I$	$-8.43568 + 3.16396I$	$-9.97493 - 3.47609I$
$b = -1.00000$		

### III. u-Polynomials

Crossings	u-Polynomials at each crossing
$c_1$	$((u - 1)^4)(u^{47} - 5u^{46} + \cdots - 8u + 1)$
$c_2$	$((u + 1)^4)(u^{47} + 21u^{46} + \cdots - 6u + 1)$
$c_3, c_7$	$u^4(u^{47} + u^{46} + \cdots + 40u + 16)$
$c_4$	$((u + 1)^4)(u^{47} - 5u^{46} + \cdots - 8u + 1)$
$c_5, c_6$	$(u^4 + u^3 + 3u^2 + 2u + 1)(u^{47} + 2u^{46} + \cdots + 4u + 1)$
$c_8$	$(u^4 + u^3 + u^2 + 1)(u^{47} - 8u^{46} + \cdots + 616u - 49)$
$c_9, c_{10}$	$(u^4 - u^3 + 3u^2 - 2u + 1)(u^{47} + 2u^{46} + \cdots + 4u + 1)$
$c_{11}$	$(u^4 - u^3 + u^2 + 1)(u^{47} - 8u^{46} + \cdots + 616u - 49)$

#### IV. Riley Polynomials

Crossings	Riley Polynomials at each crossing
$c_1, c_4$	$((y - 1)^4)(y^{47} - 21y^{46} + \dots - 6y - 1)$
$c_2$	$((y - 1)^4)(y^{47} + 15y^{46} + \dots - 262y - 1)$
$c_3, c_7$	$y^4(y^{47} + 27y^{46} + \dots - 3264y - 256)$
$c_5, c_6, c_9$ $c_{10}$	$(y^4 + 5y^3 + 7y^2 + 2y + 1)(y^{47} + 52y^{46} + \dots + 12y - 1)$
$c_8, c_{11}$	$(y^4 + y^3 + 3y^2 + 2y + 1)(y^{47} + 32y^{46} + \dots + 287140y - 2401)$