


9₁₃ ($K9a_{34}$)



Ideals for irreducible components² of X_{par}

$$I_1^u = \langle u^{18} - u^{17} + \cdots + 3u - 1 \rangle$$

* 1 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 18 representations.

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/math/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$\text{I. } I_1^u = \langle u^{18} - u^{17} + 9u^{16} - 8u^{15} + 32u^{14} - 25u^{13} + 55u^{12} - 36u^{11} + 43u^{10} - 19u^9 + 9u^8 + 4u^7 + 2u^5 + 4u^4 - 2u^3 - u^2 + 3u - 1 \rangle$$

(i) **Arc colorings**

$$a_5 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_4 = \begin{pmatrix} u \\ u^3 + u \end{pmatrix}$$

$$a_1 = \begin{pmatrix} u^2 + 1 \\ u^4 + 2u^2 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} u^3 + 2u \\ u^3 + u \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -u^7 - 4u^5 - 4u^3 \\ -u^7 - 3u^5 - 2u^3 + u \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -u^6 - 3u^4 - 2u^2 + 1 \\ -u^8 - 4u^6 - 4u^4 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -u^{17} - 8u^{15} - 25u^{13} - 36u^{11} - 19u^9 + 4u^7 + 2u^5 - 2u^3 + 3u \\ -u^{17} + u^{16} + \dots + 3u - 1 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -u^{17} - 8u^{15} - 25u^{13} - 36u^{11} - 19u^9 + 4u^7 + 2u^5 - 2u^3 + 3u \\ -u^{17} + u^{16} + \dots + 3u - 1 \end{pmatrix}$$

(ii) **Obstruction class** = -1

$$(iii) \text{ Cusp Shapes} = -4u^{17} + 4u^{16} - 36u^{15} + 28u^{14} - 124u^{13} + 72u^{12} - 196u^{11} + 72u^{10} - 120u^9 + 8u^7 - 36u^6 + 8u^5 - 4u^4 - 16u^3 + 8u - 18$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_2, c_6	$u^{18} + u^{17} + \cdots - u - 1$
c_3	$u^{18} - u^{17} + \cdots - 13u - 5$
c_4, c_8, c_9	$u^{18} + u^{17} + \cdots - 3u - 1$
c_5, c_7	$u^{18} - 3u^{17} + \cdots - 3u + 3$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_6	$y^{18} - 15y^{17} + \cdots - 7y + 1$
c_3	$y^{18} + 5y^{17} + \cdots - 39y + 25$
c_4, c_8, c_9	$y^{18} + 17y^{17} + \cdots - 7y + 1$
c_5, c_7	$y^{18} + 13y^{17} + \cdots - 75y + 9$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.215059 + 1.214380I$	$-2.15328 + 3.22673I$	$-11.05526 - 3.62956I$
$u = -0.215059 - 1.214380I$	$-2.15328 - 3.22673I$	$-11.05526 + 3.62956I$
$u = 0.678984 + 0.355286I$	$-1.88880 - 5.71427I$	$-11.06596 + 6.05983I$
$u = 0.678984 - 0.355286I$	$-1.88880 + 5.71427I$	$-11.06596 - 6.05983I$
$u = -0.590027 + 0.406016I$	$2.42619 + 1.88569I$	$-5.68331 - 3.99357I$
$u = -0.590027 - 0.406016I$	$2.42619 - 1.88569I$	$-5.68331 + 3.99357I$
$u = 0.482433 + 0.528989I$	$-1.12877 + 1.78695I$	$-9.23943 + 0.02251I$
$u = 0.482433 - 0.528989I$	$-1.12877 - 1.78695I$	$-9.23943 - 0.02251I$
$u = 0.076050 + 1.298790I$	$3.35362 - 1.57187I$	$-5.80878 + 4.22070I$
$u = 0.076050 - 1.298790I$	$3.35362 + 1.57187I$	$-5.80878 - 4.22070I$
$u = -0.663049$	-5.83256	-16.3720
$u = 0.17132 + 1.45278I$	$5.14514 - 0.55896I$	$-5.51114 - 0.25710I$
$u = 0.17132 - 1.45278I$	$5.14514 + 0.55896I$	$-5.51114 + 0.25710I$
$u = 0.25789 + 1.44398I$	$3.89024 - 9.13509I$	$-6.98695 + 5.86478I$
$u = 0.25789 - 1.44398I$	$3.89024 + 9.13509I$	$-6.98695 - 5.86478I$
$u = -0.22144 + 1.45044I$	$8.38729 + 4.87394I$	$-2.47320 - 3.60136I$
$u = -0.22144 - 1.45044I$	$8.38729 - 4.87394I$	$-2.47320 + 3.60136I$
$u = 0.382766$	-0.621918	-15.9800

II. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1, c_2, c_6	$u^{18} + u^{17} + \cdots - u - 1$
c_3	$u^{18} - u^{17} + \cdots - 13u - 5$
c_4, c_8, c_9	$u^{18} + u^{17} + \cdots - 3u - 1$
c_5, c_7	$u^{18} - 3u^{17} + \cdots - 3u + 3$

III. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_6	$y^{18} - 15y^{17} + \cdots - 7y + 1$
c_3	$y^{18} + 5y^{17} + \cdots - 39y + 25$
c_4, c_8, c_9	$y^{18} + 17y^{17} + \cdots - 7y + 1$
c_5, c_7	$y^{18} + 13y^{17} + \cdots - 75y + 9$