


12a₀₀₃₆ (K12a₀₀₃₆)



Ideals for irreducible components² of X_{par}

$$I_1^u = \langle -3.71327 \times 10^{48} u^{35} + 4.04621 \times 10^{50} u^{34} + \cdots + 1.08619 \times 10^{53} d - 2.28969 \times 10^{52}, \\ 5.92442 \times 10^{49} u^{35} - 1.37260 \times 10^{51} u^{34} + \cdots + 2.17237 \times 10^{53} c - 1.61328 \times 10^{53}, \\ 1.56429 \times 10^{50} u^{35} - 1.90035 \times 10^{51} u^{34} + \cdots + 1.08619 \times 10^{53} b + 1.05897 \times 10^{53}, \\ - 1.77635 \times 10^{50} u^{35} + 8.48997 \times 10^{50} u^{34} + \cdots + 1.08619 \times 10^{53} a - 1.56178 \times 10^{53}, \\ u^{36} - 3u^{35} + \cdots - 64u + 32 \rangle$$

$$I_2^u = \langle 1.13466 \times 10^{15} au^{27} - 3.06796 \times 10^{15} u^{27} + \dots + 2.79964 \times 10^{15} a - 1.43321 \times 10^{16}, \\ - 6.13592 \times 10^{15} au^{27} - 7.57070 \times 10^{15} u^{27} + \dots - 2.86643 \times 10^{16} a - 2.56902 \times 10^{16}, b + 1, \\ 2.13254 \times 10^{15} au^{27} + 1.47729 \times 10^{16} u^{27} + \dots + 2.93914 \times 10^{16} a + 7.21477 \times 10^{16}, u^{28} + u^{27} + \dots + 8u + 4 \rangle$$

$$I_1^v = \langle a, d, c-1, b+1, v^2 + v + 1 \rangle$$

$$I_2^v = \langle a, d - 1, c + a, b + 1, v^2 + v + 1 \rangle$$

$$I_3^v = \langle a, d-1, c+a+1, b+1, v+1 \rangle$$

$$I_4^{\prime }=\langle c,\;d-1,\;cb-a-1,\;v^2ba+v^2c+v^2b+2v^2a+c v+av+2v^2+c+v,\;b^2v^2+2v^2b+bv+v^2+v+1\rangle$$

* 5 irreducible components of $\dim_{\mathbb{C}} = 0$, with total 97 representations.

* 1 irreducible components of $\dim_{\mathbb{C}} = 1$

¹The image of knot diagram is generated by the software “**Draw programme**” developed by Andrew Bartholomew(<http://www.layer8.co.uk/maths/draw/index.htm#Running-draw>), where we modified some parts for our purpose(<https://github.com/CATsTAILs/LinksPainter>).

²All coefficients of polynomials are rational numbers. But the coefficients are sometimes approximated in decimal forms when there is not enough margin.

$$\text{I. } I_1^u = \langle -3.71 \times 10^{48}u^{35} + 4.05 \times 10^{50}u^{34} + \dots + 1.09 \times 10^{53}d - 2.29 \times 10^{52}, 5.92 \times 10^{49}u^{35} - 1.37 \times 10^{51}u^{34} + \dots + 2.17 \times 10^{53}c - 1.61 \times 10^{53}, 1.56 \times 10^{50}u^{35} - 1.90 \times 10^{51}u^{34} + \dots + 1.09 \times 10^{53}b + 1.06 \times 10^{53}, -1.78 \times 10^{50}u^{35} + 8.49 \times 10^{50}u^{34} + \dots + 1.09 \times 10^{53}a - 1.56 \times 10^{53}, u^{36} - 3u^{35} + \dots - 64u + 32 \rangle$$

(i) Arc colorings

$$a_4 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix}$$

$$a_7 = \begin{pmatrix} 0.00163541u^{35} - 0.00781632u^{34} + \dots - 1.02250u + 1.43786 \\ -0.00144017u^{35} + 0.0174956u^{34} + \dots + 2.23968u - 0.974944 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -0.000272717u^{35} + 0.00631846u^{34} + \dots + 0.563935u + 0.742637 \\ 0.0000341864u^{35} - 0.00372516u^{34} + \dots - 0.444645u + 0.210801 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -0.0000432971u^{35} + 0.0122726u^{34} + \dots + 1.33646u + 0.416353 \\ 0.00188305u^{35} - 0.0189570u^{34} + \dots - 2.03361u + 0.857265 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 0.000904113u^{35} - 0.00884304u^{34} + \dots - 1.47566u + 1.66218 \\ -0.000947410u^{35} + 0.0211156u^{34} + \dots + 2.81213u - 1.24583 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} 0.00194611u^{35} - 0.00144413u^{34} + \dots + 2.53686u - 0.189589 \\ 0.0137165u^{35} - 0.0449480u^{34} + \dots - 0.634013u + 0.309094 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 0.00349030u^{35} - 0.000257565u^{34} + \dots + 3.44348u - 0.336815 \\ 0.0131839u^{35} - 0.0436398u^{34} + \dots - 1.21762u + 0.270108 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} -0.000238531u^{35} + 0.00259330u^{34} + \dots + 0.119290u + 0.953438 \\ 0.0000341864u^{35} - 0.00372516u^{34} + \dots - 0.444645u + 0.210801 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} -0.000238531u^{35} + 0.00259330u^{34} + \dots + 0.119290u + 0.953438 \\ 0.000492763u^{35} + 0.00362002u^{34} + \dots + 0.572452u - 0.270888 \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes = $0.0805025u^{35} - 0.242112u^{34} + \dots + 5.72407u - 9.52193$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$u^{36} + 17u^{35} + \cdots - 120u + 16$
c_2, c_5	$u^{36} + u^{35} + \cdots + 16u + 4$
c_3	$u^{36} - u^{35} + \cdots - 104u + 1252$
c_4, c_9	$u^{36} + 3u^{35} + \cdots + 64u + 32$
c_6, c_7, c_8 c_{10}, c_{11}, c_{12}	$u^{36} - 5u^{35} + \cdots + 3u - 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$y^{36} + 5y^{35} + \cdots - 27936y + 256$
c_2, c_5	$y^{36} + 17y^{35} + \cdots - 120y + 16$
c_3	$y^{36} - 7y^{35} + \cdots - 22103608y + 1567504$
c_4, c_9	$y^{36} - 15y^{35} + \cdots - 1024y + 1024$
c_6, c_7, c_8 c_{10}, c_{11}, c_{12}	$y^{36} - 41y^{35} + \cdots - 21y + 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.956534 + 0.246505I$ $a = -0.0142428 + 0.0350628I$ $b = 0.595799 - 0.651291I$ $c = 0.141093 - 0.856670I$ $d = 0.446161 + 0.609262I$	$-2.56616 - 0.65266I$	$-13.12638 + 3.31133I$
$u = 0.956534 - 0.246505I$ $a = -0.0142428 - 0.0350628I$ $b = 0.595799 + 0.651291I$ $c = 0.141093 + 0.856670I$ $d = 0.446161 - 0.609262I$	$-2.56616 + 0.65266I$	$-13.12638 - 3.31133I$
$u = -0.935949 + 0.527292I$ $a = -0.118634 + 0.273971I$ $b = 0.711040 + 0.301202I$ $c = 0.364553 + 1.079240I$ $d = 0.236726 - 0.726179I$	$1.36964 + 3.10356I$	$-5.16268 - 4.71165I$
$u = -0.935949 - 0.527292I$ $a = -0.118634 - 0.273971I$ $b = 0.711040 - 0.301202I$ $c = 0.364553 - 1.079240I$ $d = 0.236726 + 0.726179I$	$1.36964 - 3.10356I$	$-5.16268 + 4.71165I$
$u = -0.651308 + 0.620650I$ $a = 0.088381 + 0.613358I$ $b = 0.525117 + 0.024485I$ $c = 0.711979 + 1.024680I$ $d = 0.018859 - 0.596676I$	$2.25394 + 1.43184I$	$-2.05632 - 3.52848I$
$u = -0.651308 - 0.620650I$ $a = 0.088381 - 0.613358I$ $b = 0.525117 - 0.024485I$ $c = 0.711979 - 1.024680I$ $d = 0.018859 + 0.596676I$	$2.25394 - 1.43184I$	$-2.05632 + 3.52848I$

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.134519 + 0.831817I$ $a = -2.49461 + 0.46036I$ $b = 5.38486 - 0.94232I$ $c = -0.800905 - 0.073407I$ $d = 1.245840 + 0.061269I$	$-4.07214 - 2.53804I$	$-12.12047 + 3.16226I$
$u = -0.134519 - 0.831817I$ $a = -2.49461 - 0.46036I$ $b = 5.38486 + 0.94232I$ $c = -0.800905 + 0.073407I$ $d = 1.245840 - 0.061269I$	$-4.07214 + 2.53804I$	$-12.12047 - 3.16226I$
$u = 0.488093 + 0.675929I$ $a = 0.219986 - 0.913847I$ $b = 0.439268 + 0.155961I$ $c = 0.951048 - 1.034360I$ $d = -0.119169 + 0.531960I$	$0.92725 + 3.15352I$	$-4.07893 - 3.45921I$
$u = 0.488093 - 0.675929I$ $a = 0.219986 + 0.913847I$ $b = 0.439268 - 0.155961I$ $c = 0.951048 + 1.034360I$ $d = -0.119169 - 0.531960I$	$0.92725 - 3.15352I$	$-4.07893 + 3.45921I$
$u = 1.066500 + 0.529753I$ $a = -0.230642 - 0.205554I$ $b = 0.835462 - 0.357382I$ $c = 0.257325 - 1.157650I$ $d = 0.296959 + 0.811036I$	$-0.85359 - 7.85577I$	$-9.34096 + 8.43490I$
$u = 1.066500 - 0.529753I$ $a = -0.230642 + 0.205554I$ $b = 0.835462 + 0.357382I$ $c = 0.257325 + 1.157650I$ $d = 0.296959 - 0.811036I$	$-0.85359 + 7.85577I$	$-9.34096 - 8.43490I$

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.471804 + 1.201650I$ $a = -1.55056 + 0.44817I$ $b = 3.43387 - 0.95934I$ $c = -1.009840 - 0.265513I$ $d = 1.41953 + 0.22707I$	$-6.92260 - 4.39379I$	$-12.55418 + 2.40542I$
$u = -0.471804 - 1.201650I$ $a = -1.55056 - 0.44817I$ $b = 3.43387 + 0.95934I$ $c = -1.009840 + 0.265513I$ $d = 1.41953 - 0.22707I$	$-6.92260 + 4.39379I$	$-12.55418 - 2.40542I$
$u = 1.232210 + 0.408112I$ $a = 1.16290 - 2.88555I$ $b = 3.50484 - 0.82980I$ $c = -0.478569 + 1.058540I$ $d = -1.43441 - 0.19589I$	$-8.11992 - 1.67660I$	$-15.7568 + 0.9806I$
$u = 1.232210 - 0.408112I$ $a = 1.16290 + 2.88555I$ $b = 3.50484 + 0.82980I$ $c = -0.478569 - 1.058540I$ $d = -1.43441 + 0.19589I$	$-8.11992 + 1.67660I$	$-15.7568 - 0.9806I$
$u = -1.223950 + 0.516090I$ $a = 0.35261 + 3.02077I$ $b = 3.33397 + 0.96426I$ $c = -0.310513 - 1.254720I$ $d = -1.43061 + 0.24878I$	$-7.33125 + 7.52384I$	$-14.0983 - 6.1679I$
$u = -1.223950 - 0.516090I$ $a = 0.35261 - 3.02077I$ $b = 3.33397 - 0.96426I$ $c = -0.310513 + 1.254720I$ $d = -1.43061 - 0.24878I$	$-7.33125 - 7.52384I$	$-14.0983 + 6.1679I$

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.666134 + 0.068993I$		
$a = 0.780189 + 0.682100I$		
$b = -0.90161 - 1.52116I$	$-2.77509 - 2.72721I$	$-16.6085 + 5.9652I$
$c = -0.128853 - 0.268419I$		
$d = 0.706776 + 0.200522I$		
$u = -0.666134 - 0.068993I$		
$a = 0.780189 - 0.682100I$		
$b = -0.90161 + 1.52116I$	$-2.77509 + 2.72721I$	$-16.6085 - 5.9652I$
$c = -0.128853 + 0.268419I$		
$d = 0.706776 - 0.200522I$		
$u = 0.318666 + 1.335800I$		
$a = -1.57860 - 0.26349I$		
$b = 3.47760 + 0.56643I$	$-11.42670 + 0.95860I$	$-17.6650 + 0.2050I$
$c = -1.083360 + 0.176981I$		
$d = 1.48437 - 0.15238I$		
$u = 0.318666 - 1.335800I$		
$a = -1.57860 + 0.26349I$		
$b = 3.47760 - 0.56643I$	$-11.42670 - 0.95860I$	$-17.6650 - 0.2050I$
$c = -1.083360 - 0.176981I$		
$d = 1.48437 + 0.15238I$		
$u = 0.568640 + 1.285090I$		
$a = -1.44127 - 0.41824I$		
$b = 3.20069 + 0.90828I$	$-9.63713 + 9.42250I$	$-15.4804 - 5.9490I$
$c = -1.059710 + 0.318993I$		
$d = 1.46105 - 0.27427I$		
$u = 0.568640 - 1.285090I$		
$a = -1.44127 + 0.41824I$		
$b = 3.20069 - 0.90828I$	$-9.63713 - 9.42250I$	$-15.4804 + 5.9490I$
$c = -1.059710 - 0.318993I$		
$d = 1.46105 + 0.27427I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.579877$		
$a = 0.700882$		
$b = -0.382079$	-0.811618	-12.0290
$c = 0.313536$		
$d = 0.400496$		
$u = -1.27702 + 0.74720I$		
$a = -0.79571 + 2.33781I$		
$b = 2.94232 + 0.99996I$	-9.5381 + 11.3478I	-13.0594 - 5.6672I
$c = 0.11920 - 1.42161I$		
$d = -1.45945 + 0.36293I$		
$u = -1.27702 - 0.74720I$		
$a = -0.79571 - 2.33781I$		
$b = 2.94232 - 0.99996I$	-9.5381 - 11.3478I	-13.0594 + 5.6672I
$c = 0.11920 + 1.42161I$		
$d = -1.45945 - 0.36293I$		
$u = 1.51924$		
$a = 1.38331$		
$b = 3.36154$	-14.8609	-15.8600
$c = -0.179172$		
$d = -1.57128$		
$u = 1.29117 + 0.81558I$		
$a = -0.98989 - 2.12524I$		
$b = 2.84808 - 0.99055I$	-12.0349 - 16.8809I	-15.3736 + 9.2575I
$c = 0.22492 + 1.45088I$		
$d = -1.46798 - 0.39682I$		
$u = 1.29117 - 0.81558I$		
$a = -0.98989 + 2.12524I$		
$b = 2.84808 + 0.99055I$	-12.0349 + 16.8809I	-15.3736 - 9.2575I
$c = 0.22492 - 1.45088I$		
$d = -1.46798 + 0.39682I$		

Solutions to I_1^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.111921 + 0.451208I$ $a = 1.46514 - 0.45120I$ $b = -0.0451061 + 0.0894307I$ $c = 1.40357 - 0.27941I$ $d = -0.190856 + 0.111820I$	$-0.30692 - 1.79670I$	$-2.13838 + 3.34715I$
$u = 0.111921 - 0.451208I$ $a = 1.46514 + 0.45120I$ $b = -0.0451061 - 0.0894307I$ $c = 1.40357 + 0.27941I$ $d = -0.190856 - 0.111820I$	$-0.30692 + 1.79670I$	$-2.13838 - 3.34715I$
$u = 1.38860 + 0.67758I$ $a = -0.35175 - 2.03094I$ $b = 2.99578 - 0.83109I$ $c = 0.126360 + 1.208340I$ $d = -1.51341 - 0.32643I$	$-15.0030 - 8.0885I$	$-18.3300 + 3.8436I$
$u = 1.38860 - 0.67758I$ $a = -0.35175 + 2.03094I$ $b = 2.99578 + 0.83109I$ $c = 0.126360 - 1.208340I$ $d = -1.51341 + 0.32643I$	$-15.0030 + 8.0885I$	$-18.3300 - 3.8436I$
$u = -1.61122 + 0.12770I$ $a = 0.954593 + 0.434273I$ $b = 3.22829 + 0.14916I$ $c = 0.004513 - 0.218310I$ $d = -1.61501 + 0.06033I$	$-18.8051 + 4.7571I$	$-19.1049 - 3.2273I$
$u = -1.61122 - 0.12770I$ $a = 0.954593 - 0.434273I$ $b = 3.22829 - 0.14916I$ $c = 0.004513 + 0.218310I$ $d = -1.61501 - 0.06033I$	$-18.8051 - 4.7571I$	$-19.1049 + 3.2273I$

II.

$$I_2^u = \langle 1.13 \times 10^{15} au^{27} - 3.07 \times 10^{15} u^{27} + \dots + 2.80 \times 10^{15} a - 1.43 \times 10^{16}, -6.14 \times 10^{15} au^{27} - 7.57 \times 10^{15} u^{27} + \dots - 2.87 \times 10^{16} a - 2.57 \times 10^{16}, b + 1, 2.13 \times 10^{15} au^{27} + 1.48 \times 10^{16} u^{27} + \dots + 2.94 \times 10^{16} a + 7.21 \times 10^{16}, u^{28} + u^{27} + \dots + 8u + 4 \rangle$$

(i) Arc colorings

$$a_4 = \begin{pmatrix} 0 \\ u \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 1 \\ u^2 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -u \\ -u^3 + u \end{pmatrix}$$

$$a_7 = \begin{pmatrix} a \\ -1 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 2.97137au^{27} + 3.66617u^{27} + \dots + 13.8809a + 12.4407 \\ -1.09893au^{27} + 2.97137u^{27} + \dots - 2.71150a + 13.8809 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -1.87244u^{27} - 0.749749u^{26} + \dots - 7.03105u - 11.1694 \\ 1.09893u^{27} - 0.185171u^{26} + \dots + 5.48839u + 2.71150 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -0.333289u^{27} - 0.505230u^{26} + \dots - 0.0509032u - 3.96717 \\ -1.53915u^{27} - 0.244519u^{26} + \dots - 6.98014u - 7.20224 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -3.54326u^{27} - 0.280326u^{26} + \dots - 16.6119u - 14.7552 \\ 0.187465u^{27} - 0.126714u^{26} + \dots + 1.72320u + 0.0904714 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -2.49289u^{27} - 0.296873u^{26} + \dots - 11.3968u - 10.5619 \\ 0.0560178u^{27} - 0.0365360u^{26} + \dots + 0.841877u + 0.164880 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 1.87244au^{27} + 6.63754u^{27} + \dots + 11.1694a + 26.3216 \\ -1.09893au^{27} + 2.97137u^{27} + \dots - 2.71150a + 13.8809 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1.87244au^{27} + 6.63754u^{27} + \dots + 11.1694a + 26.3216 \\ 1.53915au^{27} + 1.87244u^{27} + \dots + 7.20224a + 9.16941 \end{pmatrix}$$

(ii) Obstruction class = -1

(iii) Cusp Shapes

$$= \frac{37274533074785}{129063433504426}u^{27} + \frac{53621307092165}{129063433504426}u^{26} + \dots + \frac{660629047616669}{129063433504426}u - \frac{645502044434670}{64531716752213}$$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1	$(u^{28} + 14u^{27} + \cdots + 2u + 1)^2$
c_2, c_5	$(u^{28} + 2u^{27} + \cdots + 2u + 1)^2$
c_3	$(u^{28} - 2u^{27} + \cdots - 22u + 17)^2$
c_4, c_9	$(u^{28} - u^{27} + \cdots - 8u + 4)^2$
c_6, c_7, c_8 c_{10}, c_{11}, c_{12}	$u^{56} - 3u^{55} + \cdots + 72u + 16$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1	$(y^{28} + 2y^{27} + \cdots + 14y + 1)^2$
c_2, c_5	$(y^{28} + 14y^{27} + \cdots + 2y + 1)^2$
c_3	$(y^{28} - 10y^{27} + \cdots - 246y + 289)^2$
c_4, c_9	$(y^{28} - 15y^{27} + \cdots - 88y + 16)^2$
c_6, c_7, c_8 c_{10}, c_{11}, c_{12}	$y^{56} - 43y^{55} + \cdots + 736y + 256$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.910131 + 0.395689I$ $a = -1.02788 + 1.10315I$ $b = -1.00000$ $c = -1.20522 + 1.86019I$ $d = -1.276980 - 0.186733I$	$-1.72215 - 4.24816I$	$-10.11355 + 6.97904I$
$u = 0.910131 + 0.395689I$ $a = -0.01361 + 2.44534I$ $b = -1.00000$ $c = 0.300677 - 0.944242I$ $d = 0.313357 + 0.643423I$	$-1.72215 - 4.24816I$	$-10.11355 + 6.97904I$
$u = 0.910131 - 0.395689I$ $a = -1.02788 - 1.10315I$ $b = -1.00000$ $c = -1.20522 - 1.86019I$ $d = -1.276980 + 0.186733I$	$-1.72215 + 4.24816I$	$-10.11355 - 6.97904I$
$u = 0.910131 - 0.395689I$ $a = -0.01361 - 2.44534I$ $b = -1.00000$ $c = 0.300677 + 0.944242I$ $d = 0.313357 - 0.643423I$	$-1.72215 + 4.24816I$	$-10.11355 - 6.97904I$
$u = -0.017123 + 0.961380I$ $a = 0.177063 - 0.402321I$ $b = -1.00000$ $c = -0.873117 - 0.009489I$ $d = 1.306640 + 0.007988I$	$-4.61196 + 1.34593I$	$-13.91932 - 0.66126I$
$u = -0.017123 + 0.961380I$ $a = 1.58097 + 1.51341I$ $b = -1.00000$ $c = 1.50836 + 1.81823I$ $d = -0.638546 - 0.533764I$	$-4.61196 + 1.34593I$	$-13.91932 - 0.66126I$

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.017123 - 0.961380I$		
$a = 0.177063 + 0.402321I$		
$b = -1.00000$	$-4.61196 - 1.34593I$	$-13.91932 + 0.66126I$
$c = -0.873117 + 0.009489I$		
$d = 1.306640 - 0.007988I$		
$u = -0.017123 - 0.961380I$		
$a = 1.58097 - 1.51341I$		
$b = -1.00000$	$-4.61196 - 1.34593I$	$-13.91932 + 0.66126I$
$c = 1.50836 - 1.81823I$		
$d = -0.638546 + 0.533764I$		
$u = -0.907099 + 0.252760I$		
$a = 0.971495 + 0.463170I$		
$b = -1.00000$	$-2.63794 + 3.28147I$	$-13.2327 - 4.9939I$
$c = 0.186210 + 0.817511I$		
$d = 0.419544 - 0.572998I$		
$u = -0.907099 + 0.252760I$		
$a = 0.704043 - 0.365524I$		
$b = -1.00000$	$-2.63794 + 3.28147I$	$-13.2327 - 4.9939I$
$c = -0.351737 - 0.579401I$		
$d = 0.859190 + 0.455166I$		
$u = -0.907099 - 0.252760I$		
$a = 0.971495 - 0.463170I$		
$b = -1.00000$	$-2.63794 - 3.28147I$	$-13.2327 + 4.9939I$
$c = 0.186210 - 0.817511I$		
$d = 0.419544 + 0.572998I$		
$u = -0.907099 - 0.252760I$		
$a = 0.704043 + 0.365524I$		
$b = -1.00000$	$-2.63794 - 3.28147I$	$-13.2327 + 4.9939I$
$c = -0.351737 + 0.579401I$		
$d = 0.859190 - 0.455166I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.387411 + 0.832689I$		
$a = 0.365520 + 0.349527I$		
$b = -1.00000$	$-1.43770 + 1.40144I$	$-7.30053 - 1.74630I$
$c = -0.795695 + 0.216194I$		
$d = 1.239370 - 0.180444I$		
$u = 0.387411 + 0.832689I$		
$a = 1.42688 - 0.89203I$		
$b = -1.00000$	$-1.43770 + 1.40144I$	$-7.30053 - 1.74630I$
$c = 1.11790 - 1.26235I$		
$d = -0.279611 + 0.583432I$		
$u = 0.387411 - 0.832689I$		
$a = 0.365520 - 0.349527I$		
$b = -1.00000$	$-1.43770 - 1.40144I$	$-7.30053 + 1.74630I$
$c = -0.795695 - 0.216194I$		
$d = 1.239370 + 0.180444I$		
$u = 0.387411 - 0.832689I$		
$a = 1.42688 + 0.89203I$		
$b = -1.00000$	$-1.43770 - 1.40144I$	$-7.30053 + 1.74630I$
$c = 1.11790 + 1.26235I$		
$d = -0.279611 - 0.583432I$		
$u = -0.387502 + 1.047530I$		
$a = 0.340954 - 0.431763I$		
$b = -1.00000$	$-3.70255 - 5.75423I$	$-11.89302 + 5.96655I$
$c = -0.920428 - 0.218491I$		
$d = 1.344350 + 0.185032I$		
$u = -0.387502 + 1.047530I$		
$a = 1.27979 + 1.06278I$		
$b = -1.00000$	$-3.70255 - 5.75423I$	$-11.89302 + 5.96655I$
$c = 1.05142 + 1.53293I$		
$d = -0.382179 - 0.735842I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.387502 - 1.047530I$		
$a = 0.340954 + 0.431763I$		
$b = -1.00000$	$-3.70255 + 5.75423I$	$-11.89302 - 5.96655I$
$c = -0.920428 + 0.218491I$		
$d = 1.344350 - 0.185032I$		
$u = -0.387502 - 1.047530I$		
$a = 1.27979 - 1.06278I$		
$b = -1.00000$	$-3.70255 + 5.75423I$	$-11.89302 - 5.96655I$
$c = 1.05142 - 1.53293I$		
$d = -0.382179 + 0.735842I$		
$u = -0.802767 + 0.244916I$		
$a = -2.05210 - 1.27314I$		
$b = -1.00000$	$-2.32218 - 0.90628I$	$-12.59768 - 1.67094I$
$c = -2.29143 - 1.82037I$		
$d = -1.229860 + 0.111546I$		
$u = -0.802767 + 0.244916I$		
$a = -0.72932 - 2.54382I$		
$b = -1.00000$	$-2.32218 - 0.90628I$	$-12.59768 - 1.67094I$
$c = 0.273098 + 0.711267I$		
$d = 0.374376 - 0.483492I$		
$u = -0.802767 - 0.244916I$		
$a = -2.05210 + 1.27314I$		
$b = -1.00000$	$-2.32218 + 0.90628I$	$-12.59768 + 1.67094I$
$c = -2.29143 + 1.82037I$		
$d = -1.229860 - 0.111546I$		
$u = -0.802767 - 0.244916I$		
$a = -0.72932 + 2.54382I$		
$b = -1.00000$	$-2.32218 + 0.90628I$	$-12.59768 + 1.67094I$
$c = 0.273098 - 0.711267I$		
$d = 0.374376 + 0.483492I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.147340 + 0.340892I$ $a = -0.224291 + 1.144170I$ $b = -1.00000$ $c = -0.471254 - 0.771751I$ $d = 0.938913 + 0.621780I$	$-5.81300 + 1.47542I$	$-13.29345 - 0.59666I$
$u = -1.147340 + 0.340892I$ $a = -1.07497 - 1.33487I$ $b = -1.00000$ $c = -0.776477 - 1.062760I$ $d = -1.393270 + 0.163069I$	$-5.81300 + 1.47542I$	$-13.29345 - 0.59666I$
$u = -1.147340 - 0.340892I$ $a = -0.224291 - 1.144170I$ $b = -1.00000$ $c = -0.471254 + 0.771751I$ $d = 0.938913 - 0.621780I$	$-5.81300 - 1.47542I$	$-13.29345 + 0.59666I$
$u = -1.147340 - 0.340892I$ $a = -1.07497 + 1.33487I$ $b = -1.00000$ $c = -0.776477 + 1.062760I$ $d = -1.393270 - 0.163069I$	$-5.81300 - 1.47542I$	$-13.29345 + 0.59666I$
$u = 0.611767 + 0.458091I$ $a = 1.235620 - 0.499323I$ $b = -1.00000$ $c = 0.669435 - 0.799308I$ $d = 0.099411 + 0.470368I$	$-0.887541 + 0.644142I$	$-7.64602 + 1.30683I$
$u = 0.611767 + 0.458091I$ $a = 0.552061 + 0.278825I$ $b = -1.00000$ $c = -0.542285 + 0.331792I$ $d = 1.027130 - 0.267751I$	$-0.887541 + 0.644142I$	$-7.64602 + 1.30683I$

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 0.611767 - 0.458091I$ $a = 1.235620 + 0.499323I$ $b = -1.00000$ $c = 0.669435 + 0.799308I$ $d = 0.099411 - 0.470368I$	$-0.887541 - 0.644142I$	$-7.64602 - 1.30683I$
$u = 0.611767 - 0.458091I$ $a = 0.552061 - 0.278825I$ $b = -1.00000$ $c = -0.542285 - 0.331792I$ $d = 1.027130 + 0.267751I$	$-0.887541 - 0.644142I$	$-7.64602 - 1.30683I$
$u = 1.175470 + 0.589984I$ $a = -0.299231 + 0.442331I$ $b = -1.00000$ $c = -0.23958 + 1.44120I$ $d = -1.40691 - 0.28559I$	$-3.88965 - 6.77427I$	$-10.22594 + 4.95962I$
$u = 1.175470 + 0.589984I$ $a = 0.42258 + 1.93124I$ $b = -1.00000$ $c = 0.209555 - 1.262070I$ $d = 0.307742 + 0.904345I$	$-3.88965 - 6.77427I$	$-10.22594 + 4.95962I$
$u = 1.175470 - 0.589984I$ $a = -0.299231 - 0.442331I$ $b = -1.00000$ $c = -0.23958 - 1.44120I$ $d = -1.40691 + 0.28559I$	$-3.88965 + 6.77427I$	$-10.22594 - 4.95962I$
$u = 1.175470 - 0.589984I$ $a = 0.42258 - 1.93124I$ $b = -1.00000$ $c = 0.209555 + 1.262070I$ $d = 0.307742 - 0.904345I$	$-3.88965 + 6.77427I$	$-10.22594 - 4.95962I$

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.312590 + 0.177484I$ $a = -0.920187 + 0.871890I$ $b = -1.00000$ $c = -0.575440 + 0.451407I$ $d = -1.47301 - 0.08468I$	$-9.90219 + 2.08114I$	$-17.7960 - 2.7886I$
$u = 1.312590 + 0.177484I$ $a = -0.113586 - 1.304510I$ $b = -1.00000$ $c = -0.391157 + 0.943404I$ $d = 0.853841 - 0.754977I$	$-9.90219 + 2.08114I$	$-17.7960 - 2.7886I$
$u = 1.312590 - 0.177484I$ $a = -0.920187 - 0.871890I$ $b = -1.00000$ $c = -0.575440 - 0.451407I$ $d = -1.47301 + 0.08468I$	$-9.90219 - 2.08114I$	$-17.7960 + 2.7886I$
$u = 1.312590 - 0.177484I$ $a = -0.113586 + 1.304510I$ $b = -1.00000$ $c = -0.391157 - 0.943404I$ $d = 0.853841 + 0.754977I$	$-9.90219 - 2.08114I$	$-17.7960 + 2.7886I$
$u = 1.262900 + 0.460239I$ $a = -0.110303 - 1.099220I$ $b = -1.00000$ $c = -0.584732 + 0.832898I$ $d = 1.027590 - 0.683591I$	$-8.61088 - 6.23266I$	$-16.1498 + 4.3008I$
$u = 1.262900 + 0.460239I$ $a = -0.73910 + 1.34519I$ $b = -1.00000$ $c = -0.341352 + 1.102730I$ $d = -1.44949 - 0.22125I$	$-8.61088 - 6.23266I$	$-16.1498 + 4.3008I$

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = 1.262900 - 0.460239I$		
$a = -0.110303 + 1.099220I$		
$b = -1.00000$	$-8.61088 + 6.23266I$	$-16.1498 - 4.3008I$
$c = -0.584732 - 0.832898I$		
$d = 1.027590 + 0.683591I$		
$u = 1.262900 - 0.460239I$		
$a = -0.73910 - 1.34519I$		
$b = -1.00000$	$-8.61088 + 6.23266I$	$-16.1498 - 4.3008I$
$c = -0.341352 - 1.102730I$		
$d = -1.44949 + 0.22125I$		
$u = -1.280370 + 0.446560I$		
$a = -0.502612 - 0.171924I$		
$b = -1.00000$	$-8.68474 + 3.62399I$	$-16.2087 - 2.7619I$
$c = -0.332736 - 1.053240I$		
$d = -1.45797 + 0.21451I$		
$u = -1.280370 + 0.446560I$		
$a = 0.25189 - 1.78789I$		
$b = -1.00000$	$-8.68474 + 3.62399I$	$-16.2087 - 2.7619I$
$c = 0.059654 + 1.225020I$		
$d = 0.436936 - 0.910395I$		
$u = -1.280370 - 0.446560I$		
$a = -0.502612 + 0.171924I$		
$b = -1.00000$	$-8.68474 - 3.62399I$	$-16.2087 + 2.7619I$
$c = -0.332736 + 1.053240I$		
$d = -1.45797 - 0.21451I$		
$u = -1.280370 - 0.446560I$		
$a = 0.25189 + 1.78789I$		
$b = -1.00000$	$-8.68474 - 3.62399I$	$-16.2087 + 2.7619I$
$c = 0.059654 - 1.225020I$		
$d = 0.436936 + 0.910395I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -0.376924 + 0.508425I$		
$a = -0.304778 + 0.396413I$		
$b = -1.00000$	$-3.51302 + 1.43304I$	$-13.5823 - 4.9760I$
$c = -0.608002 - 0.187738I$		
$d = 1.085050 + 0.152783I$		
$u = -0.376924 + 0.508425I$		
$a = 0.96605 - 5.26986I$		
$b = -1.00000$	$-3.51302 + 1.43304I$	$-13.5823 - 4.9760I$
$c = 1.01842 - 5.30238I$		
$d = -0.999265 + 0.195824I$		
$u = -0.376924 - 0.508425I$		
$a = -0.304778 - 0.396413I$		
$b = -1.00000$	$-3.51302 - 1.43304I$	$-13.5823 + 4.9760I$
$c = -0.608002 + 0.187738I$		
$d = 1.085050 - 0.152783I$		
$u = -0.376924 - 0.508425I$		
$a = 0.96605 + 5.26986I$		
$b = -1.00000$	$-3.51302 - 1.43304I$	$-13.5823 + 4.9760I$
$c = 1.01842 + 5.30238I$		
$d = -0.999265 - 0.195824I$		
$u = -1.241130 + 0.661367I$		
$a = -0.160346 - 0.333380I$		
$b = -1.00000$	$-6.41692 + 11.95450I$	$-13.0412 - 8.3222I$
$c = -0.043949 - 1.401630I$		
$d = -1.44024 + 0.32064I$		
$u = -1.241130 + 0.661367I$		
$a = 0.49740 - 1.83602I$		
$b = -1.00000$	$-6.41692 + 11.95450I$	$-13.0412 - 8.3222I$
$c = 0.199842 + 1.344140I$		
$d = 0.293882 - 0.972896I$		

Solutions to I_2^u	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$u = -1.241130 - 0.661367I$		
$a = -0.160346 + 0.333380I$		
$b = -1.00000$	$-6.41692 - 11.95450I$	$-13.0412 + 8.3222I$
$c = -0.043949 + 1.401630I$		
$d = -1.44024 - 0.32064I$		
$u = -1.241130 - 0.661367I$		
$a = 0.49740 + 1.83602I$		
$b = -1.00000$	$-6.41692 - 11.95450I$	$-13.0412 + 8.3222I$
$c = 0.199842 - 1.344140I$		
$d = 0.293882 + 0.972896I$		

$$\text{III. } I_1^v = \langle a, d, c-1, b+1, v^2 + v + 1 \rangle$$

(i) **Arc colorings**

$$a_4 = \begin{pmatrix} v \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} v \\ 0 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -1 \\ -1 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} 0 \\ -v \end{pmatrix}$$

$$a_2 = \begin{pmatrix} 1 \\ -v \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

(ii) **Obstruction class = 1**

(iii) **Cusp Shapes = $-4v - 11$**

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_3, c_5	$u^2 - u + 1$
c_2	$u^2 + u + 1$
c_4, c_8, c_9 c_{10}, c_{11}	u^2
c_6, c_7	$(u - 1)^2$
c_{12}	$(u + 1)^2$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_3 c_5	$y^2 + y + 1$
c_4, c_8, c_9 c_{10}, c_{11}	y^2
c_6, c_7, c_{12}	$(y - 1)^2$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_1^v	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$v = -0.500000 + 0.866025I$		
$a = 0$		
$b = -1.00000$	$-1.64493 + 2.02988I$	$-9.00000 - 3.46410I$
$c = 1.00000$		
$d = 0$		
$v = -0.500000 - 0.866025I$		
$a = 0$		
$b = -1.00000$	$-1.64493 - 2.02988I$	$-9.00000 + 3.46410I$
$c = 1.00000$		
$d = 0$		

$$\text{IV. } I_2^v = \langle a, d-1, c+a, b+1, v^2+v+1 \rangle$$

(i) **Arc colorings**

$$a_4 = \begin{pmatrix} v \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} v \\ 0 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} v \\ -v \end{pmatrix}$$

$$a_2 = \begin{pmatrix} v+1 \\ -v \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$$

(ii) **Obstruction class** = 1

(iii) **Cusp Shapes** = $-4v - 11$

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_3, c_5	$u^2 - u + 1$
c_2	$u^2 + u + 1$
c_4, c_6, c_7 c_9, c_{12}	u^2
c_8	$(u - 1)^2$
c_{10}, c_{11}	$(u + 1)^2$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_3 c_5	$y^2 + y + 1$
c_4, c_6, c_7 c_9, c_{12}	y^2
c_8, c_{10}, c_{11}	$(y - 1)^2$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_2^v	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$v = -0.500000 + 0.866025I$		
$a = 0$		
$b = -1.00000$	$-1.64493 + 2.02988I$	$-9.00000 - 3.46410I$
$c = 0$		
$d = 1.00000$		
$v = -0.500000 - 0.866025I$		
$a = 0$		
$b = -1.00000$	$-1.64493 - 2.02988I$	$-9.00000 + 3.46410I$
$c = 0$		
$d = 1.00000$		

$$\mathbf{V}. \quad I_3^v = \langle a, d-1, c+a+1, b+1, v+1 \rangle$$

(i) Arc colorings

$$a_4 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

(ii) Obstruction class = 1

(iii) Cusp Shapes = -12

(iv) u-Polynomials at the component

Crossings	u-Polynomials at each crossing
c_1, c_2, c_3 c_4, c_5, c_9	u
c_6, c_7, c_{10} c_{11}	$u + 1$
c_8, c_{12}	$u - 1$

(v) Riley Polynomials at the component

Crossings	Riley Polynomials at each crossing
c_1, c_2, c_3 c_4, c_5, c_9	y
c_6, c_7, c_8 c_{10}, c_{11}, c_{12}	$y - 1$

(vi) Complex Volumes and Cusp Shapes

Solutions to I_3^v	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$v = -1.00000$		
$a = 0$		
$b = -1.00000$	-3.28987	-12.0000
$c = -1.00000$		
$d = 1.00000$		

VI.

$$I_4^v = \langle c, d - 1, cb - a - 1, v^2ba + v^2c + \cdots + c + v, b^2v^2 + 2v^2b + \cdots + v + 1 \rangle$$

(i) **Arc colorings**

$$a_4 = \begin{pmatrix} v \\ 0 \end{pmatrix}$$

$$a_9 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_{10} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$a_5 = \begin{pmatrix} v \\ 0 \end{pmatrix}$$

$$a_7 = \begin{pmatrix} -1 \\ b \end{pmatrix}$$

$$a_{12} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$a_1 = \begin{pmatrix} -1 \\ b+1 \end{pmatrix}$$

$$a_6 = \begin{pmatrix} -1 \\ b+1 \end{pmatrix}$$

$$a_3 = \begin{pmatrix} bv + 2v \\ -b^2v - 2bv - v \end{pmatrix}$$

$$a_2 = \begin{pmatrix} -v^2b + bv - v^2 + v \\ -b^2v - 2bv - v \end{pmatrix}$$

$$a_{11} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$a_8 = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$$

(ii) **Obstruction class** = -1

(iii) **Cusp Shapes** = $b^3v + 3b^2v - bv + v^2 - 3v - 16$

(iv) **u-Polynomials at the component** : It cannot be defined for a positive dimension component.

(v) **Riley Polynomials at the component** : It cannot be defined for a positive dimension component.

(iv) Complex Volumes and Cusp Shapes

Solution to I_4^v	$\sqrt{-1}(\text{vol} + \sqrt{-1}CS)$	Cusp shape
$v = \dots$		
$a = \dots$		
$b = \dots$	$-3.28987 - 2.02988I$	$-14.5770 - 3.5248I$
$c = \dots$		
$d = \dots$		

VII. u-Polynomials

Crossings	u-Polynomials at each crossing
c_1	$u(u^2 - u + 1)^2(u^{28} + 14u^{27} + \dots + 2u + 1)^2 \cdot (u^{36} + 17u^{35} + \dots - 120u + 16)$
c_2	$u(u^2 + u + 1)^2(u^{28} + 2u^{27} + \dots + 2u + 1)^2(u^{36} + u^{35} + \dots + 16u + 4)$
c_3	$u(u^2 - u + 1)^2(u^{28} - 2u^{27} + \dots - 22u + 17)^2 \cdot (u^{36} - u^{35} + \dots - 104u + 1252)$
c_4, c_9	$u^5(u^{28} - u^{27} + \dots - 8u + 4)^2(u^{36} + 3u^{35} + \dots + 64u + 32)$
c_5	$u(u^2 - u + 1)^2(u^{28} + 2u^{27} + \dots + 2u + 1)^2(u^{36} + u^{35} + \dots + 16u + 4)$
c_6, c_7	$u^2(u - 1)^2(u + 1)(u^{36} - 5u^{35} + \dots + 3u - 1) \cdot (u^{56} - 3u^{55} + \dots + 72u + 16)$
c_8	$u^2(u - 1)^3(u^{36} - 5u^{35} + \dots + 3u - 1)(u^{56} - 3u^{55} + \dots + 72u + 16)$
c_{10}, c_{11}	$u^2(u + 1)^3(u^{36} - 5u^{35} + \dots + 3u - 1)(u^{56} - 3u^{55} + \dots + 72u + 16)$
c_{12}	$u^2(u - 1)(u + 1)^2(u^{36} - 5u^{35} + \dots + 3u - 1) \cdot (u^{56} - 3u^{55} + \dots + 72u + 16)$

VIII. Riley Polynomials

Crossings	Riley Polynomials at each crossing
c_1	$y(y^2 + y + 1)^2(y^{28} + 2y^{27} + \dots + 14y + 1)^2 \cdot (y^{36} + 5y^{35} + \dots - 27936y + 256)$
c_2, c_5	$y(y^2 + y + 1)^2(y^{28} + 14y^{27} + \dots + 2y + 1)^2 \cdot (y^{36} + 17y^{35} + \dots - 120y + 16)$
c_3	$y(y^2 + y + 1)^2(y^{28} - 10y^{27} + \dots - 246y + 289)^2 \cdot (y^{36} - 7y^{35} + \dots - 22103608y + 1567504)$
c_4, c_9	$y^5(y^{28} - 15y^{27} + \dots - 88y + 16)^2 \cdot (y^{36} - 15y^{35} + \dots - 1024y + 1024)$
c_6, c_7, c_8 c_{10}, c_{11}, c_{12}	$y^2(y - 1)^3(y^{36} - 41y^{35} + \dots - 21y + 1) \cdot (y^{56} - 43y^{55} + \dots + 736y + 256)$